University of Groningen

Peroxisome biogenesis and maintenance in yeast

Wroblewska, Justyna

DOI:
[10.33612/diss.113500905](https://doi.org/10.33612/diss.113500905)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Wroblewska, J. (2020). *Peroxisome biogenesis and maintenance in yeast*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.113500905>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Peroxisome biogenesis and maintenance in yeast

Justyna Paulina Wróblewska

**rijksuniversiteit
 groningen**

The work presented in this thesis was carried out in the research unit Molecular Cell Biology of the Groningen Biomolecular Sciences and Biotechnology Institute (GBB) of the University of Groningen, The Netherlands.

This project was supported financially by the Marie Curie Initial Training Networks (ITN) program PerFuMe (Grant Agreement Number 316723).

ISBN digital version: 978-94-034-2365-4

ISBN printed version: 978-94-034-2366-1

© 2020 Justyna Paulina Wróblewska, Groningen, The Netherlands

All rights reserved.

Cover design: Justyna Paulina Wróblewska

Layout and design: Jules Verkade, persoonlijkproefschrift.nl

Printing: Ridderprint BV | www.ridderprint.nl

university of
 groningen

Peroxisome biogenesis and maintenance in yeast

PhD thesis

to obtain the degree of PhD at the
 University of Groningen
 on the authority of the
 Rector Magnificus Prof. C. Wijmenga
 and in accordance with
 the decision by the College of Deans.

This thesis will be defended in public on

Friday 21 February 2020 at 11.00 hours

by

Justyna Paulina Wróblewska

born on 25 February 1987
 in Racibórz, Poland

Supervisor

Prof. I.J. van der Klei

Co-supervisor

Dr. D. Devos

Assessment Committee

Prof. M. Schrader

Prof. R.A.L. Bovenberg

Prof. J. Kok

To my loving mom and the world-changing friends I made over the course of my Ph.D.
Without your love and support, this book would not have been possible.

Justyna

THANK YOU!

MY PROMOTER: PROF. IDA VAN DER KLEI

MY CO-PROMOTER: DR. DAMIEN DEVOS

THE READING COMMITTEE THE PERFUME NETWORK

MY PARANYMPHS: RENATE JANSEN & RINSE DE BOER

RITIKA SINGH ♥ MY MOMMY ♥ MY FAMILY

JANNET NIJHUIS-KAMPEN NATASHA DANDA

ANN THOMAS ARMAN AKŞIT TERRY

ADAM KAWAŁEK MAŁGOSIA KRYGOWSKA

HUALA WU FEI WU ISCO XIN CHEN

CHRIS WILLIAMS SANJEEV KUMAR

ARJEN KRIKKEN KÈVIN KNOOPS

SHRISHTI DEVARAJAN JAN KIEL

SELVA MANIVANNAN ROMY AKKERMAN

MY STUDENTS: VERA, JOEY, BASTIAAN

INDIAN PARENTS & HAPPY ANITA KRAM

EINAT ZALCKVAR MAYA SCHULDINER

KASIA GAWRON SABINA TABACZAR

DAGMARA KOGUCIUK DAVID DONKOR

MARTA RYZNER ELIZA WARSZAWIK

ILONA JÓŹWIK ŞEYDA ŞEN ASIA SZWED

MORGAN LOVE EDYTA KASZUBSKA

BASIA MATUSEWICZ SANDRA KUSTOS

SYLVIA PUTZ FRANKY DAVID FARAGO

Table of contents

Aim and outline	11
Chapter 1 Introduction: peroxisome proliferation and dynamics	15
Chapter 2 <i>Saccharomyces cerevisiae</i> cells lacking Pex3 contain membrane vesicles that harbor a subset of peroxisomal membrane proteins	33
Chapter 3 Large-scale study of the origin of peroxisomal membrane vesicles in <i>Saccharomyces cerevisiae</i> <i>pex3 atg1</i> cells	65
Chapter 4 <i>Hansenula polymorpha</i> Vac8: a vacuolar membrane protein required for vacuole inheritance and nucleus-vacuole junction formation	93
Chapter 5 Peroxisome maintenance depends on <i>de novo</i> peroxisome formation in yeast mutants defective in peroxisome fission and inheritance	111
Summary	135
Samenvatting	142
Biography of the author	148

Peroxisomes belong to an important class of sub-cellular organelles present in essentially all eukaryotes. These highly dynamic organelles are involved in a wide range of functions which depend on the organism, cell type, developmental stage as well as internal and external cues.

PEX genes encode peroxins that are responsible for peroxisome formation. Mutations in those genes in human lead to severe disorders, often lethal at very early developmental stages. Because yeast *pex* mutants are viable, they provide an ideal system for studies of the molecular bases of peroxisome biogenesis.

Numerous studies addressing peroxisome biogenesis suggest two ways of peroxisome formation. The first one proposes growth and division of the pre-existing peroxisomes while the second one indicates that peroxisomes form *de novo* with an engagement of the endoplasmic reticulum (ER). Recent studies in yeast have led to the discovery of pre-peroxisomal vesicles (PPVs), which may represent early stages of peroxisomes in the *de novo* formation pathway. The aim of this thesis is to obtain further insights into this pathway.

Chapter 1 provides an overview of the current knowledge on peroxisome formation and inheritance in yeast.

In **Chapter 2** we show that peroxisomal membrane vesicles are present in an *S. cerevisiae pex3* mutant, as was previously demonstrated for *H. polymorpha pex3* cells. This finding counters the generally accepted view that cells lacking Pex3 are devoid of any peroxisomal membrane structures. At the vesicular structures a subset of peroxisomal membrane proteins (PMPs) (Pex14, Pex13, Pex17 and Pex5) assemble into a complex similar to the PTS1 protein translocation pore of WT yeast cells. Using a combination of microscopy and biochemical approaches, we show that the identified membrane vesicles do not represent a specialized region of the ER. Our results challenge the model proposing that all PMPs are first sorted to the ER and subsequently exit that compartment in the Pex3-dependent manner.

In **Chapter 3** we addressed the origin and protein composition of the peroxisomal membrane vesicles in *S. cerevisiae pex3* cells using two genetic screens that were based on automated mating, sporulation and mutant selection approaches combined with automated fluorescence microscopy. One of these screens, aiming to determine the protein composition of the peroxisomal vesicles, resulted in a list of proteins that co-localized with the peroxisomal vesicle marker protein

Pex14. We failed to identify proteins crucial for peroxisomal vesicle formation. Some of the risks associated with high-throughput approaches are discussed.

Our co-localization screen identified Nvj2, a protein of nucleus-vacuole junctions (NVJs), as a possible candidate protein associated with pre-peroxisomal vesicles in *S. cerevisiae* *pex3* cells (**Chapter 3**). Vac8, another NVJ protein, was identified in two independent organelle proteomics studies. **Chapter 4** describes studies aiming to elucidate whether Vac8 plays a role in peroxisome biogenesis in *H. polymorpha*. First we showed that *H. polymorpha* Vac8 is required for the formation of NVJs and vacuole inheritance, like in *S. cerevisiae*. However, *HpVac8* is not required for vacuole fusion. The composition of the *H. polymorpha* NVJ differs from the one in *S. cerevisiae*, because of the absence of Nvj1, which is the second essential component for the formation of NVJs in baker's yeast. We were unable to detect any peroxisomal defect in *H. polymorpha* cells lacking Vac8, indicating that this protein most likely is not important in peroxisome biology.

Some reports suggest that peroxisomes are formed *de novo* from the ER in WT yeast cells. However, other studies indicate that peroxisomes predominantly multiply by fission and are carefully segregated over mother and bud during yeast budding. In **Chapter 5** we describe the consequences of impaired peroxisome fission and inheritance on the peroxisome population in *H. polymorpha*. Detailed fluorescence microscopy analysis revealed that peroxisome proliferation and inheritance are completely blocked in a *pex11 inp2* double deletion strain, because peroxisomes could not be detected in newly formed buds of this double mutant. At later stages, however, these structures could be identified, implying that the buds acquire them *de novo*. This study suggests that in *H. polymorpha* *de novo* peroxisome formation can occur, but serves only as a rescue mechanism for the formation of peroxisomes in mutant cells that lack these organelles.