

University of Groningen

Complete Genome Sequences of 28 Lactococcal Bacteriophages Isolated from Failed Dairy Fermentation Processes

Marcelli, Barbara; de Jong, Anne; Janzen, Thomas; Serrano, Mariela; Kok, Jan; Kuipers, Oscar P

Published in:
Microbiology resource announcements

DOI:
[10.1128/MRA.01535-19](https://doi.org/10.1128/MRA.01535-19)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Marcelli, B., de Jong, A., Janzen, T., Serrano, M., Kok, J., & Kuipers, O. P. (2020). Complete Genome Sequences of 28 Lactococcal Bacteriophages Isolated from Failed Dairy Fermentation Processes. *Microbiology resource announcements*, 9(12), [e01535-19]. <https://doi.org/10.1128/MRA.01535-19>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).


Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Complete Genome Sequences of 28 Lactococcal Bacteriophages Isolated from Failed Dairy Fermentation Processes

Barbara Marcelli,^a Anne de Jong,^a Thomas Janzen,^b Mariela Serrano,^c Jan Kok,^a  Oscar P. Kuipers^a

^aDepartment of Molecular Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Groningen, The Netherlands

^bBacterial Physiology and Improvement, R&D Discovery, Chr. Hansen A/S, Hørsholm, Denmark

^cR&D Department, CSK Food Enrichment, Wageningen, The Netherlands

ABSTRACT *Lactococcus lactis* is a Gram-positive lactic acid bacterium commonly used in the dairy industry for the production of fermented foods such as buttermilk and a wide variety of cheeses. Here, we report the complete genome sequences of 28 bacteriophages infecting different *L. lactis* industrial starter strains isolated from dairy plants throughout the world.

Bacteriophage infection of *Lactococcus lactis* strains, which are used as starter cultures in dairy fermentation processes, is one of the main causes of fermentation failure and causes great economic losses for dairy industries (1, 2). Bacteriophages infecting *L. lactis* have been divided into 10 species (3), and those belonging to species c2, 936, or P335 are more commonly encountered in dairy plants (4, 5). However, isolates belonging to other species have also been reported to cause dairy fermentation halts (6–10). Studying lactococcal bacteriophages is of crucial importance for understanding phage-host interactions in dairy environments and for preventing the spread of infections in production lines. Here, we present the complete genome sequences of 28 lactococcal bacteriophages isolated over the past 3 decades from failed fermentations in dairy plants located throughout the world.

The bacteriophages were isolated from whey samples and plaque purified three times on their industrial lactococcal hosts at 30°C in M17 medium using the soft-agar overlay assay (11). A single plaque was finally propagated in liquid M17 medium on the same host to obtain a pure phage lysate. Phage purification was achieved by polyethylene glycol 800 (PEG 800) precipitation, and genomic DNA was extracted via phenol-chloroform purification using a previously described method (10). Samples were prepared for sequencing using a standard Illumina genomic library. The sequencing process delivered 5 million paired-end reads (2 × 150 bp) per sample. Quality control of the total sequence reads was performed with FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc>), and the sequence reads were trimmed using Trimmomatic v0.36 (12). Genome assembly was performed using the A5-miseq pipeline with default parameters (13). The sequences of the contigs obtained were subjected to a BLAST search against the total genomes of known lactococcal strains and bacteriophages present in the NCBI databases. Contigs that contained contaminating host chromosomal DNA were discarded, and the contigs that entailed a full phage genome sequence were annotated using RASTtk with default parameters (14). Bacteriophage sequences were subsequently assigned to known species using two previously described multiplex PCR methods (15, 16). Fifteen and nine isolates could be assigned to the c2 and 936 species, respectively, using this approach. Following a previously proposed classification scheme (17), bacteriophages belonging to the c2 species were further classified into the subspecies c2 and bIL67. The analysis was conducted by

Citation Marcelli B, de Jong A, Janzen T, Serrano M, Kok J, Kuipers OP. 2020. Complete genome sequences of 28 lactococcal bacteriophages isolated from failed dairy fermentation processes. *Microbiol Resour Announc* 9:e01535-19. <https://doi.org/10.1128/MRA.01535-19>.

Editor Simon Roux, DOE Joint Genome Institute

Copyright © 2020 Marcelli et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Oscar P. Kuipers, o.p.kuipers@rug.nl.

Received 29 December 2019

Accepted 26 February 2020

Published 19 March 2020

TABLE 1 Characteristics of and accession numbers for the bacteriophages studied here

Bacteriophage name	Genome length (kb)	Sequencing coverage (1,000 fold)	No. of ORFs	Species ^a	GenBank accession no.	Host species	Host strain name
CHPC116	21.86	68.6	37	c2 (c2)	MN689507	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC20
CHPC122	22.1	67.8	41	c2 (c2)	MN689512	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC21
CHPC134	21.98	68.2	38	c2 (c2)	MN689515	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC22
CHPC966	21.74	69	37	c2 (bIL67)	MN689526	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC23
CHPC967	22.4	67	42	c2 (c2)	MN689527	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC24
CHPC972	23.28	64.4	40	c2 (c2)	MN689528	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC25
CHPC973	22.3	67.2	36	c2 (c2)	MN689529	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC26
CHPC1020	22.41	67	36	c2 (bIL67)	MN689505	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC27
CHPC1161	21.34	70.3	34	c2 (c2)	MN689506	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC28
CHPC1170	21.74	69	40	c2 (c2)	MN689508	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC29
CHPC1182	20.75	72.3	34	c2 (bIL67)	MN689510	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC30
CHPC1183	20.03	74.8	40	c2 (c2)	MN689511	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC31
CHPC1242	21.14	71	35	c2 (c2)	MN689513	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC32
5171F	21.07	71.2	37	c2 (c2)	MN689503	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	MG1363
5205F	21.21	70.7	35	c2 (c2)	MN689504	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	MG1363
CHPC52	29.65	50.6	54	936	MN689519	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC33
CHPC129	30.83	48.6	55	936	MN689514	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC34
CHPC361	30.16	49.7	55	936	MN689517	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC35
CHPC362	27.64	54.2	46	936	MN689518	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC36
CHPC781	29.21	51.3	56	936	MN689520	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC37
CHPC958	32.65	45.5	62	936	MN689522	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC38
CHPC959	29.34	51.1	59	936	MN689523	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC39
CHPC964	29.92	50.1	56	936	MN689524	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC40
CHPC965	25.32	59.2	47	936	MN689525	<i>Lactococcus lactis</i> subsp. <i>cremoris</i>	CH_LC41
CHPC148	33.55	44.7	51	BK5-T	MN689516	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC42
CHPC836	36.48	41.1	57	BK5-T	MN689521	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC43
CHPC974	33.79	44.4	60	BK5-T	MN689530	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC44
CHPC1175	36.33	41.3	52	BK5-T	MN689509	<i>Lactococcus lactis</i> subsp. <i>lactis</i>	CH_LC45

^a The subspecies of c2 bacteriophages is given.

comparing the C and N termini of their predicted receptor-binding proteins (RBP) (as shown by RASTtk annotation) and the complete sequences of the proteins encoded by the two adjacent open reading frames (ORFs) with those of the reference phages c2 and D4410. Based on the high (>80%) nucleotide similarity of their aforementioned ORFs to those of the two reference lactococcal bacteriophages, 12 isolates proved to belong to the c2 subspecies; the remaining 3 were shown to be part of the bIL67 subspecies. In cases in which no species-level results were obtained with the multiplex PCR approach, phage species were assigned via BLAST comparison of total genome sequences with publicly available lactococcal bacteriophage genomes. Four Bk5-T bacteriophages were identified in this way, based on >72% whole-genome similarity and a conserved genome organization, compared with known Bk5-T lactococcal bacteriophages.

The GC contents of the analyzed bacteriophages range from 34 to 36.4%. The genome lengths of the isolates range from 20 to 23.2 kb for the c2 phages, from 25.3 to 32.6 kb for phages of the 936 species, and from 25.3 to 32.6 kb for the Bk5-T members. The numbers of predicted ORFs range from 34 to 42 among members of the c2 species, from 46 to 62 among 936 phages, and from 51 to 60 for Bk5-T isolates.

Data availability. The complete genomic sequences of the 28 bacteriophages described here are available at GenBank under the accession numbers reported in Table 1. The SRA data for each genome reported here are available at the NCBI under BioProject accession number [PRJNA606016](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA606016).

REFERENCES

1. Marco MB, Moineau S, Quiberoni A. 2012. Bacteriophages and dairy fermentations. *Bacteriophage* 2:149–158. <https://doi.org/10.4161/bact.21868>.
2. Song AA-L, In LLA, Lim SHE, Rahim RA. 2017. A review on *Lactococcus lactis*: from food to factory. *Microb Cell Fact* 16:55. <https://doi.org/10.1186/s12934-017-0669-x>.
3. Mahony J, Van Sinderen D. 2014. Current taxonomy of phages infecting lactic acid bacteria. *Front Microbiol* 5:7. <https://doi.org/10.3389/fmicb.2014.00007>.
4. Deveau H, Labrie SJ, Chopin M-C, Moineau S. 2006. Biodiversity and classification of lactococcal phages. *Appl Environ Microbiol* 72:4338–4346. <https://doi.org/10.1128/AEM.02517-05>.

5. Oliveira J, Mahony J, Hanemaaijer L, Kouwen T, van Sinderen D. 2018. Biodiversity of bacteriophages infecting *Lactococcus lactis* starter cultures. *J Dairy Sci* 101:96–105. <https://doi.org/10.3168/jds.2017-13403>.
6. Fortier L-C, Bransi A, Moineau S. 2006. Genome sequence and global gene expression of Q54, a new phage species linking the 936 and c2 phage species of *Lactococcus lactis*. *J Bacteriol* 188:6101–6114. <https://doi.org/10.1128/JB.00581-06>.
7. Garneau JE, Tremblay DM, Moineau S. 2008. Characterization of 1706, a virulent phage from *Lactococcus lactis* with similarities to prophages from other Firmicutes. *Virology* 373:298–309. <https://doi.org/10.1016/j.virol.2007.12.002>.
8. Chopin A, Deveau H, Ehrlich SD, Moineau S, Chopin M-C. 2007. KSY1, a lactococcal phage with a T7-like transcription. *Virology* 365:1–9. <https://doi.org/10.1016/j.virol.2007.03.044>.
9. Samson JE, Moineau S. 2010. Characterization of *Lactococcus lactis* phage 949 and comparison with other lactococcal phages. *Appl Environ Microbiol* 76:6843–6852. <https://doi.org/10.1128/AEM.00796-10>.
10. Marcelli B, de Jong A, Karsens H, Janzen T, Kok J, Kuipers OP. 2019. A specific sugar moiety in the *Lactococcus lactis* cell wall pellicle is required for infection by CHPC971, a member of the rare 1706 phage species. *Appl Environ Microbiol* 85:e01224-19. <https://doi.org/10.1128/AEM.01224-19>.
11. Schuster H. 1962. Bacteriophages, von M. H. Adams. Interscience Publishers, Inc., New York-London 1959. 1. Aufl., XVIII, 592 S., 26 Tab., 16 Abb., geb. £6.50. *Angew Chem* 74:164. <https://doi.org/10.1002/ange.19620740437>.
12. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
13. Coil D, Jospin G, Darling AE. 2015. A5-miseq: an updated pipeline to assemble microbial genomes from Illumina MiSeq data. *Bioinformatics* 31:587–589. <https://doi.org/10.1093/bioinformatics/btu661>.
14. Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>.
15. Labrie S, Moineau S. 2000. Multiplex PCR for detection and identification of lactococcal bacteriophages. *Appl Environ Microbiol* 66:987–994. <https://doi.org/10.1128/aem.66.3.987-994.2000>.
16. del Rio B, Binetti AG, Martín MC, Fernández M, Magadán AH, Alvarez MA. 2007. Multiplex PCR for the detection and identification of dairy bacteriophages in milk. *Food Microbiol* 24:75–81. <https://doi.org/10.1016/j.fm.2006.03.001>.
17. Millen AM, Romero DA. 2016. Genetic determinants of lactococcal C2viruses for host infection and their role in phage evolution. *J Gen Virol* 97:1998–2007. <https://doi.org/10.1099/jgv.0.000499>.