

University of Groningen

Surgeons' Ability to Predict the Extent of Surgery Prior to Cytoreductive Surgery With Hyperthermic Intraperitoneal Chemotherapy

Hentzen, Judith; Plas, van der, Willemijn; Been, Lukas; Hoogwater, Frederik; Ginkel, van, Robert; Dam, van, Go; Hemmer, Patrick; Kruijff, Schelto

Published in:
Annals of Surgical Oncology

DOI:
[10.1245/s10434-020-08237-8](https://doi.org/10.1245/s10434-020-08237-8)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hentzen, J., Plas, van der, W., Been, L., Hoogwater, F., Ginkel, van, R., Dam, van, G., ... Kruijff, S. (2020). Surgeons' Ability to Predict the Extent of Surgery Prior to Cytoreductive Surgery With Hyperthermic Intraperitoneal Chemotherapy. *Annals of Surgical Oncology*, 27(8), 2997-3008. <https://doi.org/10.1245/s10434-020-08237-8>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record.

Please cite this article as <https://doi.org/10.4097/kja.20294>

Epub ahead of print

Postoperative neurocognitive disorders

Setayesh R. Tasbihgou¹ and Anthony R. Absalom²

1. PhD student, Department of Anesthesiology, University Medical Centre Groningen, University of Groningen, The Netherlands

2. Professor, Department of Anesthesiology, University Medical Centre Groningen, University of Groningen, The Netherlands

Running title: Postoperative neurocognitive disorders

Corresponding author: Setayesh R. Tasbihgou, M.D., Department of Anesthesiology, University Medical Centre Groningen, University of Groningen, The Netherlands

Email: s.r.tasbihgou@umcg.nl

Address: Hanzeplein 1, Groningen , 9713 GZ, Groningen, The Netherlands

Prior presentations: not applicable

Conflict of interest: No potential conflict of interest relevant to this article was reported

Funding: the writing and submission of this article was sponsored by the Korean Journal of Anesthesiology

Acknowledgments: not applicable

IRB number: not applicable

Clinical Trial registration number: not applicable

Postoperative neurocognitive disorders

Running title: Postoperative neurocognitive disorders

Epub ahead of print

Abstract

A decline in cognitive function is a frequent complication after major surgery. Postoperative cognitive impairments have generally been divided into short (postoperative delirium) and long-term disturbances (postoperative cognitive dysfunction [POCD]). Long-term impairments are often subtle and overlooked. They need to be objectively assessed with neuropsychological tests in order to be diagnosed. Although POCD has been the subject of considerable research over the past decades, it remains uncertain why some patients do not return to preoperative levels of cognitive function. Surgery and anesthesia have both been implicated in playing a role in developing POCD, and certain patient-related factors, such as advanced age and low preoperative baseline cognitive function have consistently been found to predict postoperative cognitive decline. This article will provide an overview of POCD, and its etiology, and provide advice on possible strategies on preventing it.

Keywords: Aged; Delirium; Frail elderly; Inflammation; Neurocognitive disorders; Perioperative care; Postoperative cognitive complications.

1 **Introduction**

2 It has been over five decades since P.D. Bedford observed that surgery in elderly patients was
3 followed by a significant cognitive decline that lasted for an extensive period of time [1]. Through
4 interviews (with patients and relatives) and subjective assessment, he found that 7% of his
5 (elderly) patients who underwent surgery and received general anesthesia had developed signs of
6 cognitive impairment. He published these findings in The Lancet, concluding that 'the allegation
7 "He's never been the same since his operation " is sometimes true, and that an irreversible gross
8 dementia is occasionally the aftermath of surgical operations under general anesthesia [1].'

9 By the late 1980s psychometric tests were being used to objectively assess cognitive decline after
10 surgery, particularly in patients undergoing cardiac surgery [2]. These studies too, consistently
11 documented long-term cognitive disorder in elderly patients, although with varying incidences and
12 severity. As a result, the concept postoperative cognitive dysfunction (POCD) developed as a
13 diagnosis based on these objective measurements. Although surgery and anesthesia have improved
14 dramatically since then, our exact understanding of when, how and why some patients do not return
15 to baseline cognitive function remains elusive. As cognitive dysfunction, in the form of delirium
16 has been shown to be important for perioperative outcome and mortality [3–5], it is also important
17 to consider the effects of long-term cognitive impairment and its possible risk factors. In this
18 review, we present a brief overview of POCD and its etiology, and provide advice on possible
19 strategies on preventing it.

21 **Postoperative cognitive impairment; delirium and POCD**

22 Cognitive impairment after surgery is common, particularly in elderly patients. These
23 impairments have generally been divided into short (delirium) and long-term disturbances
24 (postoperative cognitive dysfunction [POCD]) [6]. The former is familiar among many clinicians

1 and well-defined according to the Diagnostic and Statistical Manual of Medical Disorders (DSM)-5
2 [7]. It states that delirium consists of impairments in attention, awareness and cognition. Cognition
3 is considered to be a dynamic state, involving multiple domains such as memory, orientation,
4 language, visuospatial ability or perception [8]. It fluctuates throughout the day, and is affected by
5 both endogenous and exogenous factors [9]. The incidence of postoperative delirium is reported to
6 be between 20–45% of the older adult surgery patients [10,11].

7 The term POCD on the other hand, has been used to refer to any signs of new cognitive
8 impairment that exceeds the expected length of time needed to recover from the acute effects of
9 surgery and anesthesia [6,12,13]. Unlike delirium, which is a relatively simple and recognizable
10 syndrome, POCD is clinically far less apparent as it often only manifests as mild cognitive decline
11 in one or more cognitive domains [6,14,15]. Furthermore, the DSM-5 does not list POCD as a
12 diagnosis. In 2018, this prompted an expert panel of scientists and clinicians, The International
13 Perioperative Cognition Nomenclature Working Group, to address, clarify and give structure to
14 POCD and other perioperative cognitive impairments, whilst proposing new nomenclature to be
15 used in relation to these terms [16].

16 This working group stated that all cognitive changes associated with surgery and anesthesia
17 should be summarized under the term ‘perioperative neurocognitive disorders,’ thus aligning these
18 impairments with the clinical diagnostic criteria for ‘neurocognitive disorders (NCD)’ already
19 applied in the DSM-5 [7,16]. The working group recommends assessing POCD at least 30 days
20 after surgery, at which point most patients are expected to have recovered, physically,
21 physiologically and emotionally from surgery and hospitalization [16]. If assessed too early, the
22 effects of POCD may be overshadowed by acute postoperative delirium or by other cognitive
23 complications that may arise from immobility, sleep deprivation and ongoing pharmacological

1 interventions [2]. When cognitive impairment manifests itself beyond 12 months after surgery,
2 mild or major [e.g. dementia] Neurocognitive Disorders should be considered over POCD [16].

3 The term ‘delayed neurocognitive recovery’ may be used to describe a cognitive disorder that is
4 detected within 30 days after surgery when delirium has been excluded. Table 1 summarizes the
5 recommendations offered by the working group [9,16].

6

7 **Assessing POCD**

8 Unlike delirium, the diagnosis POCD has primarily been confined to research. Its diagnosis relies
9 on objectively measurable cognitive decline assessed with neuropsychological tests [12,13,17].
10 Subjective reports of cognitive changes by patients or proxies are also relevant, however, most
11 studies comparing cognitive complaints and neuropsychological test results were unable to find a
12 significant correlation [13,18]. Certain cognitive functions may be less relevant to a patient’s daily
13 life, and as such, any dysfunction may be overlooked by the patient. There is no agreed upon
14 definition for POCD, but it generally refers to impairment of memory, learning, concentration,
15 attention or psychomotor performance [12,16]. Neuropsychological tests are often specific to one
16 of these cognitive domains.

17 Neuropsychological tests that were used in a key international multicenter study on POCD (the
18 International Study of Post-Operative Cognitive Dysfunction, ISPOCD 1) are described in Table 2.
19 There are a wide variety of neuropsychological tests, which all have different levels of sensitivity,
20 and reliability. The ISPOCD mostly used written tests. Our research group, however, favors
21 computerized tests, such as the Cogstate Computerized Cognitive Test Battery ®, because of its
22 ease of use, versatility and the availability of age-matched control group test data.

23 Certain tests are more vulnerable to the effects of practice, and have a poor test-retest reliability
24 [9,13,17]. Others, notoriously suffer from floor and ceiling-effects resulting from tests being either

1 too difficult or too easy to detect subtle changes [13]. The method with which these test results are
2 interpreted also varies throughout literature [2]. Test batteries, consisting of multiple tests are able
3 to assess various cognitive domains and are recommended as they are able to describe the brain
4 functions in more detail and sensitivity [2,13]. To measure cognitive decline, investigators should
5 determine the change between baseline preoperative cognitive function and postoperative cognitive
6 function. In order to correct for age-related test-retest variability, determining the change in
7 cognitive function with the use of the reliable change index (RCI) is recommended, as it calculates
8 this change with reference to the expected change found within an age-matched control group [13].
9

10 **Incidence of POCD**

11 The incidence of POCD ranges from 20 to 50% of older patients 3 months after cardiac surgery
12 and in 5 to 55% of those undergoing major noncardiac surgeries [12,19–23]. This large variation is
13 the result of the methodological differences between studies, making comparison of data often
14 difficult. In addition to the various types of test that may be administered for measuring cognitive
15 change, the degree of change and cut-offs necessary for determining POCD have also varied
16 throughout literature. Generally, POCD is divided into mild or major neurocognitive decline, if
17 testing exhibits a decline of > 1 or > 2 standard deviations of cognitive function compared to
18 preoperative cognitive performance, respectively. As described above, the timing of tests is also a
19 known source of variability; the later the cognitive assessment is conducted and the more stringent
20 statistical criteria for identifying POCD, the lower the reported incidence [13].

21 This point is illustrated by the large multicenter ISPOCD study conducted in 1998, which
22 observed 1000 patients (> 60 years) undergoing various noncardiac surgeries [12]. A
23 comprehensive neuropsychological test battery was administered with a strict criterion for POCD.
24 This study found that 25.8% (95% CI, 23.1–28.5) of patients showed signs of cognitive dysfunction

1 1 week after surgery. Cognitive dysfunction at 3 months after surgery was 9.9% (95% CI, 8.1–
2 12.0). A later study by Monk et al., found similar incidences of POCD in 365 patients undergoing
3 noncardiac surgery; 41.4% (95% CI, 36.2–46.7) at discharge and 12.7% (95% CI, 8.9–16.4) at 3
4 months [23]. A recent systematic review of 24 studies found that the incidence of POCD at 3
5 months was 11.7% (95% CI 10.9–12.5), although they concluded that major differences in
6 methodology and definitions accounted for variations in the results [24].

7

8 **Pathogenesis of POCD**

9 Despite a growing volume of research concerning POCD, the exact etiology for cognitive decline
10 after surgery and anesthesia is still not well understood. Surgery-, anesthesia- and patient-related
11 factors have all been implicated in playing a role in developing POCD, and support for various
12 hypotheses have changed markedly over the years. Historically, a poor cognitive outcome after
13 surgery was often regarded to be the consequence of cerebral hypoperfusion and hypoxemia [2,14].
14 Indeed, inadequate cerebral oxygenation will result in brain damage and cognitive decline.
15 Although intuitively compelling, no strong evidence has been found in favor of POCD being the
16 direct consequence of impaired cerebral hemodynamics and oxygenation [2,22,25]. This was also
17 confirmed by the ISPOCD, which monitored perioperative blood pressure and oxygenation and
18 showed that POCD occurred in the absence of perioperative hypoxemia or hypotension [12].

19 Factors such as the type and duration of surgery and anesthesia, have also often been presumed to
20 be associated with the incidence of POCD. However, this has not yet been conclusive. A
21 comprehensive study by Evered et al. [19] compared the incidence of POCD after coronary
22 angiography under sedation, total hip replacement and coronary artery bypass graft under general
23 anesthesia. Interestingly, the incidence of POCD was similar and thus independent of the nature of
24 surgery or type of anesthesia administered. Furthermore, evidence on whether volatile or

1 intravenous anesthetics may be related to POCD has also been controversial and conflicting [26].
2 Moreover, other studies have not found any correlation between regional or general anesthesia and
3 the incidence of POCD, which further supports the argument that the type of anesthesia appears to
4 be unrelated to the occurrence of POCD [27,28]. It is therefore unlikely that POCD is solely caused
5 by anesthesia or surgery.

6 A recurring theme and the current rationale for the pathogenesis of cognitive dysfunction
7 encompasses the role of an inflammatory response to surgery and anesthesia [2,14,25]. It is
8 commonly known that inflammatory processes, such as those associated with pneumonia or a
9 urinary tract infection, are regularly accompanied by cognitive decline, particularly in the elderly
10 [29,30]. Extending this model to postoperative cognitive dysfunction, it is thought that the release
11 of proinflammatory mediators, triggered by peripheral surgical stress or trauma, may result in an
12 exaggerated systemic inflammatory response leading to neuroinflammation in vulnerable
13 individuals [14,25,31]. The release of inflammatory cytokines is known to lead to endothelial
14 dysfunction, and also the disruption of tight junctions, which results in an increased blood-brain-
15 barrier (BBB) permeability [25,32]. Consequently, systemic inflammatory cytokines will penetrate
16 the BBB, triggering neuroinflammation and the activation of the neuronal immune system,
17 including microglia and astrocytes [25,31,33]. Inflammatory mediators are also produced within
18 the brain, as a result of peripheral-to-central signaling via humoral and neuronal pathways [34].
19 The consequences of this immune response are healing, but if excessive, it may also result in further
20 (cerebral) tissue damage in the form of increased synaptic dysfunction, inhibition of neurogenesis,
21 and neuronal death [25].

22 In mouse models, surgery caused hippocampal-dependent memory impairment that was
23 associated with increased expression of plasma cytokines, as well as reactive microgliosis and
24 interleukin (IL)-1 β transcription and expression in the hippocampus [35,36]. By inhibiting IL-1 β ,

1 these neuroinflammatory changes were mitigated. Another study showed that tumor necrosis factor
2 (TNF)- α inhibition was also able to limit the release of IL-1 and prevent neuroinflammation and
3 cognitive decline in mice [37]. Thus, peripheral surgical injury can result in inflammation and
4 neuroinflammation. However, interpreting and judging the significance of an inflammatory
5 markers is challenging, as inflammation is a normal physiological response to injury [25].

6 Generally, inflammation is only harmful when proinflammatory responses outweigh the anti-
7 inflammatory response. Certain patient-related factors are known to exacerbate proinflammatory
8 responses or result in some patients being more vulnerable to the effects of inflammation.
9 Advanced age has been consistently associated with POCD throughout the literature [2,9].
10 Structural cerebral changes, such as a reduction in grey matter volume and myelinated axon length
11 are normal changes that occur with aging [25,38]. The normal decline of cognitive function in the
12 elderly might possibly be further exacerbated by the loss of neuronal dendrite spines, as well as
13 alterations in synaptic transmission and receptors [39]. Furthermore, blood-brain-barrier
14 dysfunction has also been found in older patients even in the absence of surgery [40]. This decline
15 in ‘cognitive reserve’ may thus explain how elderly patients are more susceptible to effects of
16 inflammation and therefore neuronal injury. A low preoperative cognitive function, and lower
17 education level have also been frequently associated with POCD, also suggesting the vulnerability
18 of a reduced ‘cognitive reserve’ [2,12,23,41].

19 Predisposing patient factors may also exaggerate an inflammatory response, as a result of
20 ‘immune priming’ [25]. For instance, normal aging without any comorbidities has been associated
21 with a low-grade inflammatory activity and increased levels of plasma tumor necrosis factor- α and
22 IL-6 compared to younger patients [42]. Elderly patients are also more susceptible for sepsis [43].
23 It is unsurprising that patients of advanced age may be more likely to develop an exaggerated
24 inflammatory response as a consequence of surgery. The immune system activation caused by

1 atherosclerosis, or neurogenerative disorders such as Alzheimer's and Parkinson's, may also prime
2 individuals to develop an excessive inflammatory response [44,45]. The presence of Alzheimer's
3 dementia biomarkers in cerebrospinal fluid has been shown to be associated with POCD at 3
4 months, which has also led to the notion that they may involve similar mechanisms [46].

5 Considering the role of inflammation, several studies have attempted to prevent postoperative
6 cognitive dysfunction with anti-inflammatory drugs [14,47]. One study on the effects of high-dose
7 intraoperative dexamethasone administration in cardiac surgery, showed that it did not reduce the
8 risk of POCD [48]. Furthermore, other studies have found that lidocaine, magnesium and
9 complement cascade inhibitors also failed to prevent POCD [49–51]. These negative findings and
10 the understanding that not all elderly patients undergoing major surgery develop POCD or not all
11 patients with atherosclerosis develop POCD after cardiac surgery, reflects the pathophysiological
12 complexity of POCD.

14 **Prevention of POCD**

15 Although a firm understanding for the causes of POCD is absent, improving cognitive outcome
16 after surgery remains an important objective for anesthesiologists and surgeons alike. To date, no
17 pharmacological intervention has convincingly been shown to mitigate the incidence or magnitude
18 of POCD [14]. Dexmedetomidine, an anesthetic agent with neural anti-inflammatory effects, has
19 been found to be potentially effective at reducing the incidence of postoperative delirium, however
20 any evidence that it may be effective at reducing POCD is incomplete [52,53]. Deep sedation has
21 also been identified as a risk factor for delirium, and several studies have found that measuring the
22 depth of anesthesia (with electroencephalogram monitors) was effective at reducing postoperative
23 delirium, however, there is conflicting evidence that POCD can also be prevented with the same
24 measures [54–58]. If possible, deliriogenic [pre]medications such as benzodiazepines should also

1 be avoided [59,60]. Pain and increased post-operative opioid consumption are known to increase
2 the risk of delirium and have also been associated with POCD [60]. Although sufficient pain-
3 management is mandatory, opioid-sparing analgesia may be an effective measure at alleviating
4 some of this risk. Early postoperative mobilization, and a fast-track postoperative approach may
5 help in this respect [61].

6 Generally, for [non-pharmacological] preventive measures to be significantly effective, multiple
7 (interdisciplinary) interventions, covering various domains, should be considered [2,9,14]. Patients
8 with a possible high risk for POCD should be identified preoperatively and cognitively assessed.
9 When possible, predisposing factors should be modified and adjusted so that patients are
10 sufficiently prepared for surgery. Preparing patients, and their relatives, adequately by informing
11 them about possible postoperative cognitive changes is also beneficial [62]. Extended periods of
12 preoperative fasting and dehydration should be avoided, as should unnecessary postponement of
13 surgery [63]. Peri-and postoperative patient (re)orientation is essential. Encouraging patients to
14 wear their glasses and hearing aids, and early removal of catheters and lines are known to be
15 effective at reducing postoperative delirium, and will help orientate patients and mobilize them
16 earlier, which may likely be effective at preventing POCD [60].

17 Numerous novel approaches that have been shown to improve cognitive function in older adults
18 have also been proposed as possible interventions that may prevent, or protect patients against
19 postoperative cognitive dysfunction. These proposed interventions involve diet interventions,
20 physical exercise programs, and brain stimulation and cognitive training [14,64]. Although these
21 strategies are known to improve overall cognition, few of them have been investigated as potential
22 and feasible interventions for POCD [2]. One study by Kawano et al. [65] found that preoperative
23 environmental enrichment (PEE), consisting of both cognitive and physical activity, was able to
24 attenuate neuroinflammation and improve cognitive function in old rats after abdominal surgery. In

1 humans, there is some evidence that preoperative physical status may improve postoperative
2 morbidity, however, cognitive advantages, if any, are not yet known [66,67]. Nonetheless, for
3 treatments for cognitive decline there appears to be some potential in improving lifestyle-based
4 factors, although further investigation is necessary.

5

6 **Conclusion**

7 In summary, many studies have drawn attention to neurocognitive dysfunction after surgery by
8 using neuropsychological assessments prior to and after surgery. Results on the incidence and
9 severity of postoperative cognitive decline is varying, mostly due to varying definitions for
10 diagnosing postoperative cognitive dysfunction. The incidence of POCD in older patients at 3
11 months ranges from 20 to 50% after cardiac surgery and 5 to 55% major noncardiac surgeries
12 [12,19–23].

13 Although the etiology of postoperative cognitive dysfunction is still not fully understood,
14 inflammatory-processes are currently considered to be central to its genesis. At the present time, no
15 clear anesthetic and surgical components have been found to influence POCD. Nevertheless,
16 several patient-related factors, such as advanced age, have been associated with an increased risk
17 for cognitive decline. As the age of the general population undergoing surgery is growing older,
18 investigations on preventive measures and interventions are warranted and they should be aptly
19 applied.

References

1. Bedford PD. Adverse cerebral effects of anaesthesia on old people. *Lancet* 1955; 269: 259-63.
2. Evered LA, Silbert BS. Postoperative Cognitive Dysfunction and Noncardiac Surgery. *Anesth Analg* 2018; 127: 496-505.
3. Kiely DK, Jones RN, Bergmann MA, Murphy KM, Orav EJ, Marcantonio ER. Association between delirium resolution and functional recovery among newly admitted postacute facility patients. *J Gerontol A Biol Sci Med Sci.* 2006; 61: 204-8.
4. Kiely DK, Marcantonio ER, Inouye SK, Shaffer ML, Bergmann MA, Yang FM, et al. Persistent delirium predicts greater mortality. *J Am Geriatr Soc.* 2009; 57: 55-61.
5. Deiner S, Silverstein JH. Postoperative delirium and cognitive dysfunction. *Br J Anaesth.* 2009; 103 Suppl 1: i41-6.
6. Monk TG, Price CC. Postoperative cognitive disorders. *Curr Opin Crit Care* 2011; 17: 376-81.
7. American Psychiatric Association., American Psychiatric Association. DSM-5 Task Force. Diagnostic and statistical manual of mental disorders : DSM-5. 5th ed. Washington, D.C.: American Psychiatric Association; 2013. xliv, 947 p. p.
8. Sachdev PS, Blacker D, Blazer DG, Ganguli M, Jeste DV, Paulsen JS, et al. Classifying neurocognitive disorders: the DSM-5 approach. *Nat Rev Neurol* 2014; 10: 634-42.
9. Olotu C. Postoperative neurocognitive disorders. *Curr Opin Anaesthesiol* 2020; 33: 101-8.
10. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet* 2014; 383: 911-22.

11. Rudolph JL, Marcantonio ER. Review articles: postoperative delirium: acute change with long-term implications. *Anesth Analg* 2011; 112: 1202-11.
12. Moller JT, Cluitmans P, Rasmussen LS, Houx P, Rasmussen H, Canet J, et al. Long-term postoperative cognitive dysfunction in the elderly ISPOCD1 study. ISPOCD investigators. International Study of Post-Operative Cognitive Dysfunction. *Lancet* 1998; 351: 857-61.
13. Rasmussen LS, Larsen K, Houx P, Skovgaard LT, Hanning CD, Moller JT, et al. The assessment of postoperative cognitive function. *Acta Anaesthesiol Scand* 2001; 45: 275-89.
14. Berger M, Terrando N, Smith SK, Browndyke JN, Newman MF, Mathew JP. Neurocognitive function after cardiac surgery: from phenotypes to mechanisms. *Anesthesiology* 2018; 129: 829-51.
15. Daiello LA, Racine AM, Yun Gou R, Marcantonio ER, Xie Z, Kunze LJ, et al. Postoperative delirium and postoperative cognitive dysfunction: overlap and divergence. *Anesthesiology* 2019; 131: 477-91.
16. Evered L, Silbert B, Knopman DS, Scott DA, DeKosky ST, Rasmussen LS, et al. Recommendations for the nomenclature of cognitive change associated with anaesthesia and surgery-2018. *Br J Anaesth* 2018; 121: 1005-12.
17. Rudolph JL, Schreiber KA, Culley DJ, McGlinchey RE, Crosby G, Levitsky S, et al. Measurement of post-operative cognitive dysfunction after cardiac surgery: a systematic review. *Acta Anaesthesiol Scand* 2010; 54: 663-77.
18. Newman S, Klinger L, Venn G, Smith P, Harrison M, Treasure T. Subjective reports of cognition in relation to assessed cognitive performance following coronary artery bypass surgery. *J Psychosom Res* 1989; 33: 227-33.

19. Evered L, Scott DA, Silbert B, Maruff P. Postoperative cognitive dysfunction is independent of type of surgery and anesthetic. *Anesth Analg* 2011; 112: 1179-85.
20. McDonagh DL, Mathew JP, White WD, Phillips-Bute B, Laskowitz DT, Podgoreanu MV, et al. Cognitive function after major noncardiac surgery, apolipoprotein E4 genotype, and biomarkers of brain injury. *Anesthesiology* 2010; 112: 852-9.
21. Newman MF, Kirchner JL, Phillips-Bute B, Gaver V, Grocott H, Jones RH, et al. Longitudinal assessment of neurocognitive function after coronary-artery bypass surgery. *N Engl J Med* 2001; 344: 395-402.
22. Kok WF, van Harten AE, Koene BM, Mariani MA, Koerts J, Tucha O, et al. A pilot study of cerebral tissue oxygenation and postoperative cognitive dysfunction among patients undergoing coronary artery bypass grafting randomised to surgery with or without cardiopulmonary bypass*. *Anaesthesia* 2014; 69: 613-22.
23. Monk TG, Weldon BC, Garvan CW, Dede DE, van der Aa MT, Heilman KM, et al. Predictors of cognitive dysfunction after major noncardiac surgery. *Anesthesiology* 2008; 108: 18-30.
24. Paredes S, Cortinez L, Contreras V, Silbert B. Post-operative cognitive dysfunction at 3 months in adults after non-cardiac surgery: a qualitative systematic review. *Acta Anaesthesiol Scand* 2016; 60: 1043-58.
25. van Harten AE, Scheeren TW, Absalom AR. A review of postoperative cognitive dysfunction and neuroinflammation associated with cardiac surgery and anaesthesia. *Anaesthesia* 2012; 67: 280-93.
26. Miller D, Lewis SR, Pritchard MW, Schofield-Robinson OJ, Shelton CL, Alderson P, et al. Intravenous versus inhalational maintenance of anaesthesia for postoperative

- cognitive outcomes in elderly people undergoing non-cardiac surgery. *Cochrane Database Syst Rev* 2018; 8: CD012317.
27. Rasmussen LS, Johnson T, Kuipers HM, Kristensen D, Siersma VD, Vila P, et al. Does anaesthesia cause postoperative cognitive dysfunction? A randomised study of regional versus general anaesthesia in 438 elderly patients. *Acta Anaesthesiol Scand* 2003; 47: 260-6.
 28. Wu CL, Hsu W, Richman JM, Raja SN. Postoperative cognitive function as an outcome of regional anesthesia and analgesia. *Reg Anesth Pain Med* 2004; 29: 257-68.
 29. Dunn N, Mullee M, Perry VH, Holmes C. Association between dementia and infectious disease: evidence from a case-control study. *Alzheimer Dis Assoc Disord* 2005; 19: 91-4.
 30. Perry VH, Cunningham C, Holmes C. Systemic infections and inflammation affect chronic neurodegeneration. *Nat Rev Immunol* 2007; 7: 161-7.
 31. Eckenhoff RG, Laudansky KF. Anesthesia, surgery, illness and Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 2013; 47: 162-6.
 32. Hovens IB, Schoemaker RG, van der Zee EA, Absalom AR, Heineman E, van Leeuwen BL. Postoperative cognitive dysfunction: Involvement of neuroinflammation and neuronal functioning. *Brain Behav Immun* 2014; 38: 202-10.
 33. Alam A, Hana Z, Jin Z, Suen KC, Ma D. Surgery, neuroinflammation and cognitive impairment. *EBioMedicine* 2018; 37: 547-56.
 34. Steinberg BE, Sundman E, Terrando N, Eriksson LI, Olofsson PS. Neural control of inflammation: implications for perioperative and critical care. *Anesthesiology* 2016; 124: 1174-89.
 35. Cibelli M, Fidalgo AR, Terrando N, Ma D, Monaco C, Feldmann M, et al. Role of interleukin-1beta in postoperative cognitive dysfunction. *Ann Neurol* 2010; 68: 360-8.

36. Terrando N, Rei Fidalgo A, Vizcaychipi M, Cibelli M, Ma D, Monaco C, et al. The impact of IL-1 modulation on the development of lipopolysaccharide-induced cognitive dysfunction. *Crit Care* 2010; 14: R88.
37. Terrando N, Monaco C, Ma D, Foxwell BM, Feldmann M, Maze M. Tumor necrosis factor-alpha triggers a cytokine cascade yielding postoperative cognitive decline. *Proc Natl Acad Sci U S A* 2010; 107: 20518-22.
38. Fjell AM, Walhovd KB. Structural brain changes in aging: courses, causes and cognitive consequences. *Rev Neurosci* 2010; 21: 187-221.
39. Dickstein DL, Kabaso D, Rocher AB, Luebke JI, Wearne SL, Hof PR. Changes in the structural complexity of the aged brain. *Aging Cell* 2007; 6: 275-84.
40. Hainsworth AH, Minett T, Andoh J, Forster G, Bhide I, Barrick TR, et al. Neuropathology of white matter lesions, blood-brain barrier dysfunction, and dementia. *Stroke* 2017; 48: 2799-804.
41. Gao L, Taha R, Gauvin D, Othmen LB, Wang Y, Blaise G. Postoperative cognitive dysfunction after cardiac surgery. *Chest* 2005; 128: 3664-70.
42. Bruunsgaard H, Pedersen BK. Age-related inflammatory cytokines and disease. *Immunol Allergy Clin North Am* 2003; 23: 15-39.
43. Martin GS, Mannino DM, Moss M. The effect of age on the development and outcome of adult sepsis. *Crit Care Med* 2006; 34: 15-21.
44. Bartels AL, Willemsen AT, Doorduyn J, de Vries EF, Dierckx RA, Leenders KL. [11C]-PK11195 PET: quantification of neuroinflammation and a monitor of anti-inflammatory treatment in Parkinson's disease? *Parkinsonism Relat Disord* 2010; 16: 57-9.

45. Doorduyn J, de Vries EF, Willemsen AT, de Groot JC, Dierckx RA, Klein HC. Neuroinflammation in schizophrenia-related psychosis: a PET study. *J Nucl Med* 2009; 50: 1801-7.
46. Evered L, Silbert B, Scott DA, Ames D, Maruff P, Blennow K. Cerebrospinal fluid biomarker for alzheimer disease predicts postoperative cognitive dysfunction. *Anesthesiology* 2016; 124: 353-61.
47. Rasmussen LS. Postoperative cognitive dysfunction: incidence and prevention. *Best Pract Res Clin Anaesthesiol* 2006; 20: 315-30.
48. Ottens TH, Dieleman JM, Sauer AM, Peelen LM, Nierich AP, de Groot WJ, et al. Effects of dexamethasone on cognitive decline after cardiac surgery: a randomized clinical trial. *Anesthesiology* 2014; 121: 492-500.
49. Mathew JP, Mackensen GB, Phillips-Bute B, Grocott HP, Glower DD, Laskowitz DT, et al. Randomized, double-blinded, placebo controlled study of neuroprotection with lidocaine in cardiac surgery. *Stroke* 2009; 40: 880-7.
50. Mathew JP, Shernan SK, White WD, Fitch JC, Chen JC, Bell L, et al. Preliminary report of the effects of complement suppression with pexelizumab on neurocognitive decline after coronary artery bypass graft surgery. *Stroke* 2004; 35: 2335-9.
51. Mathew JP, White WD, Schinderle DB, Podgoreanu MV, Berger M, Milano CA, et al. Intraoperative magnesium administration does not improve neurocognitive function after cardiac surgery. *Stroke* 2013; 44: 3407-13.
52. Carr ZJ, Cios TJ, Potter KF, Swick JT. Does dexmedetomidine ameliorate postoperative cognitive dysfunction? a brief review of the recent literature. *Curr Neurol Neurosci Rep* 2018; 18: 64.

53. Hu J, Vacas S, Feng X, Lutrin D, Uchida Y, Lai IK, et al. Dexmedetomidine prevents cognitive decline by enhancing resolution of high mobility group box 1 protein-induced inflammation through a vagomimetic action in mice. *Anesthesiology* 2018; 128: 921-31.
54. Lu X, Jin X, Yang S, Xia Y. The correlation of the depth of anesthesia and postoperative cognitive impairment: A meta-analysis based on randomized controlled trials. *J Clin Anesth* 2018; 45: 55-9.
55. Chan MT, Cheng BC, Lee TM, Gin T, Group CT. BIS-guided anesthesia decreases postoperative delirium and cognitive decline. *J Neurosurg Anesthesiol* 2013; 25: 33-42.
56. Radtke FM, Franck M, Lendner J, Kruger S, Wernecke KD, Spies CD. Monitoring depth of anaesthesia in a randomized trial decreases the rate of postoperative delirium but not postoperative cognitive dysfunction. *Br J Anaesth* 2013; 110 Suppl 1: i98-105.
57. Punjasawadwong Y, Chau-In W, Laopaiboon M, Punjasawadwong S, Pin-On P. Processed electroencephalogram and evoked potential techniques for amelioration of postoperative delirium and cognitive dysfunction following non-cardiac and non-neurosurgical procedures in adults. *Cochrane Database Syst Rev* 2018; 5: CD011283.
58. Bocskai T, Kovacs M, Szakacs Z, Gede N, Hegyi P, Varga G, et al. Is the bispectral index monitoring protective against postoperative cognitive decline? A systematic review with meta-analysis. *PLoS One* 2020; 15: e0229018.
59. Aldecoa C, Bettelli G, Bilotta F, Sanders RD, Audisio R, Borozdina A, et al. European Society of Anaesthesiology evidence-based and consensus-based guideline on postoperative delirium. *Eur J Anaesthesiol* 2017; 34: 192-214.
60. Inouye SK. A practical program for preventing delirium in hospitalized elderly patients. *Cleve Clin J Med* 2004; 71: 890-6.

61. Awada HN, Luna IE, Kehlet H, Wede HR, Høevsgaard SJ, Aasvang EK. Postoperative cognitive dysfunction is rare after fast-track hip- and knee arthroplasty - But potentially related to opioid use. *J Clin Anesth* 2019; 57: 80-6.
62. Hsieh TT, Yang T, Gartaganis SL, Yue J, Inouye SK. Hospital elder life program: systematic review and meta-analysis of effectiveness. *Am J Geriatr Psychiatry* 2018; 26: 1015-33.
63. Radtke FM, Franck M, MacGuill M, Seeling M, Lutz A, Westhoff S, et al. Duration of fluid fasting and choice of analgesic are modifiable factors for early postoperative delirium. *Eur J Anaesthesiol* 2010; 27: 411-6.
64. Ngandu T, Lehtisalo J, Solomon A, Levalahti E, Ahtiluoto S, Antikainen R, et al. A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER: a randomised controlled trial. *Lancet* 2015; 385: 2255-63.
65. Kawano T, Eguchi S, Iwata H, Tamura T, Kumagai N, Yokoyama M. Impact of preoperative environmental enrichment on prevention of development of cognitive impairment following abdominal surgery in a rat model. *Anesthesiology* 2015; 123: 160-70.
66. Robinson TN, Wu DS, Pointer L, Dunn CL, Cleveland JC, Jr., Moss M. Simple frailty score predicts postoperative complications across surgical specialties. *Am J Surg* 2013; 206: 544-50.
67. Robinson TN, Wu DS, Sauaia A, Dunn CL, Stevens-Lapsley JE, Moss M, et al. Slower walking speed forecasts increased postoperative morbidity and 1-year mortality across surgical specialties. *Ann Surg* 2013; 258: 582-8.

68. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975; 12: 189-98.
69. Brand N, Jolles J. Learning and retrieval rate of words presented auditorily and visually. *J Gen Psychol* 1985; 112: 201-10.
70. Reitan RM. Validity of the trail making test as an indicator of organic brain damage. *Percept Mot Skills* 1958: 271-6.
71. Bohnen N, Twijnstra A, Jolles J. Performance in the Stroop color word test in relationship to the persistence of symptoms following mild head injury. *Acta Neurol Scand* 1992; 85: 116-21.
72. Axelrod BN. Validity of the Wechsler abbreviated scale of intelligence and other very short forms of estimating intellectual functioning. *Assessment* 2002; 9: 17-23.
73. Houx PJ, Jolles J. Age-related decline of psychomotor speed: effects of age, brain health, sex, and education. *Percept Mot Skills* 1993; 76: 195-211.

Table 1. A Summary of the Recommendations for the New Nomenclature of Perioperative Disorders, from The International Perioperative Cognition Nomenclature Working Group [16].

Terms	Time period
Neurocognitive disorder [mild or major [e.g. dementia]]	Pre-existing/preoperative cognitive impairment <i>or</i> cognitive impairment developing after 12 months of surgery.
Emergence delirium	Delirium diagnosed within minutes or hours after surgery.
Postoperative delirium	Delirium diagnosed within days after surgery, up to 1 week or until discharge.
Delayed neurocognitive recovery	Cognitive decline up to 30 days after surgery.
Postoperative [neuro]cognitive dysfunction	Cognitive impairment detected between 30 days up to 12 months after surgery.

Table 2. Neuropsychological Tests Used in the in the International Study of Post-Operative Cognitive Dysfunction 1 (ISPOCD 1) Study

Tests	
Mini-mental state examination [MMSE] [68]	A commonly used assessment, initially developed to evaluate dementia. It assesses multiple cognitive domains including attention, memory and orientation.
Visual verbal learning test [69]	Based on Rey's auditive recall test. It assesses verbal memory, by asking patients to recall a list of words that they were presented with earlier.
Concept Shifting test [trail-making test] [70]	Also known as the trail-making tests A and B. It is used to assess executive function and attention, by asking subjects to connect a series of consecutive numbers, letters or both as quickly as possible.
Stroop color word interference test [71]	This test evaluates the ability to inhibit cognitive interreference from multiple congruent and incongruent stimuli.
Letter-digit coding test [symbol-digit substitution task] [72]	Used to assess executive function. Patients are presented with a series of digits and letters that are paired and another list of only digits. They are then asked to write the corresponding letter as fast as possible.
Four boxes test [73]	This test is computer-based. It is used to measure reaction time, by asking patients to select a black circle in one of four boxes on a screen as quickly as possible.
