

University of Groningen

Lessons learnt from scoring adjuvant colon cancer trials and meta-analyses using the ESMO-Magnitude of Clinical Benefit Scale V.1.1

Knapen, Daan Geert; Cherny, Nathan I; Zygoura, Panagiota; Latino, Nicola Jane; Douillard, Jean-Yves; Dafni, Urania; de Vries, E. G. E.; Groot, de, Derk Jan

Published in:
 ESMO Open

DOI:
[10.1136/esmoopen-2020-000681](https://doi.org/10.1136/esmoopen-2020-000681)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Knapen, D. G., Cherny, N. I., Zygoura, P., Latino, N. J., Douillard, J-Y., Dafni, U., de Vries, E. G. E., & Groot, de, D. J. (2020). Lessons learnt from scoring adjuvant colon cancer trials and meta-analyses using the ESMO-Magnitude of Clinical Benefit Scale V.1.1. *ESMO Open*, 5(5), [e000681]. <https://doi.org/10.1136/esmoopen-2020-000681>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Lessons learnt from scoring adjuvant colon cancer trials and meta-analyses using the ESMO-Magnitude of Clinical Benefit Scale V.1.1

Daan Geert Knapen,¹ Nathan I Cherny,² Panagiota Zygoura ,³ Nicola Jane Latino,⁴ Jean-Yves Douillard,⁴ Urania Dafni,^{5,6} Elisabeth G E de Vries,¹ Derk Jan de Groot ¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/esmoopen-2020-000681>).

To cite: Knapen DG, Cherny NI, Zygoura P, *et al*. Lessons learnt from scoring adjuvant colon cancer trials and meta-analyses using the ESMO-Magnitude of Clinical Benefit Scale V.1.1. *ESMO Open* 2020;5:e000681. doi:10.1136/esmoopen-2020-000681

Received 13 January 2020
Revised 12 March 2020
Accepted 13 March 2020



► <http://dx.doi.org/10.1136/esmoopen-2020-000743>

© Author (s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. Published by BMJ on behalf of the European Society for Medical Oncology.

For numbered affiliations see end of article.

Correspondence to
Dr Derk Jan de Groot;
d.j.a.de.groot@umcg.nl

[Click here to listen to the Podcast](#)

ABSTRACT

Background Form 1 of the European Society for Medical Oncology-Magnitude of Clinical Benefit Scale (ESMO-MCBS) serves to grade therapies with curative intent. Hitherto only few trials with curative intent have been field tested using form 1. We aimed to evaluate the applicability of the scale and to assess the reasonableness of the generated scores in early colon cancer, in order to identify shortcomings that may be rectified in future amendments.

Methods Adjuvant studies were identified in PubMed, Food and Drug Administration and European Medicines Agency registration sites, as well as ESMO and National Comprehensive Cancer Network guidelines. Studies meeting inclusion criteria were graded using form 1 of the ESMO-MCBS V.1.1 and field tested by ESMO Colorectal Cancer Faculty. Shortcomings of the scale were identified and evaluated.

Results Eighteen of 57 trials and 7 out of 14 meta-analyses identified met criteria for ESMO-MCBS V.1.1 grading. In stage III colon cancer, randomised clinical trials and meta-analyses of modulated 5-fluorouracil (5-FU) based chemotherapy versus surgery scored ESMO-MCBS grade A and randomised controlled trials (RCTs) and meta-analyses comprising oxaliplatin added to this 5-FU backbone showed a more modest additional overall survival benefit (grade A and B). For stage II colon cancer, the findings are less consistent. The fluoropyrimidine trials in stage II were graded 'no evaluable benefit' but the most recent meta-analysis demonstrated a 5.4% survival advantage after 8 years follow-up (grade A). RCTs and a meta-analysis adding oxaliplatin demonstrated no added benefit. Exploratory toxicity evaluation and annotation was problematic given inconsistent toxicity reporting and limited results of late toxicity. Field testers (n=37) reviewed the scores, 25 confirmed their reasonableness, 12 found them mostly reasonable. Moreover, they identified the inability of crediting improved convenience in non-inferiority trials as a shortcoming.

Conclusion Form 1 of the ESMO-MCBS V.1.1 provided very reasonable grading for adjuvant colon cancer studies.

Key questions

What is already known about this subject?

► Form 1 of the European Society for Medical Oncology-Magnitude of Clinical Benefit Scale (ESMO-MCBS) serves to grade therapies with curative intent. Hitherto only few trials with curative intent have been field tested using form 1.

What does this study add?

► We evaluated the applicability of the scale and assessed the reasonableness of the generated scores in early colon cancer. Form 1 of the ESMO-MCBS V.1.1 provided very reasonable grading for adjuvant colon cancer studies. Our exploratory analysis indicated that toxicity annotation is feasible but that the prevailing convention of physician reported toxicity may underestimate the true level of patient burden from both acute and late toxicity. The inability of crediting improved convenience in non-inferiority trials was identified as a shortcoming.

How might this impact on clinical practice?

► Future revisions of form 1 of the ESMO-MCBS will be cognoscente of these findings.

INTRODUCTION

Colorectal cancer is the third most common tumour in men, the second in women and second place in cancer-related cause of death in the world.¹ Mortality has declined over the years for several reasons, including colorectal cancer screening and more effective systemic therapies in both the adjuvant setting and metastatic disease.¹

Adjuvant therapies for colon cancer have evolved over the past 40 years. Early studies failed to show overall survival (OS) benefit of single agent therapy including 5-fluorouracil (5-FU) monotherapy compared with surgery




Figure 1 CONSORT diagram for RCTs and meta-analyses eligible for analysis

DFS, disease-free survival; ESMO-MCBS, European Society for Medical Oncology-Magnitude of Clinical Benefit Scale; MOF, Lomustine, Vincristine, and 5-Fluorouracil; OS, overall survival; PVI, portal vein infusion; RCT, randomised controlled trial; UFT, uracil and tegafur.

alone.² Adjuvant leucovorin modulated 5-FU (5-FU/LV) did, however, improve relative OS, but not absolute OS due to the increased incidence, and has been the standard of care since the mid-nineties. As of 2004, standard adjuvant therapy consists of a 5-FU/LV-based backbone to which oxaliplatin was added. Oxaliplatin did improve disease-free survival (DFS) and OS in stage III patients but it commonly caused substantial late toxicity (LT) with peripheral sensory neurotoxicity (PSN).³ Other agents including irinotecan, cetuximab and bevacizumab tested in the adjuvant setting, failed to show additional OS benefit.⁴⁻¹¹

The European Society for Medical Oncology (ESMO) has developed a validated and reproducible tool to assess the magnitude of clinical benefit of anticancer therapies of solid tumours. The ESMO-Magnitude of Clinical Benefit Scale (ESMO-MCBS) was initially published in 2015¹² and a revised V.1.1 was issued in 2017.¹³ The ESMO-MCBS incorporates different grading approaches for interventions with either curative intent, such as adjuvant treatment in colon cancer or with non-curative intent. Form 1 has been developed for assessing new approaches to adjuvant therapy or new potentially curative therapies. The scale ranges from A to C, with grades

A and B representing a substantial level of clinical benefit.^{12, 13} Currently, form 1 does not apply penalties for toxicity. However, patient advocates in consultation with the ESMO-MCBS working group have suggested that toxicity annotations should be introduced for treatments with a high prevalence of strong acute toxicity (AT) or LT and this is currently under consideration.

The validity of the ESMO-MCBS is predicated on adherence to the public policy ethical standard of ‘accountability for reasonableness’.¹²⁻¹⁵ Whereas the grading of treatments of advanced and incurable cancer using forms 2a, b, c and 3 of the ESMO-MCBS has been extensively field tested and reviewed for reasonableness, hitherto only 13 trials with curative intent, including adjuvant therapies, have been field tested using form 1.¹⁶ The main purpose was to evaluate the applicability of the scale in adjuvant colon cancer trials and further assess the reasonableness of the generated scores, in order to identify shortcomings that may be rectified in future amendments.

METHODS

Randomised controlled trials (RCTs) and meta-analyses in the adjuvant treatment of stage II/III colon cancer, published since the review of negative studies by Buyse *et al*² up to September 2019 were identified. Data were collected by electronic searches of PubMed (medical headings “colonic neoplasms” OR “colorectal neoplasms”, and the text words “adjuvant therapy” OR “adjuvant chemotherapy” OR “early colon cancer”) and by a manual review of Food and Drug Administration and European Medicines Agency registration sites and ESMO and National Comprehensive Cancer Network guidelines.^{3, 17} Reference lists of included studies were also analysed.

We included also trials that investigated regimens, which are currently seen as obsolete, to ensure the most comprehensive overview of the treatment of early colon cancer over three decades. These obsolete regimens often serve as the control arm in newer trials. Furthermore, scoring these older trials might give valuable information regarding the applicability of the scale and identify possible shortcomings. Trials investigating adjuvant treatment regimens that resulted in only negative results were excluded from the analysis. However, a trial with negative results per se, if the regimen investigated had positive outcome in other trials, was not an exclusion criterion. In addition, trials including rectal cancer without a predefined colon cancer subgroup, were excluded from the analysis since radiotherapy is instrumental in (neo) adjuvant rectal cancer treatment which would make it difficult to assess the impact of chemotherapy. Meta-analyses that were not scoreable by the ESMO-MCBS scale were excluded for analysis as well (Consolidated Standards of Reporting Trials (CONSORT) diagram [figure 1](#)).

All studies meeting the inclusion and exclusion criteria were graded using form 1 of the ESMO-MCBS V.1.1 based on OS or DFS results. Additionally, for non-inferiority trials, the grading was influenced by toxicity, quality of

life (QoL) and costs. If there were up to three predefined subgroups included in the trial and there was an appropriate adjustment for multiplicity, these subgroups were graded individually. Trials that did not meet the criteria for scoring due to insufficient benefit (negative studies) have been designated as trials with 'no evaluable benefit' (NEB). Negative non-inferiority (NNI) studies were labelled as NNI. Extracted data and grading were reviewed by the ESMO-MCBS Working Group for accuracy.

An exploratory analysis of reported toxicity data was undertaken to determine the feasibility of toxicity annotations. Side effects during treatment or within 3 months after treatment completion were defined as AT. LT was defined as all events that occurred 3 months after treatment completion in accordance with Common Toxicity Criteria.¹⁸ AT as well as LT was annotated as overall less (-), equal (=) or more (+) toxicity for the intervention versus the control group. When there is insufficient data reported to draw conclusions, not reported (NR) is annotated.

The scores generated in this field testing were reviewed by the ESMO Gastro-Intestinal Tumours Faculty for reasonableness.

RESULTS

The literature search yielded 57 RCTs and 14 meta-analyses, with 18 RCTs and 7 meta-analyses finally found eligible and were included in the analysis. Reasons for exclusion for final analyses are summarised in the CONSORT diagram [figure 1](#) and excluded studies and meta-analyses can be found in the supplementary references.

ESMO-MCBS grading

Information for the selected trials is summarised in [table 1](#) for fluoropyrimidine regimens and in [table 2](#) for oxaliplatin added to fluoropyrimidine regimens. Results in the tables are categorised to combined stage II and III, stage II and stage III.

Fluoropyrimidine regimens

Four trials and a meta-analysis compared 5-FU/LV chemotherapy with MOF combination chemotherapy (lomustine (MeCCNU), vincristine and non-modulated 5-FU)¹⁹ or surgery only for combined stage II and III colon cancers.^{20–23} They showed OS gain ranging from 5% to 14% at 3.0–5.0 years, resulting in the highest-grade ESMO-MCBS grade (A). These results were confirmed in three successive meta-analyses by the Adjuvant Colon Cancer End-points (ACCENT) Group showing a 7.0%–7.2% OS advantage at 5–8 years follow-up (grade A).^{24–26} Since 5-FU/levamisole (LEV) was included in these meta-analyses, the OS benefit of 5-FU/LV was probably underestimated since LEV was subsequently found to be inferior to LV as a 5-FU modulator.^{27–30} [Table 1](#).

The two trials with 318 and 500 patients^{31 32} and a meta-analysis³³ with 1016 patients evaluated adjuvant 5-FU/LV versus no adjuvant therapy in stage II colon cancer. None of these three studies demonstrated OS benefit and all were

annotated as NEB. A 2004 meta-analysis involving 1440 patients²⁵ demonstrated a non-significant 5-year survival gain of 1% (ESMO-MCBS grade NEB), however, a subsequent 2009 evaluation by the same group²⁶ with more mature data reported a 5.4% OS benefit at 8 years (grade A). This discrepancy is addressed in the discussion below.

In grade III colon cancer, the observed OS benefit was 13.5% and 10.3% at 5 and 8 years, respectively, resulting in a grade A on the ESMO-MCBS.^{26 34}

Uracil and tegafur (UFT)/LV in combined stage II and III colon cancer,^{35–37} capecitabine³⁸ and S-1^{39 40} in stage III colon cancer all did not provide an OS or DFS benefit compared with 5-FU/LV. Three of the four trials were non-inferiority trials. Although non-inferiority was proven, since neither QoL nor toxicity was improved; all studies were graded NEB.

Fluoropyrimidines with oxaliplatin combinations

Oxaliplatin added to 5-FU based regimens was evaluated in the MOSAIC⁴¹ and NSABP C-07⁴² trials including stage II and III patients and the NO16968 trial confined to stage III patients.⁴³ Greater clinical benefit (grade A) was observed in the trial confined to stage III compared with the other two trials which were graded B and NEB, respectively. The ACCENT group published a meta-analysis of these studies in 2016.⁴⁴ Based on 5-year OS data their analysis demonstrated an insignificant 0.8% OS gain for stage II (NEB) and a 4.2% OS advantage for stage III colon cancer (grade B). [Table 2](#).

In 2018 the International Duration Evaluation of Adjuvant (IDEA) consortium reported the planned combined analysis of 6 individual RCTs, with a non-inferiority design, comparing folinic acid/5-FU/oxaliplatin (FOLFOX) and capecitabine/oxaliplatin (CAPOX) for 3 vs 6 months.⁴⁵ The 3-year DFS rate was very similar but non-inferiority was not proven for the intention to treat population resulting in a NNI. A preplanned subgroup analysis showed that 3 months CAPOX was non-inferior compared with 6 months. The 3 months treatment arm received a grade B based on non-inferiority in combination with less toxicity.^{46–48} T4 versus T1-3 and N2 versus N1 subgroups were prespecified however their combinations in subgroups and its interaction test was not significant, thus these subgroup analyses were post hoc and could not be graded.

In one meta-analysis, in stage III patients, capecitabine with or without oxaliplatin versus 5-FU/LV with or without oxaliplatin was examined. As no OS and DFS benefit was seen, and neither QoL nor toxicity was improved, it was graded NEB.⁴⁹

Toxicity, QoL and cost

AT and LT reported in the included trials are summarised in the online supplementary table 1. All trials reported AT using several different approaches to toxicity evaluation: one trial did not use any grading system,⁵⁰ five did NR the grading system used,^{19 22 26–29} five used the WHO toxicity scoring system^{20 23 30 33 35} and seven the common terminology



Table 1 ESMO-MCBS grades of fluoropyrimidine treatment for combined stage II and III, stage III and stage II colon cancer

Trial name, year first publication	Intervention versus control	N	Primary outcome	Median follow-up	DFS		OS	Toxicity*	QoL	ESMO-MCBS V1.1	Ref.		
					control group	OS control group							
Fluoropyrimidine treatment versus surgery or no active adjuvant therapy													
Combined stage II and III													
NSABP C-03 1993	5-FU/LV versus MOF	1041	DFS	3 years	64% at 3 years	9%	P=0.0004	77% at 3 years	7%	P=0.003	AT-LT NR	A	19
Siena 1994	5-FU/LV versus surgery	239	DFS	4.5 years	59% at 5 years	15%		65% at 5 years	14%		AT+LT NR	A	20
Meta-analysis IMPACT1995	5-FU/LV versus placebo	1493		5-FU/LV 3.3 yrsPlacebo 3.1 years	62% at 3 years	9%	0.67 (0.56–0.80)	78% at 3 years	5%	0.77 (0.62–0.96)	AT+LT NR	B	21
NCCCTG 874651 1997	5-FU/LV versus placebo	309		6 years	0.58 at 5 years	16%		0.63at 5 years	11%	P=0.02	AT+LT NR	A	22
GIVIO-SITAC 011998	5-FU/LV versus placebo	869		5-FU/LV 5.4 yrsPlacebo 5.3 years	0.54 at 5 years	12%		0.65at 5 years	7%		AT+LT NR	A	23
Meta-analysis Sargent et al. 2001	5-FU/LV or 5-FU/LEV versus surgery alone	3351	OS	5.17–8.54 years	0.58 at 5 years	11%	0.68 (0.60–0.76)	0.64at 5 years	7%	0.76 (0.68–0.85)		A	24
Meta-analysis Gill et al 2004 ²⁵	5-FU/LV or 5-FU/LEV versus surgery	3302	DFS and OS	5 years	0.55 at 5 years	12%	0.7 (0.63–0.78)	0.64 at 5 years	7%	0.74 (0.66–0.83)		A	25
Meta-analysis Sargent et al 2009 ²⁶	5-FU/LV or 5-FU/LEV versus surgery	4922		8 years			NS (HR=0.61)	0.543 at 8 years	7.20%	HR=0.74 P ≤0.001		A	26
Stage II													
Intergroup 00351990	5-FU/LEV versus surgery	318	OS	7 years	71% at 7 years	8%		72% at 7 years	0%		AT+LT NR	NEB	31
Meta-analysis IMPACT B2 1999	5-FU/LV versus placebo	1016	DFS†	5.8 years	73% at 5 years	3%	0.83 (0.68–1.01)	80% at 5 years	2%	0.86 (0.64–1.01)	AT+LT NR	NEB	33
Meta-analysis Gill et al 2004 ²⁵	5-FU/LV or 5-FU/LEV versus surgery	1440	DFS and OS	–	72% at 5 years	4%	p=0.049	80% at 5 years	1%	NS p=0.11		NEB	25
ABCSCG Schippinger et al 2007 ³²	5-FU/LV versus surgery	500	OS	95.6 m (~8 years)	69.4% at 7 years	0.80%	0.95 (0.69–1.31)	76.6% at 7 years	1.60%	0.88 (0.61–1.27)	AT+LT NR	NEB	32
Meta-analysis Sargent et al 2009 ²⁶	5-FU/LV or 5-FU/LEV versus surgery			8 years				66.8% at 8 years	5.40%	P=0.026		A	26
Stage III													
Intergroup 00351990	5-FU/LEV versus surgery alone	619	OS	6.5 years	43.8% at 5 years	17.10%		46.7% at 5 years	13.50%	P ≤0.001	AT+LT?	A	34

Continued

Table 1 Continued

Trial name, year first publication	Intervention versus control	N	Primary outcome	Median follow-up	DFS control group	DFS gain	DFS HR	OS control group	OS gain	OS HR	Toxicity*	QoL	ESMO-MCBS V1.1	Ref.
Meta-analysis Sargent <i>et al</i> 2009 ²⁶	5-FU/LV or 5-FU/LEV versus surgery alone			8 years				42.7% at 8 years	10.30%	$P \leq 0.001$			A	26
Duration of therapy and/or difference in fluoropyrimidine modulator														
Combined stage II and III														
NCCTG-N01C8946511998	5-FU/LEV/LV 12 m versus 5-FU/LEV 12 m	446	OS	5.1 years	63% at 5 years	-6%		68% at 5 years	-5%		AT+LT NR		NEB	27
	5-FU/LEV/LV 6 m versus 5-FU/LEV 12 m	443	OS	5.1 years	63% at 5 years	0%		68% at 5 years	2%		AT-LT NR		NEB	27
	5-FU/LEV 6 m versus 5-FU/LEV 12 m	442	OS	5.1 years	63% at 5 years	-5%		68% at 5 years	-8%		AT-LT NR		NEB	27
NSABP C-041999	5-FU/LV/LEV versus 5-FU/LV	1387	DFS and OS	5 years	65% at 5 years	-1%	NS $p=0.67$	74% at 5 years	-1%	NS $p=0.99$	AT=LT NR		NEB	28
Intergroup 0089 2004	RPMI± versus 5-FU/LEV 12 m	1568	OS	10 years	45% at 10 years	2%		50% at 10 years	2%		AT-LT NR		C	29
	Mayo Clinic§ versus 5-FU/LEV 12 m	1579	OS	10 years	45% at 10 years	4%		50% at 10 years	2%	$P=0.04$	AT-LT NR		C	29
	Mayo Clinic+LEV versus 5-FU/LEV 12 m	1658	OS	10 years	45% at 10 years	23%		50% at 10 years	9%		AT-LT NR		A	29
Stage III														
adjCCCA-012001	5-FU/LV versus 5-FU/LEV	680	OS	82 m (~7 years)	54% at 5 years	8%		60.8% at 5 years	9.20%	$P=0.01$	AT=LT NR		A	30
Convenience of therapy														
Stage II and III														
Kim <i>et al</i> 2003 ³⁵	UFT/LV versus 5-FU/LV	122	DFS and OS	28 m (~3 years)	84.1% at 28 m	3.40%	NS	92.5% at 28 m	2.40%	NS	AT+LT-	=	NEB	35
NSABP-C062006non inferiority trial	UFT/LV versus 5-FU/LV	1551	DFS and OS (margin not clear)	62.3 m (~5 years)	68.2% at 5 years	-1.20%	1 (0.85-1.19)	78.7% at 5 years	-0.20%	1.01 (0.83-1.25)	AT=LT NR	=	NEB	36 37
Stage III														
X-ACT 2005non inferiority trial	cap versus 5-FU/LV	1987	DFS (margin DFS 1.20, OS 1.25)	3.8 years	60.6 at 3 years	3.60%	0.87 (0.75-1.00)	77.6% at 3 years	3.70%	0.84 (0.69-1.01)	AT=LT NR	=	NEB	38
ACTS-CC 2014non-inferiority trial	S-1 versus UFT/LV	1518	DFS (margin DFS 1.29)	41.3 m (~3.5 years)	72.5% at 3 years	3%	0.85 (0.70-1.03)	92.7% at 3 years	0.90%		AT=LT NR		NEB	39 40

Continued



Table 1 Continued

Trial name, year first publication	Intervention versus control		N	Primary outcome	Median follow-up	DFS control group		DFS gain	OS control group		OS gain	OS HR	Toxicity*	QoL	ESMO-MCBS V1.1		Ref.
	control	control				DFS HR	OS HR		MCBS	MCBS							

*All toxicity annotations are exploratory; not part of the latest ESMO-MCBS forms. Toxicity of experimental arm versus control arm, more or less acute toxicity (duration of treatment and thus exposure time to drug and toxicity accounted for as well) of the experimental arm versus the control arm is shown as AT+ or LT+, more or less late toxicity is shown as LT+ or LT-. All toxicity data are summarised in online supplementary table 1. †The endpoint in the IMPACT B2 meta-analysis was EFS, however the given definition 'time from randomisation to first event (ie, either a first recurrence, second tumour, or death from any cause)' is not different from the definition of DFS 'time to any event, irrespective of cause. All events are included, except lost to follow-up'. For better readability DFS is shown in the table instead of EFS. ‡RPMI; 5-FU+LDLV for four courses. §Mayo clinic; 5-FU+LDLV for six courses.

AT, acute toxicity; DFS, disease-free survival; EFS, event free survival; ESMO-MCBS, European Society for Medical Oncology-Magnitude of Clinical Benefit Scale; 5-FU, 5-fluorouracil; HDLV, high dose leucovorin; HR, hazard ratio; LDLV, low dose leucovorin; LEV, levamisole; LT, late toxicity; LV, leucovorin; MOF, lomustine (MeCCNU)/vincristine/5-fluorouracil; N, number of patients; NEB, no evaluable benefit; NR, not reported; NS, not significant; OS, overall survival; QoL, quality of life; RPMI, Roswell Park Memorial Institute; S-1, tegafur/gimeracil/oteracil; UFT, uracil and tegafur; vs, versus.

criteria for adverse events (CTCAE) V.1–3.^{36 38 41 42 44 45} Reporting was more complete in the latest reported studies. The most recent IDEA consortium trial was the most complete in acute adverse events reporting, summarised in online supplementary table 1.⁴⁵ This non-inferiority trial was the only trial in which AT data influenced the grade, as one prespecified subgroup with non-inferior efficacy was rewarded for having less AT to a B grade.

Reporting of LT was very limited. Five trials (two were individual trials within the IDEA collaboration), reported late sensory neuropathy graded with the CTCAE V.1–3.^{41–43 45–47} In all trials, this was investigator reported data and the assessment times and follow-up period differed. Overall, the reported prevalence of late neuropathy was low. With regard to oxaliplatin treatment duration, in the IDEA France trial, at a median follow-up of 3.6 years, the prevalence of grade 3–4 neuropathy was 0.5% among patients exposed to 3 months of oxaliplatin versus 2% among those who received 6 months of oxaliplatin.⁴⁶ In the ACHIEVE trial, at a median follow-up of 3 years, the prevalence all grade neuropathy was 23.3% vs 10%, while grade 3 was only 0.3 vs 0%.⁴⁷ In the ACHIEVE trial, it was also observed that the incidence of any grade PSN was lower for patients treated with CAPOX compared with FOLFOX in both the 6 months and 3 months treatment groups. All other studies did NR any LT.

QoL data were only available for 5 of the 20 RCT (one was an individual trial within the IDEA collaboration).^{23 35–38 48} There was no consistency in the scales used. The only trial to report differences in QoL between the treatment arms was the SCOT trial of the IDEA consortium which compared 3–6 months of oxaliplatin based adjuvant therapy. After 3 months to 5 years of follow-up, there was major difference ($p < 0.001$) in neuropathy-related QoL evaluated using the Functional Assessment of Cancer Therapy/Gynecologic Oncology Group-Neurotoxicity questionnaire. Patients receiving 6 months oxaliplatin reported a worse QoL at 1, 3 and 5 years compared with those receiving 3 months oxaliplatin, and this disparity was associated with major differences in Global QoL between 3 and 6 months gradually attenuating over subsequent months and years.⁴⁸

None of the trials did a formal cost analysis and could therefore not be used for grading of non-inferiority trials. However, non-inferiority of a shorter treatment duration most likely leads to reduction in treatment cost.

Expert peer review of the generated scores

Thirty-seven experts from the ESMO Gastro-Intestinal Tumours Faculty reviewed the generated scores. Twenty-five (67.6%) confirmed the reasonableness of the scores and 12 (34.4%) found the scores mostly reasonable. Experts pointed out that it was striking that the current recommended oxaliplatin-based treatment for stage III disease was only once graded with the highest A grade.

Two experts commented on non-inferiority trials that offer a similar efficacy despite evaluating a more convenient oral mode of administration. They expressed

Table 2 ESMO-MCBS grades of oxaliplatin added to fluoropyrimidine treatment for combined stage II and III, stage III and stage II colon cancer

Trial name, year first publication	Intervention versus control	N	Primary outcome	Median follow-up	DFS control group	DFS gain	DFS HR	OS control group	OS Gain	OS HR	Toxicity*	QoL	ESMO-MCBS V1.1	Ref.
Oxaliplatin added to fluoropyrimidine vs fluoropyrimidine treatment alone														
Stage II and III														
MOSAIC 2004	FOLFOX4 versus 5-FU/LV	2246	DFS	9.5 yrs	61.7% at 10 yrs	5.80%	0.82 (0.71–0.95)	67.1% at 10 yrs	4.60%	0.85 (0.73–0.99)	AT+ LT+		B	41
NSABP C-07 2007	FLOX versus 5-FU/LV	2409	DFS	8 yrs	64.2% at 5 yrs	5.20%	0.82 (0.72–0.93)	78.4% at 5 yrs	1.80%	0.88 (0.75–1.02)	AT+ LT+		NEB	42
Meta-analysis Shah et al 2016 ⁴⁴	5-FU/LV + OX versus 5-FU/LV	6468	OS	6 yrs				77.7% at 5 yrs	2.30%	Significant (HR not presented)			C	44
Stage II														
Meta-analysis Shah et al 2016 ⁴⁴	5-FU/LV + OX versus 5-FU/LV	1600	OS	6 yrs				89.8% at 5 yrs	0.80%	NS			NEB	44
Stage III														
NO16968 2011	CAPOX versus 5-FU/LV	1886	DFS	7 yrs	56% at 7 yrs	7%	0.8 (0.69–0.93)	67% at 7 yrs	6%	0.83 (0.70–0.99)	AT+ LT+		A	43
Meta-analysis Shah et al 2016 ⁴⁴	5-FU/LV + OX versus 5-FU/LV	4868	OS	6 yrs				73.7% at 5 yrs	4.20%	Significant (HR not provided)			B	44
Duration of therapy														
Stage III														
IDEA consortium NCT00958737 2018 non inferiority trial	CAPOX/FOLFOX 3 versus 6 m	12834	DFS (margin 1.12 for DFS)	41.8 m (~3.5 yrs)	75.5% at 3 yrs	-0.90%	1.07 (1.00–1.15)				AT- LT-†	Overall = ‡ PSN - ‡	NNI	45–48
	CAPOX 3 versus 6 m	5071	DFS	74.8% at 3 yrs	1.10%		0.95 (0.85–1.06)				AT- LT-†		B	
	FOLFOX 3 versus 6 m	7763	DFS	76% at 3 yrs	-2.40%		1.16 (1.06–1.26)				AT- LT-		NNI	
Convenience of therapy														
Stage III														
Meta-analysis Schmolli et al 2014 ⁴⁹	CAP ± OX versus 5-FU/LV ± OX	5819	DFS	CAP ± OX 6.2 yrs 5-FU/LV ± OX 5.7 yrs	62.8% at 5 yrs	0%	1.01 (0.92–1.10)	73.9% at 5 yrs	-1.3%	1.02 (0.92–1.14)	AT= LT=		NEB	49

*All toxicity annotations are exploratory; not part of the latest ESMO-MCBS forms. Toxicity of experimental arm versus control arm, more or less acute toxicity (duration of treatment and thus exposure time to drug and toxicity accounted for as well) of the experimental arm versus the control arm is shown as AT+ or AT-, more or less late toxicity is shown as LT+ or LT-. All toxicity data are summarised in Supplementary.

†Late toxicity data were not reported in the IDEA consortium pooled analyses. However, subsequently two of the six individual trials reported late toxicity data^{45,46} which is shown here.

‡QoL data were not reported in the IDEA consortium pooled analyses. However, subsequently one of six individual trials reported QoL data⁴⁷ which is reported here.

AT, acute toxicity; CAP, capecitabine; CAPOX, capecitabine/oxaliplatin; DFS, disease free survival; ESMO-MCBS, European Society for Medical Oncology-Magnitude of Clinical Benefit Scale; FLOX, bolus 5-fluorouracil/leucovorin/oxaliplatin;

FOLFOX, folinic acid/5-fluorouracil/oxaliplatin; 5-FU, 5-fluorouracil; IDEA, International Duration Evaluation of Adjuvant; LT, late toxicity; LV, leucovorin; NEB, no evaluable benefit; NNI, negative non-inferiority; NS, not significant; OS, overall survival; PSN, peripheral sensory neuropathy; QoL, quality of life; yrs, years.



concern that the failure to reward this difference in convenience may fail to credit a true benefit. The ESMO-MCBS V.1.1 of form 1 does not offer the means to credit convenience.

DISCUSSION

This paper has evaluated the applicability of form 1 of the ESMO-MCBS V.1.1 to the adjuvant therapies for early colon cancer. Overall, the experience has been positive insofar as the scoring of adjuvant approaches in early colon cancer are considered reasonable (67.6%) or mostly reasonable (32.4%) by all experts.

For patients with stage III colon cancer, RCTs and meta-analyses of modulated 5-FU-based chemotherapy versus surgery only, consistently scored A in the ESMO-MCBS and the RCTs and meta-analysis comprising oxaliplatin added to this 5-FU backbone showed a modest additional OS benefit (grade B).

For stage II, the findings were less consistent. Whereas fluoropyrimidine trials in patients with stage II colon cancer consistently were graded NEB, the most recent meta-analysis demonstrated a 5.4% survival advantage after 8 years follow-up (grade A). The ACCENT investigators have subsequently cautioned that conclusions derived from older trials of FU-based adjuvant therapy in stage II colon cancer may be biased by stage migration over time.⁵¹ To date, there are no subgroup analyses restricted to stage II in trials with patients that were adequately staged by contemporary standards. RCTs and meta-analysis adding oxaliplatin demonstrated no added benefit for patients with stage II colon cancer.

Several meta-analyses analysed efficacy in stage II/III^{21 24–26 44} as well as separately in II^{25 26 32 44} or stage III.^{26 44 49} Four of these were performed by the ACCENT Collaborative Group^{24–26 44} which, as of 2016, included detailed information collected from over 40 000 patients from 27 adjuvant colon cancer trials including patient demographics and disease characteristics, treatment data, biomarkers for selected studies, adverse events, as well as long term recurrence and survival follow-up for all patients. This has facilitated the capacity to undertake robust analysis of pooled individual patient data in meta-analyses and in the evaluation of the validity of surrogate outcomes.^{52 53}

Regarding the surrogacy of DFS as a predictor of OS, analysis by the ACCENT Collaborative Group demonstrated a robust relationship for 2, 3, 5 and 6 years DFS and OS for stage III colon cancer^{52 53} but this was not the case for stage II disease and indeed even 6 years DFS was only weakly associated with OS.⁵³ Consequently, they concluded that unless DFS is considered a clinically relevant endpoint, OS should be regarded as the most appropriate endpoint for trials in unselected stage II disease.⁵³

The ESMO-MCBS V.1.1 has no defined rules regarding the minimum quality prerequisites for a meta-analysis to be evaluated. In future amendments of the scale, formal definitions of quality and improved clarity regarding the issue of multiplicity when there are several subgroup

analysis will be important. In general, an impactful and valid meta-analysis should include at least the following ingredients: investigation of a plausible question based on randomised evidence using an exhaustive review of relevant studies; evaluation of consistency across studies regarding population of interest, relevant patient characteristics and control arm, coupled with lack of bias (publication, selective reporting); exploration of heterogeneity and clear description of limitations.⁵⁴

Reporting of toxicity and QoL effects of new adjuvant systemic treatment modalities, especially if long-lasting, is important to optimally inform patients. A penalty system for toxicities, such as used in the non-curative setting in the ESMO-MCBS V.1.1 (forms 2 and 3), is not appropriate for the curative setting (form 1) since patients may accept higher toxicity trade-off when treatment is with curative intent. Representatives of patient advocacy groups, in consultation with the ESMO-MCBS Working Group, have indicated preference for annotation of high likelihood of AT or LT versus penalties which may mask the magnitude of curative potential. We strongly believe toxicity annotations should indeed be introduced for treatments with a high prevalence of AT and especially LT.

Our exploratory evaluation of toxicity highlighted that toxicity evaluation and annotation is challenging in the setting of inconsistent methods of toxicity reporting, a high prevalence of apparent under reporting and minimum reporting of LT. The chronic neurotoxicity induced by oxaliplatin is a cumulative, dose-dependent, sensory, symmetric distal axonal neuropathy.^{55 56} Tingling is the most prominent symptom, but numbness and pain can also occur.⁵⁵ In our review of the toxicity data, late grade 3/4 PSN was reported in only 0.5%–2% of patients, substantially lower than the prevalence data derived from patient reported outcome data.⁵⁴ This highlights the risk of under-reporting of toxicities by physicians.⁵⁷ In addition, even several years after adjuvant oxaliplatin-based chemotherapy, in some situations distal neurotoxicity symptoms are reported as re-induced by cold temperature or repeated use of fingers like key-board typing, piano playing or exercising precise finger movements. This is general not mentioned in the toxicity report of clinical trial but has a potential negative impact on QoL or professional career.

In our analyses, only 5 out of 18 trials evaluated QoL.^{23 35–38 48} The low rate of inclusion of QoL evaluation has been examined in a study comprised by phase III RCTs in cancer performed between 2012 and 2016 published in 11 major journals. In 210 of the 446 trials (47.1%), QoL was not included as an endpoint. The non-inclusion was even higher for RCTs in (neo)adjuvant disease as 81 of the 124 trials (65.3%) did not include QoL as an endpoint.⁵⁸ Most of the adjuvant trials reporting QoL showed no difference between the investigational and control arm: 5-FU/LV or placebo,²³ UFT/LV or 5-FU/LV^{35–37} and capecitabine or 5-FU/LV.³⁸ The findings of the SCOT trial⁴⁸ which demonstrated worse QoL for PSN

at 1, 3 and 5 years for patients treated 6 vs 3 months were salient ($p < 0.001$).

UFT/LV did not show OS benefit³⁵ and a non-inferior OS³⁶ for stage II/III colon cancer compared with 5-FU/LV in two trials and neither QoL nor toxicity was improved. Both trials were graded NEB^{35–38} as was the trial of capecitabine versus 5-FU/LV.³⁸ While it is plausible that oral medication may be more convenient than intravenous treatment, there are no data that it actually improves QoL compared with conventional parenteral administration. Convenience is not credited in the current version of form 1 of the ESMO-MCBS.

Our findings confirm that form 1 was highly applicable to the studies of adjuvant systemic therapies of early-stage colon cancer and it provided very reasonable grading for adjuvant colon cancer studies. The exploratory analysis indicated that toxicity annotation is feasible but the prevailing convention of physician reported toxicity may underestimate the true level of patient burden from both AT and LT. Since patients in the curative setting potentially live decades after treatment, late and prolonged adverse effects that may undermine QoL should be annotated to optimally inform patients of recognised risks. Future revisions of form 1 of the ESMO-MCBS will be cognoscente of these findings.

Author affiliations

¹Medical Oncology, University Medical Centre Groningen, Groningen, Groningen, Netherlands

²Medical Oncology, Shaare Zedek Medical Center, Jerusalem, Jerusalem, Israel

³Statistics, Frontier Science Foundation-Hellas, Statistics, Athens, Zografou, Greece

⁴ESMO-MCBS Working Group, European Society for Medical Oncology, Viganella, Switzerland

⁵Nursing, National and Kapodistrian University of Athens, Goudi-Athens, Greece

⁶University of Athens, Athens, Greece

Acknowledgements The authors wish to thank and acknowledge our oncology colleagues who participated in the field testing for 'reasonableness' of scorings using form 1 of the ESMO-MCBS and who have agreed to place their names in this publication (online supplementary table 2). We also thank those who wished to remain anonymous.

Contributors Conception of the work.

Funding ESMO (no grant numbers apply).

Competing interests EGEDV reports Institutional Financial Support for her advisory role from Daiichi Sankyo, Merck, NSABP, Pfizer, Sanofi, Synthon and Institutional Financial Support for clinical trials or contracted research from Amgen, AstraZeneca, Bayer, Chugai Pharma, CytomX Therapeutics, G1 Therapeutics, Genentech, Nordic Nanovector, Radius Health, Regeneron, Roche, Synthon, all outside the submitted work.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as ONLINE supplementary information.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, any changes made are indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Panagiota Zygoura <http://orcid.org/0000-0002-4867-8676>

Derk Jan de Groot <http://orcid.org/0000-0002-8306-6835>

REFERENCES

- GLOBOCAN website. Available: globocan.iarc.fr [Accessed 5 Nov 2018].
- Buyse M, Zeleniuch-Jacquette A, Chalmers TC. Adjuvant therapy of colorectal cancer. Why we still don't know. *JAMA* 1988;259:3571–8.
- Labianca R, Nordlinger B, Beretta GD, et al. Early colon cancer: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2013;24 Suppl 6:vi64–72.
- Saltz LB, Niedzwiecki D, Hollis D, et al. Irinotecan fluorouracil plus leucovorin is not superior to fluorouracil plus leucovorin alone as adjuvant treatment for stage III colon cancer: results of CALGB 89803. *J Clin Oncol* 2007;25:3456–61.
- Van Cutsem E, Labianca R, Bodoky G, et al. Randomized phase III trial comparing biweekly infusional fluorouracil/leucovorin alone or with irinotecan in the adjuvant treatment of stage III colon cancer: PETACC-3. *J Clin Oncol* 2009;27:3117–25.
- Ychou M, Raoul J-L, Douillard J-Y, et al. A phase III randomised trial of LV5FU2 + irinotecan versus LV5FU2 alone in adjuvant high-risk colon cancer (FNCLCC Accord02/FFCD9802). *Ann Oncol* 2009;20:674–80.
- Alberts SR, Sargent DJ, Nair S, et al. Effect of oxaliplatin, fluorouracil, and leucovorin with or without cetuximab on survival among patients with resected stage III colon cancer: a randomized trial. *JAMA* 2012;307:1383–93.
- Taieb J, Balogoun R, Le Malicot K, et al. Adjuvant FOLFOX +/- cetuximab in full Ras and BRAF wildtype stage III colon cancer patients. *Ann Oncol* 2017;28:824–30.
- Allegra CJ, Yothers G, O'Connell MJ, et al. Phase III trial assessing bevacizumab in stages II and III carcinoma of the colon: results of NSABP protocol C-08. *J Clin Oncol* 2011;29:11–16.
- de Gramont A, Van Cutsem E, Schmoll H-J, et al. Bevacizumab plus oxaliplatin-based chemotherapy as adjuvant treatment for colon cancer (AVANT): a phase 3 randomised controlled trial. *Lancet Oncol* 2012;13:1225–33.
- Kerr RS, Love S, Segelov E, et al. Adjuvant capecitabine plus bevacizumab versus capecitabine alone in patients with colorectal cancer (QUASAR 2): an open-label, randomised phase 3 trial. *Lancet Oncol* 2016;17:1543–57.
- Cherny NI, Sullivan R, Dafni U, et al. A standardised, generic, validated approach to stratify the magnitude of clinical benefit that can be anticipated from anti-cancer therapies: the European Society for medical oncology magnitude of clinical benefit scale (ESMO-MCBS). *Ann Oncol* 2015;26:1547–73.
- Cherny NI, Dafni U, Bogaerts J, et al. ESMO-Magnitude of clinical benefit scale version 1.1. *Ann Oncol* 2017;28:2340–66.
- Daniels N. Decisions about access to health care and accountability for Reasonableness. *J Urban Health* 1999;76:176–91.
- Daniels N. Accountability for Reasonableness. *BMJ* 2000;321:1300–1.
- ESMO website. Available: <https://www.esmo.org/guidelines/esmo-mcbs/esmo-magnitude-of-clinical-benefit-scale> [Accessed 5 Nov 2019].
- Benson AB, Venook AP, Al-Hawary MM, et al. NCCN guidelines colon cancer, version 4, 2018. Available: https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf
- Trotti A, Colevas AD, Setser A, et al. CTCAE v3.0: development of a comprehensive grading system for the adverse effects of cancer treatment. *Semin Radiat Oncol* 2003;13:176–81.
- Wolmark N, Rockette H, Fisher B, et al. The benefit of leucovorin-modulated fluorouracil as postoperative adjuvant therapy for primary colon cancer: results from national surgical adjuvant breast and bowel project protocol C-03. *J Clin Oncol* 1993;11:1879–87.
- Francini G, Petrioli R, Lorenzini L, et al. Folinic acid and 5-fluorouracil as adjuvant chemotherapy in colon cancer. *Gastroenterology* 1994;106:899–906.
- Labianca R, Marsoni S, Pancera G, et al. Efficacy of adjuvant fluorouracil and folinic acid in colon cancer. International multicentre pooled analysis of colon cancer trials (impact) Investigators. *Lancet* 1995;345:939–44.
- O'Connell MJ, Mailliard JA, Kahn MJ, et al. Controlled trial of fluorouracil and low-dose leucovorin given for 6 months as postoperative adjuvant therapy for colon cancer. *J Clin Oncol* 1997;15:246–50.
- Zaniboni A, Labianca R, Marsoni S, et al. GIOVIO-SITAC 01: A randomized trial of adjuvant 5-fluorouracil and folinic acid administered to patients with colon carcinoma—long term results and evaluation of the indicators of health-related quality of life. Gruppo



- Italiano Valutazione Interventi in Oncologia. Studio Italiano Terapia Adiuvante Colon. *Cancer* 1998;82:2135–44.
- 24 Sargent DJ, Goldberg RM, Jacobson SD, *et al*. A pooled analysis of adjuvant chemotherapy for resected colon cancer in elderly patients. *N Engl J Med* 2001;345:1091–7.
- 25 Gill S, Loprinzi CL, Sargent DJ, *et al*. Pooled analysis of fluorouracil-based adjuvant therapy for stage II and III colon cancer: who benefits and by how much? *J Clin Oncol* 2004;22:1797–806.
- 26 Sargent D, Sobrero A, Grothey A, *et al*. Evidence for cure by adjuvant therapy in colon cancer: observations based on individual patient data from 20,898 patients on 18 randomized trials. *J Clin Oncol* 2009;27:872–7.
- 27 O'Connell MJ, Laurie JA, Kahn M, *et al*. Prospectively randomized trial of postoperative adjuvant chemotherapy in patients with high-risk colon cancer. *J Clin Oncol* 1998;16:295–300.
- 28 Wolmark N, Rockette H, Mamounas E, *et al*. Clinical trial to assess the relative efficacy of fluorouracil and leucovorin, fluorouracil and levamisole, and fluorouracil, leucovorin, and levamisole in patients with Dukes' B and C carcinoma of the colon: results from national surgical adjuvant breast and bowel project C-04. *J Clin Oncol* 1999;17:3553–9.
- 29 Haller DG, Catalano PJ, Macdonald JS, *et al*. Phase III study of fluorouracil, leucovorin, and levamisole in high-risk stage II and III colon cancer: final report of intergroup 0089. *J Clin Oncol* 2005;23:8671–8.
- 30 Arkenau HT, Bermann A, Rettig K, *et al*. 5-Fluorouracil plus leucovorin is an effective adjuvant chemotherapy in curatively resected stage III colon cancer: long-term follow-up results of the adjCCA-01 trial. *Ann Oncol* 2003;14:395–9.
- 31 Moertel CG, Fleming TR, Macdonald JS, *et al*. Intergroup study of fluorouracil plus levamisole as adjuvant therapy for stage II/Dukes' B2 colon cancer. *J Clin Oncol* 1995;13:2936–43.
- 32 Schippinger W, Samonigg H, Schaberl-Moser R, *et al*. A prospective randomised phase III trial of adjuvant chemotherapy with 5-fluorouracil and leucovorin in patients with stage II colon cancer. *Br J Cancer* 2007;97:1021–7.
- 33 Efficacy of adjuvant fluorouracil and folinic acid in B2 colon cancer. International multicentre pooled analysis of B2 colon cancer trials (impact B2) Investigators. *J Clin Oncol* 1999;17:1356–63.
- 34 Moertel CG, Fleming TR, Macdonald JS, *et al*. Fluorouracil plus levamisole as effective adjuvant therapy after resection of stage III colon carcinoma: a final report. *Ann Intern Med* 1995;122:321–6.
- 35 Kim DJ, Kim TI, Suh JH, *et al*. Oral tegafur-uracil plus folinic acid versus intravenous 5-fluorouracil plus folinic acid as adjuvant chemotherapy of colon cancer. *Yonsei Med J* 2003;44:665–75.
- 36 Lembersky BC, Wieand HS, Petrelli NJ, *et al*. Oral uracil and tegafur plus leucovorin compared with intravenous fluorouracil and leucovorin in stage II and III carcinoma of the colon: results from national surgical adjuvant breast and bowel project protocol C-06. *J Clin Oncol* 2006;24:2059–64.
- 37 Kopec JA, Yothers G, Ganz PA, *et al*. Quality of life in operable colon cancer patients receiving oral compared with intravenous chemotherapy: results from national surgical adjuvant breast and bowel project trial C-06. *J Clin Oncol* 2007;25:424–30.
- 38 Twelves C, Wong A, Nowacki MP, *et al*. Capecitabine as adjuvant treatment for stage III colon cancer. *N Engl J Med* 2005;352:2696–704.
- 39 Yoshida M, Ishiguro M, Ikejiri K, *et al*. S-1 as adjuvant chemotherapy for stage III colon cancer: a randomized phase III study (ACTS-CC trial). *Ann Oncol* 2014;25:1743–9.
- 40 Mochizuki I, Takiuchi H, Ikejiri K, *et al*. Safety of UFT/LV and S-1 as adjuvant therapy for stage III colon cancer in phase III trial: ACTS-CC trial. *Br J Cancer* 2012;106:1268–73.
- 41 André T, de Gramont A, Vernerey D, *et al*. Adjuvant fluorouracil, leucovorin, and oxaliplatin in stage II to III colon cancer: updated 10-year survival and outcomes according to BRAF mutation and mismatch repair status of the mosaic study. *J Clin Oncol* 2015;33:4176–87.
- 42 Yothers G, O'Connell MJ, Allegra CJ, *et al*. Oxaliplatin as adjuvant therapy for colon cancer: updated results of NSABP C-07 trial, including survival and subset analyses. *J Clin Oncol* 2011;29:3768–74.
- 43 Schmoll H-J, Tabernero J, Maroun J, *et al*. Capecitabine plus oxaliplatin compared with fluorouracil/folinic acid as adjuvant therapy for stage III colon cancer: final results of the NO16968 randomized controlled phase III trial. *J Clin Oncol* 2015;33:3733–40.
- 44 Shah MA, Renfro LA, Allegra CJ, *et al*. Impact of patient factors on recurrence risk and time dependency of oxaliplatin benefit in patients with colon cancer: analysis from modern-era adjuvant studies in the adjuvant colon cancer end points (accent) database. *J Clin Oncol* 2016;34:843–53.
- 45 Grothey A, Sobrero AF, Shields AF, *et al*. Duration of adjuvant chemotherapy for stage III colon cancer. *N Engl J Med Overseas Ed* 2018;378:1177–88.
- 46 André T, Vernerey D, Mineur L, *et al*. Three versus 6 months of oxaliplatin-based adjuvant chemotherapy for patients with stage III colon cancer: disease-free survival results from a randomized, open-label, International duration evaluation of adjuvant (idea) France, phase III trial. *J Clin Oncol* 2018;36:1469–77.
- 47 Yoshino T, Yamanaka T, Oki E, *et al*. Efficacy and long-term peripheral sensory neuropathy of 3 vs 6 months of oxaliplatin-based adjuvant chemotherapy for colon cancer: the achieve phase 3 randomized clinical trial. *JAMA Oncol* 2019. doi:10.1001/jamaoncol.2019.2572. [Epub ahead of print: 12 Sep 2019].
- 48 Iveson TJ, Kerr RS, Saunders MP, *et al*. 3 versus 6 months of adjuvant oxaliplatin-fluoropyrimidine combination therapy for colorectal cancer (SCOT): an international, randomised, phase 3, non-inferiority trial. *Lancet Oncol* 2018;19:562–78.
- 49 Schmoll H-J, Twelves C, Sun W, *et al*. Effect of adjuvant capecitabine or fluorouracil, with or without oxaliplatin, on survival outcomes in stage III colon cancer and the effect of oxaliplatin on post-relapse survival: a pooled analysis of individual patient data from four randomised controlled trials. *Lancet Oncol* 2014;15:1481–92.
- 50 Moertel CG, Fleming TR, Macdonald JS, *et al*. Levamisole and fluorouracil for adjuvant therapy of resected colon carcinoma. *N Engl J Med* 1990;322:352–8.
- 51 Shi Q, Andre T, Grothey A, *et al*. Comparison of outcomes after fluorouracil-based adjuvant therapy for stages II and III colon cancer between 1978 to 1995 and 1996 to 2007: evidence of stage migration from the accent database. *J Clin Oncol* 2013;31:3656–63.
- 52 Sargent DJ, Wieand HS, Haller DG, *et al*. Disease-Free survival versus overall survival as a primary end point for adjuvant colon cancer studies: individual patient data from 20,898 patients on 18 randomized trials. *J Clin Oncol* 2005;23:8664–70.
- 53 Sargent D, Shi Q, Yothers G, *et al*. Two or three year disease-free survival (DFS) as a primary end-point in stage III adjuvant colon cancer trials with fluoropyrimidines with or without oxaliplatin or irinotecan: data from 12,676 patients from mosaic, X-ACT, PETACC-3, C-06, C-07 and C89803. *Eur J Cancer* 2011;47:990–6.
- 54 Moher D, Liberati A, Tetzlaff J, *et al*. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- 55 Pachman DR, Qin R, Seisler DK, *et al*. Clinical course of oxaliplatin-induced neuropathy: results from the randomized phase III trial N08CB (Alliance). *J Clin Oncol* 2015;33:3416–22.
- 56 Park SB, Lin CSY, Krishnan AV, *et al*. Long-Term neuropathy after oxaliplatin treatment: challenging the dictum of reversibility. *Oncologist* 2011;16:708–16.
- 57 Di Maio M, Basch E, Bryce J, *et al*. Patient-Reported outcomes in the evaluation of toxicity of anticancer treatments. *Nat Rev Clin Oncol* 2016;13:319–25.
- 58 Marandino L, La Salvia A, Sonetto C, *et al*. Deficiencies in health-related quality-of-life assessment and reporting: a systematic review of oncology randomized phase III trials published between 2012 and 2016. *Ann Oncol* 2018;29:2288–95.