

University of Groningen

## Heterologous expression and secretion of Clostridium perfringens [beta]-toxoid by Bacillus subtilis

Nijland, Reindert

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*  
2007

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Nijland, R. (2007). Heterologous expression and secretion of Clostridium perfringens [beta]-toxoid by Bacillus subtilis. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# List of publications

- Nijland R**, Veening JW, and Kuipers OP (2007) A de-repression system based on the *Bacillus subtilis* sporulation pathway offers dynamic control of heterologous gene expression. *Appl. Environ. Microbiol.*, published ahead of print on 9 February 2007, doi:10.1128/AEM.02266-06 - *RN and JWV contributed equally to this work*
- Nijland R**, Heerlien R, Hamoen LW, and Kuipers OP (2007) Changing a single amino acid in *Clostridium perfringens* {beta}-toxin affects efficiency of heterologous secretion by *Bacillus subtilis*. *Appl. Environ. Microbiol.*, 73(5): 1586-1593
- Nijland R**, Lindner C, van Hartskamp M, Hamoen LW, and Kuipers OP (2007) Heterologous production and secretion of *Clostridium perfringens* [beta]-toxin in closely related Gram-positive hosts. *Journal of Biotechnology*, 127, 361-372
- Gravato-Nobre MJ, Nicholas HR, **Nijland R**, O'Rourke D, Whittington DE, Yook KJ, and Hodgkin J (2005) Multiple genes affect sensitivity of *Caenorhabditis elegans* to the bacterial pathogen *Microbacterium nematophilum*. *Genetics*, 171, 1033-1045.
- Lindner C, **Nijland R**, van Hartskamp M, Bron S, Hamoen LW, and Kuipers OP (2004) Differential expression of two paralogous genes of *Bacillus subtilis* encoding single-stranded DNA binding protein. *J Bacteriol*, 186, 1097-1105. *CL and RN contributed equally to this work*
- Jongbloed JD, Grieger U, Antelmann H, Hecker M, **Nijland R**, Bron S, and van Dijl JM (2004) Two minimal Tat translocases in *Bacillus*. *Mol Microbiol*, 54, 1319-1325.
- Jongbloed JD, Antelmann H, Hecker M, **Nijland R**, Bron S, Airaksinen U, Pries F, Quax WJ, van Dijl JM, and Braun PG (2002) Selective contribution of the twin-arginine translocation pathway to protein secretion in *Bacillus subtilis*. *J Biol Chem*, 277, 44068-44078.

