

## University of Groningen

### Correction to

Santema, Bernadet T.; Chan, Michelle M. Y.; Tromp, Jasper; Dokter, Martin; van der Wal, Haye H.; Emmens, Johanna E.; Takens, Janny; Samani, Nilesh J.; Ng, Leong L.; Lang, Chim C.

*Published in:*  
Clinical Research in Cardiology

*DOI:*  
[10.1007/s00392-020-01762-2](https://doi.org/10.1007/s00392-020-01762-2)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2020

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Santema, B. T., Chan, M. M. Y., Tromp, J., Dokter, M., van der Wal, H. H., Emmens, J. E., Takens, J., Samani, N. J., Ng, L. L., Lang, C. C., van der Meer, P., ter Maaten, J. M., Damman, K., Dickstein, K., Cleland, J. G., Zannad, F., Anker, S. D., Metra, M., van der Harst, P., ... Voors, A. A. (2020). Correction to: The influence of atrial fibrillation on the levels of NT-proBNP versus GDF-15 in patients with heart failure. *Clinical Research in Cardiology*. <https://doi.org/10.1007/s00392-020-01762-2>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*



## Correction to: The influence of atrial fibrillation on the levels of NT-proBNP versus GDF-15 in patients with heart failure

Bernadet T. Santema<sup>1</sup> · Michelle M. Y. Chan<sup>2</sup> · Jasper Tromp<sup>1,2</sup> · Martin Dokter<sup>1</sup> · Haye H. van der Wal<sup>1</sup> · Johanna E. Emmens<sup>1</sup> · Janny Takens<sup>1</sup> · Nilesh J. Samani<sup>3,4</sup> · Leong L. Ng<sup>3,4</sup> · Chim C. Lang<sup>5</sup> · Peter van der Meer<sup>1</sup> · Jozine M. ter Maaten<sup>1</sup> · Kevin Damman<sup>1</sup> · Kenneth Dickstein<sup>6,7</sup> · John G. Cleland<sup>8,9</sup> · Faiez Zannad<sup>10</sup> · Stefan D. Anker<sup>11</sup> · Marco Metra<sup>12</sup> · Pim van der Harst<sup>1</sup> · Rudolf A. de Boer<sup>1</sup> · Dirk J. van Veldhuisen<sup>1</sup> · Michiel Rienstra<sup>1</sup> · Carolyn S. P. Lam<sup>1,2</sup> · Adriaan A. Voors<sup>1</sup> 

© The Author(s) 2020

### Correction to:

**Clinical Research in Cardiology (2020) 109:331–338**  
<https://doi.org/10.1007/s00392-019-01513-y>

The original version of this article unfortunately contained a mistake. Section “Results”, second paragraph, first sentence should read:

The plasma levels of NT-proBNP were significantly higher in patients who had AF at baseline, with a median of 3417 pg/mL (1897–6486), as compared with patients who were in SR at baseline; both those who had a history of AF (1788 pg/mL [682–3870],  $p < 0.001$ ) and those who never had AF before (2231 pg/mL [902–5270],  $p < 0.001$ ) (Table 1), also after multivariable adjustment (Table 2).

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

The original article can be found online at <https://doi.org/10.1007/s00392-019-01513-y>.

✉ Adriaan A. Voors  
a.a.voors@umcg.nl

<sup>1</sup> Department of Cardiology, University of Groningen, University Medical Centre Groningen, Hanzplein 1, 9713 GZ Groningen, The Netherlands

<sup>2</sup> Department of Cardiology, National Heart Centre Singapore, Singapore Duke-NUS Graduate Medical School, Singapore, Singapore

<sup>3</sup> Department of Cardiovascular Sciences, University of Leicester, Groby Road, Leicester LE3 9QP, UK

<sup>4</sup> NIHR Leicester Biomedical Research Centre, Glenfield Hospital, Groby Road, Leicester LE3 9QP, UK

<sup>5</sup> Division of Molecular and Clinical Medicine, School of Medicine University of Dundee, Ninewells Hospital and Medical School, Dundee DD1 9SY, UK

<sup>6</sup> University of Bergen, Bergen, Norway

<sup>7</sup> Stavanger University Hospital, Stavanger, Norway

<sup>8</sup> National Heart & Lung Institute, Royal Brompton & Harefield Hospitals, Imperial College, Sydney St, Chelsea, London SW3 6NP, UK

<sup>9</sup> Robertson Institute of Biostatistics and Clinical Trials Unit, University of Glasgow, University Avenue, Glasgow G12 8QQ, UK

<sup>10</sup> Inserm CIC 1433, Université de Lorraine, CHU de Nancy, Nancy, France

<sup>11</sup> Department of Cardiology (CVK), and Berlin Institute of Health Center for Regenerative Therapies (BCRT), German Centre for Cardiovascular Research (DZHK) partner site Berlin, Charité Universitätsmedizin, Berlin, Germany

<sup>12</sup> Department of Medical and Surgical Specialties, Radiological Sciences and Public Health, Institute of Cardiology, University of Brescia, Brescia, Italy