University of Groningen

Oxidative Methylation of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiX}$ ($\text{X} = \text{Cl}, \text{OMe}, \text{Me}$) with Group 12 Dimethyl Complexes to Diamagnetic $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiX}(\text{Me})$

Luinstra, Gerrit A.; Teuben, Jan H.

Published in:
Journal of Organometallic Chemistry

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1991

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Luinstra, G. A., & Teuben, J. H. (1991). Oxidative Methylation of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiX}$ ($\text{X} = \text{Cl}, \text{OMe}, \text{Me}$) with Group 12 Dimethyl Complexes to Diamagnetic $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiX}(\text{Me})$. *Journal of Organometallic Chemistry*, 420(3).

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Preliminary Communication

Recl. Trav. Chim. Pays-Bas **110**, 57-58 (1991)

0165-0513/91/02057-02\$1.00

OXIDATIVE METHYLATION OF $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiX}$ ($\text{X} = \text{Cl, OMe, Me}$) WITH GROUP 12 DIMETHYL COMPLEXES TO DIAMAGNETIC $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiX(Me)}$

Gerrit A. Luinstra and Jan H. Teuben*, Groningen Center for Catalysis and Synthesis, Department of Chemistry, Nijenborgh 16, University of Groningen, 9747 AG Groningen, The Netherlands.
(Received January 2nd, 1991)

Abstract. Tervalent titanium compounds $(\text{C}_5\text{Me}_5)_2\text{TiX}$ ($\text{X} = \text{OMe, Cl, Me}$) are selectively oxidized and alkylated at room temperature by ZnMe_2 to diamagnetic monomethyl derivatives $(\text{C}_5\text{Me}_5)_2\text{TiX(Me)}$.

Recently we reported the clean oxidation of tervalent titanium compounds $(\text{C}_5\text{Me}_5)_2\text{TiR}$ ($\text{R} = \text{alkyl, aryl, H, OR etc}$) to diamagnetic $(\text{C}_5\text{Me}_5)_2\text{TiR(Cl)}$ complexes by PbCl_2 ¹. This reaction is very useful since consecutive halide substitution with alkyl alkali metal reagents gives mixed complexes $(\text{C}_5\text{Me}_5)_2\text{TiR}^1\text{R}^2$. We now have developed a direct route to complexes $(\text{C}_5\text{Me}_5)_2\text{TiX(Me)}$ ($\text{X} = \text{OMe, Cl, Me}$), starting from $(\text{C}_5\text{Me}_5)_2\text{TiX}$ by oxidative alkylation with ZnMe_2 .

Analogous to the clean oxidation of $(\text{C}_5\text{Me}_5)_2\text{TiX}$ derivatives with PbCl_2 , the synthetic strategy is based on simultaneous oxidation of the metal and introduction of an alkyl group, using homoleptic metal alkyls (eq 1).

bond in $(\text{C}_5\text{Me}_5)_2\text{TiX(Me)}$ is stronger than the mean M-C bond strength in MMe_n . Literature data on the titanium methyl bond indicate a bond dissociation energy of approximately 280 kJ/mol², making the readily available homoleptic group 12 complexes with mean bond dissociation energies of 126 kJ/mol (Hg) to 184 kJ/mol (Zn) particularly suitable³.

Dimethyl mercury was not particularly reactive towards $(\text{C}_5\text{Me}_5)_2\text{TiX}$ possibly for kinetic reasons, although oxidative alkylations are known for the former compound (e.g. reaction with lithium metal⁴). In contrast ZnMe_2 was found to react stoichiometrically with $(\text{C}_5\text{Me}_5)_2\text{TiX}$ ($\text{X} = \text{OMe, Cl, Me}$) to form $(\text{C}_5\text{Me}_5)_2\text{TiX(Me)}$ and zinc metal[†]. To the best of our knowledge, the oxidizing capacity

$\text{X} = \text{OMe (1), Cl (2), Me (3)}$

The thermodynamic boundary conditions which this reaction has to meet is that the titanium carbon

of zinc compounds was unknown prior to this work. Zinc alkyls have been widely used as non reductive

substitution reagents⁵, but reduction to zinc metal has not been reported before.

The oxidative alkylation proceeds smoothly in toluene at room temperature and reactions usually are complete within 24h. Remarkably, the oxidative alkylation is kinetically favoured over substitution. Reaction of $(\text{C}_5\text{Me}_5)_2\text{TiCl}$ with half a mole of ZnMe_2 leads to selective formation of $(\text{C}_5\text{Me}_5)_2\text{TiCl}(\text{Me})$ (**2**). Neither products originating from salt metathesis, like $(\text{C}_5\text{Me}_5)_2\text{TiMe}$, nor its consecutive oxidation alkylation product $(\text{C}_5\text{Me}_5)_2\text{TiMe}_2$ (**3**) (*vide infra*) is present in the reaction mixture⁶. In addition, (**2**) is not reduced by zinc metal formed *in situ*, although zinc is reported to be capable of reducing tetravalent titanium halides⁷.

Treatment of $(\text{C}_5\text{Me}_5)_2\text{TiMe}$ with half a mole of ZnMe_2 at room temperature quantitatively yields (**3**) within 2h. However, reactions of ZnMe_2 with other $(\text{C}_5\text{Me}_5)_2\text{TiR}$ ($R \neq \text{Me}$) are complicated by extensive disproportionation. This presumably results from rapid alkyl exchange between trivalent $(\text{C}_5\text{Me}_5)_2\text{TiR}$ and ZnMe_2 , followed by irreversible oxidation to a mixture of tetravalent $(\text{C}_5\text{Me}_5)_2\text{TiR}(\text{Me})$, $(\text{C}_5\text{Me}_5)_2\text{TiMe}_2$ and $(\text{C}_5\text{Me}_5)_2\text{TiR}_2$ species. For example, reaction of $(\text{C}_5\text{Me}_5)_2\text{TiPh}$ with ZnMe_2 results in the isolation of a mixture of (**3**), $(\text{C}_5\text{Me}_5)_2\text{TiPh}(\text{Me})$ ⁸ and $(\text{C}_5\text{Me}_5)_2\text{TiPh}_2$ ⁹. The oxidative alkylation strategy for the synthesis of pure $(\text{C}_5\text{Me}_5)_2\text{TiR}^1\text{R}^2$ from $(\text{C}_5\text{Me}_5)_2\text{TiR}^1$ seems to be restricted to the case where $\text{R}^1 = \text{R}^2 = \text{Me}$.

We are currently investigating the scope of oxidative alkylations, especially for the preparation of pure mixed $(\text{C}_5\text{Me}_5)_2\text{TiR}^1\text{R}^2$ compounds. The method of introducing alkyl groups at low-valent (early) transition metal center through the oxidative alkylation pathway reported here is a promising alternative for the preparation of alkyl complexes.

† Reaction of vanadocene with dialkylcadmium complexes has been reported to give Cp_2VR_2 . We observed that compounds $(\text{C}_5\text{Me}_5)_2\text{TiX}$ are also oxidatively alkylated by CdR_2 . However, the use of CdR_2 has some disadvantages over ZnR_2 owing to the photosensitivity of the former compounds. *G.A. Razuvaev, V.N. Latyaeva, L.I. Vishinskaya, V.K. Cherkasov, S.P. Korneva, N.N. Spiridonova, J. Organomet. Chem.* **129**, 169 (1977).

References.

1. *G.A. Luinstra and J.H. Teuben*, *J. Chem. Soc., Chem. Commun.* 1470 (1990).
2. *A. Dias, M.S. Salema, J.A. Martinho Simões, J.W. Pattiasina and J.H. Teuben*, *J. Organomet. Chem.* **346**, C4 (1988),
3. *H.A. Skinner*, *Adv. Organomet. Chem.* **2**, 49 (1964).
4. *W. Schlenk and J. Holtz*, *Chem. Ber.* **50**, 271 (1917).
- 5a. *J. Boersma*, in "Comprehensive Organometallic Chemistry", Vol. 2, *G. Wilkinson, F.G.A. Stone, E.W. Abel*, eds., Pergamon, Oxford, 1982, p. 823.
- b. *R.R. Schrock*, *Acc. Chem. Res.* **12**, 98 (1979).
6. *J.E. Bercaw and H.H. Brintzinger*, *J. Am. Chem. Soc.* **93**, 2045 (1971).
7. C.f. *M. Bottrill, P.D. Gavens, J.W. Kelland and J. McMeeking*, in "Comprehensive Organometallic Chemistry", Vol. 3, Chapter 22.2, *G. Wilkinson, F.G.A. Stone, E.W. Abel*, eds., Pergamon, Oxford, 1982.
8. *G.A. Luinstra and J.H. Teuben*, manuscript in preparation.
9. *H.S. Tung and C.H. Brubaker Jr.* *Inorg. Chim. Acta.* **52**, 197 (1981).