

University of Groningen

Heritability of telomere length in the Zebra Finch

Atema, Els; Mulder, Ellis; Dugdale, Hannah L.; Briga, Michael; van Noordwijk, Arie J.; Verhulst, Simon

Published in:
Journal of Ornithology

DOI:
[10.1007/s10336-015-1212-7](https://doi.org/10.1007/s10336-015-1212-7)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Atema, E., Mulder, E., Dugdale, H. L., Briga, M., van Noordwijk, A. J., & Verhulst, S. (2015). Heritability of telomere length in the Zebra Finch. *Journal of Ornithology*, *156*(4), 1113-1123.
<https://doi.org/10.1007/s10336-015-1212-7>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Heritability of telomere length in the Zebra Finch

Els Atema^{1,2} · Ellis Mulder¹ · Hannah L. Dugdale³ · Michael Briga¹ ·
Arie J. van Noordwijk² · Simon Verhulst¹

Received: 28 January 2015 / Revised: 12 March 2015 / Accepted: 16 March 2015
© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract Telomere length predicts survival in birds, and many stressors that presumably reduce fitness have also been linked to telomere length. The response to selection of telomere length will be largely determined by the heritability of this trait; however, little is known about the genetic component of telomere length variation in animals other than humans. Moreover, published heritability estimates of telomere length are based on telomere measurements with techniques that do not distinguish between terminal telomeres, which are susceptible to age and stress, and the interstitial telomeric repeats, which are relatively inert. Heritability estimates that combine interstitial and terminal telomeres are difficult to interpret in species such as birds, where interstitial telomeres are often numerous. We estimated the heritability of terminal telomere length in a captive Zebra Finch population of cross-fostered (half-)siblings using data obtained with an electrophoresis technique that excludes the interstitial repeats from the

measurements. We used both a Bayesian quantitative genetic ‘animal’ model and a frequentist sibling regression approach to estimate heritability. With the animal model, we estimated a high heritability of telomere length ($h^2 = 0.99$, 95 % credible interval = 0.87–1), but had insufficient statistical power to separate parental and permanent environment effects. The frequentist approach yielded similar heritability estimates, although with large confidence intervals. We used general linear mixed models to disentangle variance components of telomere length. The relative contributions of the individual, mother and father to telomere length variation were statistically indistinguishable at 23–31 %. Chicks were cross-fostered 4-days after hatching, and no effect of rearing nest was found, indicating that any undetected environmental effects exerted their influence prior to, or soon after, hatching. Thus, we conclude that telomere length resemblance between relatives is high and proportional to their relatedness, but we cannot conclusively distinguish between genetic and other forms of inheritance.

Communicated by M. Wink.

Electronic supplementary material The online version of this article (doi:10.1007/s10336-015-1212-7) contains supplementary material, which is available to authorized users.

✉ Els Atema
e.atema@rug.nl

¹ Department of Behavioural Biology, Groningen Institute of Evolutionary Life Sciences, University of Groningen, P.O. Box 11103, 9700 CC Groningen, The Netherlands

² Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), P.O. Box 50, 6700 AB Wageningen, The Netherlands

³ Department of Animal and Plant Sciences, Alfred Denny Building, University of Sheffield, Western Bank, Sheffield S10 2TN, UK

Keywords TRF · Avian · (Half-)siblings · Cross-fostered · Natal environment · ‘Animal’ model

Zusammenfassung

Erblichkeit der Telomerlänge beim Zebrafinch *Taeniopygia guttata*

Anhand der Telomerlänge sind bei Vögeln Prognosen zur Überlebensdauer möglich, und viele Stressfaktoren, welche vermutlich die Fitness verringern, konnten ebenfalls mit der Telomerlänge in Verbindung gebracht werden. Wie die Telomerlänge auf Selektion reagiert, wird hauptsächlich von der Erblichkeit dieses Merkmales abhängen, allerdings

weiß man mit Ausnahme des Menschen bislang nur wenig über die genetische Komponente der Telomerlängen-Variabilität bei Tieren. Darüber hinaus basieren veröffentlichte Schätzwerte für die Erbllichkeit der Telomerlänge häufig auf Messwerten, welche mit Techniken erhoben wurden, die nicht zwischen terminalen Telomeren, welche anfällig für Alter und Stress sind, und den relativ inerten interstitiellen telomerischen Sequenzen unterscheiden. Schätzwerte für die Erbllichkeit, die interstitielle und terminale Telomere zusammenfassen, sind bei solchen Arten wie den Vögeln, bei denen interstitielle Telomere oft zahlreich vorhanden sind, schwer zu interpretieren. Wir schätzten die Erbllichkeit der terminalen Telomerlänge bei einer Volierenpopulation von Zebrafincken- (Halb-)geschwistern, die in Pflegenestern aufgezogen worden waren, anhand von Daten, die mit einer Elektrophorese-technik gewonnen wurden, welche die interstitiellen Wiederholungselemente von den Messungen ausschloss. Zur Schätzung der Erbllichkeit verwendeten wir sowohl ein Bayes'sches quantitatives genetisches Tiermodell als auch einen frequentistischen Geschwister-Regressionsansatz. Für das Tiermodell ergaben sich hohe Schätzwerte für die Erbllichkeit der Telomerlänge ($h^2 = 0.99$, 95 %-Kreditintervall = 0.87–1), dieses hatte allerdings nur eine unzureichende statistische Aussagekraft bei der Trennung von elterlichen Faktoren und ständigen Umwelteinflüssen. Der frequentistische Ansatz ergab ähnlich Erbllichkeitsschätzwerte, jedoch mit großen Konfidenzintervallen. Wir verwendeten gemischte lineare Modelle (General Linear Mixed Models, GLMM), um die Varianzkomponenten der Telomerlänge aufzuschlüsseln. Der relative Anteil, den das Individuum beziehungsweise dessen Mutter und Vater zur Telomerlängen-Variabilität beitrugen, ließ sich statistisch nicht unterscheiden und lag zwischen 23–31 %. Vier Tage nach dem Schlüpfen wurden die Küken gegen solche aus einem anderen Nest ausgetauscht, es war aber kein Effekt des Aufzuges festzustellen, was darauf hindeutet, dass etwaige unbemerkte Umweltfaktoren vor oder kurz nach dem Schlupf zur Wirkung kommen. Wir schlussfolgern daher, dass die Ähnlichkeit der Telomerlänge zwischen Verwandten hoch ist und in Proportion zum Verwandtschaftsgrad steht; wir können aber nicht mit Sicherheit zwischen genetischer und anderen Formen der Erbllichkeit unterscheiden.

Introduction

Telomeres are non-coding repeats of the highly conserved DNA sequence 5'-TTAGGG-3' (Meyne et al. 1989) that are important in the protection and stabilization of linear chromosomes (Blackburn 1991). Because of the end

replication problem, the fact that DNA polymerase is not able to completely replicate the 3' end of the DNA strand, and other contributing factors such as oxidative stress, telomeres tend to shorten with age (Olovnikov 1973; von Zglinicki 2002). Telomere shortening can be accelerated by various forms of stress encountered throughout life (Epel et al. 2004; Kotrschal et al. 2007; Gilley et al. 2008; Bauch et al. 2013; Boonekamp et al. 2014). Short telomeres eventually lead to replicative senescence of the cell (Blackburn 2005), and individuals with longer telomeres have a higher probability of survival in numerous species (Joeng et al. 2004; Haussmann et al. 2005; Bize et al. 2009; Salomons et al. 2009), including humans (Boonekamp et al. 2013). Furthermore, individual Zebra Finches with longer telomeres at the end of the nestling period have a longer lifespan (Heidinger et al. 2012). Therefore, telomeres have been suggested as biomarkers of the levels of stress that individuals have experienced over their lifetime (e.g. Monaghan 2014).

Given the association between telomere length and survival, it is of interest to unravel the causes of variation in telomere length. Variation in telomere length is determined by (1) initial telomere length in the zygote, (2) the rate of shortening and (3) the degree of telomere maintenance (Bischoff et al. 2005), and all three factors may be influenced by both genetic and environmental factors. Quantifying the contribution of genetics to the outcome of these processes (i.e., heritability) is important because it will determine the response to selection (Falconer and Mackay 1996; Visscher et al. 2008). New statistical methods also enable separation of individual phenotypes into genetic and environmental components (Kruuk and Hadfield 2007), allowing estimation of the importance of various factors in shaping the variation in a trait. Heritability of human telomere length has repeatedly been investigated, and heritability estimates are in the range of 0.36 to 1.28 (Table 1). Based on the largest data set, the heritability of telomere length was estimated at 0.70 (95 % CI 0.64–0.76, Broer et al. 2013), which falls close to the mean of the other estimates. In five animal populations, heritability of telomere length was estimated to range from 0.09 to 1 (Table 1). The number of studies is low, however, and they differ widely in many respects (e.g., in estimate and measurement technique). Hence, our knowledge of the heritability of telomere length in non-human animals is limited.

In the present paper, we report heritability estimates of telomere length in a population of captive Zebra Finches. Resemblance between relatives can have a genetic basis, but can also be due to a shared environment. In order to obtain an unbiased estimate of heritability, genetic and environmental effects need to be separated (Kruuk and Hadfield 2007). To reduce shared environment effects, we cross-fostered 75 % of the individuals 4 days after the first

Table 1 Published estimates of telomere length heritability

Species	Telomere analysis	Relationship	Sample size	h^2 (95 % CI)	Parental effects	References
Human <i>Homo sapiens</i>	Southern blot	Twin	115	0.78 (0.69–0.87)	Not reported	Slagboom et al. (1994)
Human <i>Homo sapiens</i>	Southern blot	Twin	98	0.84 (0.73–0.91)	Not reported	Jeanclos et al. (2000)
Human <i>Homo sapiens</i>	Southern blot	Family-based	327	0.99	Not reported	Nawrot et al. (2004)
Human <i>Homo sapiens</i>	Southern blot	Twin	574	0.36 (0.22–0.48)	Not reported	Bischoff et al. (2005)
Human <i>Homo sapiens</i>	Southern blot	Sibling	383	0.82 (0.59–1.05)	Not reported	Vasa-Nicotera et al. (2005)
Human <i>Homo sapiens</i>	Southern blot	Twin	2050	0.36 (0.18–0.48)	Not reported	Andrew et al. (2006)
Human <i>Homo sapiens</i>	qPCR	Family-based	907	0.44 (0.32–0.56)	Paternal	Njajou et al. (2007)
Human <i>Homo sapiens</i>	qPCR	Mother-offspring	129	0.30 (– 0.05–0.62)	Paternal	Nordfjäll et al. (2010)
Human <i>Homo sapiens</i>	qPCR	Father-offspring	98	0.91 (0.56–1.20)	(Paternal)	Al-Attas et al. (2012)
Human <i>Homo sapiens</i>	qPCR	Parent-offspring	254	1.28	(Paternal)	Al-Attas et al. (2012)
Human <i>Homo sapiens</i>	qPCR	Meta-analysis	19,713	0.70 (0.64–0.76)	Maternal	Broer et al. (2013)
Human <i>Homo sapiens</i>	Southern blot	(6 different populations)			paternal age	
Sand Lizard <i>Lacerta agilis</i>	Southern blot	Mother-daughter	55	0.52 (0.09–0.95)	Paternal	Olsson et al. (2011)
		Father-son	40	1.23 (0.81–1.67)		
Kakapo <i>Strigopshabroptilus</i>	Southern blot	Mother-offspring	29 pairs	0.84	(Maternal)	Horn et al. (2011)
Collared Flycatcher <i>Ficedula albicollis</i>	qPCR	Siblings (cross-fostering)	74 broods	0.09 (–0.04–0.15)	Not reported	Voillemot et al. (2012)
King Penguin <i>Aptenodytes patagonicus</i>	qPCR	Mid-parent-offspring	53 breeding pairs	0.2 (–0.02–0.42)	(Maternal)	Reichert et al. (2014)
Great Reed Warbler <i>Acrocephalus arundinaceus</i>	qPCR	‘Animal’ model	193	0.48 (0.25–0.72)	Maternal	Asghar et al. (2015)
		Mother-mid-offspring		1.08		
		Father-mid-offspring		0.28		
Zebra Finch <i>Taeniopygia guttata</i>	TRF	‘Animal’ model (half-)Siblings (cross-fostering)	125	0.999 (0.87–1) Details: Table 4	None	This study

The sample size is the number of phenotyped individuals in the analyses, unless otherwise specified. Parental effects between brackets are not significant, or significance was not tested

Missing CI in the table were not reported

h^2 Heritability, 95 % CI 95 % confidence interval

hatching in a nest, enabling us to separate birth and rear nest effects, at least from cross-fostering onwards. Furthermore, we used a sample containing both full siblings and half-siblings to separate covariance of different

degrees of relatives statistically. Specifically, we included both maternal and paternal half-siblings to quantify parental effects. We estimated heritability with two methods. First we applied a so-called ‘animal’ model, a general

linear mixed model using a pedigree of individuals in the data set, to estimate heritability (Kruuk 2004) in a Bayesian framework (Hadfield 2010). ‘Animal’ models have the advantage of using all relationships in the data set, but the disadvantage of being data consuming. Second, we used frequentist methods to calculate intraclass correlations and heritabilities comparing both full and half sibships to investigate parental effects (Falconer and Mackay 1996). Finally, we disentangled environmental and parental effects on telomere length with general linear mixed models (GLMM).

In contrast to humans, many avian species have numerous interstitial telomeric repeats, which are in addition to ‘terminal’ telomeric repeats (Delany et al. 2000; Foote et al. 2013). Terminal telomeres are susceptible to ageing and environmental factors and are involved in the protection and stabilization of the chromosomes. Although there is evidence that interstitial telomeric repeats are involved in DNA repair, chromosome stabilization and the regulation of gene transcription, the exact function is not yet fully understood (Kilburn et al. 2001; Rivero et al. 2004; Yang et al. 2011). It is also not known whether interstitial telomeric repeats change in length within an animal’s lifetime, given that this would involve two double strand breaks. The number of interstitial telomeric repeats and the length of terminal telomeres are therefore in essence different traits, despite their superficial similarity. Laboratory techniques differ in the type of telomeres included in the measurements, and published heritability estimates are based on techniques that pool interstitial and terminal telomeric repeats in one estimate (qPCR, southern blot; Table 1). Thus, it is not known to what extent published heritability estimates in species other than humans provide information on variation in interstitial versus terminal telomeres (there are few interstitial repeats in the human genome). We therefore measured telomeres with in-gel hybridization, labelling the single-stranded overhang of telomeres (Haussmann and Vleck 2002), i.e., the telomeric loop at the end of linear chromosomes that is an evolutionary, well-conserved aspect of telomere biology (Stansel et al. 2001). Hence, we measure only terminal telomeres (see “Materials and methods” for details), and our heritability estimates are specifically for terminal telomeres only.

Materials and methods

Study species and sampling

We used 125 Zebra Finches (66 females; 57 males), originating from 73 broods, reared from stock that were part of a long-term experiment, in which natal brood size

and energy expenditure required for foraging were manipulated (De Coster et al. 2011; Koetsier and Verhulst 2011). With respect to the foraging cost manipulation, we only used control birds that had easy access to food. Parents were paired randomly and housed in pairs during breeding; hence, paternity was known with certainty. The inbreeding level is low in our Groningen Zebra Finch population (Forstmeier et al. 2007). Four days after the first chick of a brood hatched, we conducted a brood size manipulation, in which brood size was standardized to either 2 or 6 young (both within the natural range). Our aim was to cross-foster all individuals in this procedure, but due to logistic constraints, we cross-fostered 75 % of the chicks ($N = 94$).

We measured telomere lengths in DNA from red blood cells. Blood was collected from the brachial vein into heparinised capillaries. Samples were suspended in 2 % EDTA buffer, and within 2 days the red blood cells were spun down, and the pellet was stored in glycerol buffer at $-80\text{ }^{\circ}\text{C}$ after snap freezing. We used blood samples collected in 2006–2010. Storage time prior to analysis (0–6 years) did not affect telomere length ($F_{6,152} = 0.66$, $p = 0.68$). The samples were analysed, divided over seven gels, and timing (batch) of analysis did not affect telomere measurements ($F_{6,152} = 1.48$, $p = 0.19$). Hence, storage time and timing of analysis were not included in the analyses.

On average, individuals were 132 days old ($SE = 11.6$, range = 9–636) when a blood sample was collected. For 18 individuals, we analysed two to three samples, as these individuals were sampled multiple times in life, resulting in a total of 158 telomere length estimates. On average, the samples after the first sample were taken at an age of 940 days ($SE = 52.5$, range = 609–1572). To simplify the models for estimating heritability, and because we had repeated measurements of only a subset of all individuals (18 out of 125 individuals), we used the average telomere length as a Terminal Restriction Fragment (TRF) estimate per individual to calculate heritability. In order to estimate variance components that might influence phenotypic similarity, we used the complete data set, including repeated measurements. Because telomere length generally declines with age, we controlled for age in all analyses.

TRF assay

The TRF assay was conducted following Salomons et al. (2009). In summary, 5 μl of red blood cells were suspended in an agarose solution to form an agarose plug (0.8 %; following the manufacturer’s protocol, CHEF Mammalian Genomic DNA Plug kit, Bio-Rad Laboratories, Inc., USA). The cells in half a plug were digested overnight at $50\text{ }^{\circ}\text{C}$ with Proteinase K. DNA was then digested overnight at

37 °C using a mixture of three restriction enzymes, *Hind*III (60 U), *Hinf*I (30 U) and *Msp*I (60 U), in NEB2 buffer (New England Biolabs, Inc., Beverly MS, USA).

The restricted DNA and the size standards (Molecular Weight Marker XV, Roche and 1 kb DNA ladder, New England Biolabs) were electrophoresed through a 0.8 % agarose gel by pulsed field gel electrophoresis at 14 °C for 24 h (3.5 V/cm, initial switch time 0.5 s, final switch time 7.0 s). Gels were dried with a gel dryer (Bio-Rad, model 538) and hybridized overnight with ³²P-labelled oligo (5-CCCTAA-3)₄, which labelled the single-stranded overhang of the telomeres. Since the DNA was not denatured as in Southern blot techniques, no ³²P-labelled oligo marked interstitial repeats. The radioactive signal of the marker was detected by a phosphor screen (PerkinElmer Inc., USA), and analysed using a phosphor imager (Cyclone TM Storage Phosphor System, PerkinElmer).

Telomere length varies among cells and chromosomes (Lansdorp et al. 1996); and hence, the TRF assay results in a smear, instead of a clear band. The distribution of telomere lengths was calculated based on densitometry (Hausmann and Mauck 2008) in the open-source software IMAGEJ v. 1.38x (Salomons et al. 2009). The average labelled telomere length per lane was calculated as: $\Sigma (OD_i \times L_i) / \Sigma (OD_i)$, where OD_i is the optical density output at position i , and L_i is the length of the DNA (bp) at position i . OD is corrected for the background by subtracting the average grey value of non-DNA containing gel in IMAGEJ. Our lower limit was 2.3 kb, which falls within the smallest band of the 1 kb DNA ladder, which is 1 kb, and our upper limit was an extrapolated value of 80 kb based on the Molecular Weight Marker XV, which has a range of 2.4–48.5 kb, because telomere lengths of the Zebra Finches exceeded the Molecular Weight Marker XV. Note, however, that the extrapolation comprised < 1.5 cm on the gel (± 7 % of the total length used), and that there was a strong correlation between calculations of telomere length based on the Molecular Weight Marker XV (up to 48.5 kb) and the same samples quantified with the extrapolated marker (up to 80 kb) ($r = 0.82$). Based on the repeated measures of 18 individuals, individual variation in TRF assays was 78 % of the total variance in telomere lengths. Since repeat abilities of our TRF assays are high (Jeanclos et al. 2000; Hausmann and Mauck 2008; Salomons et al. 2009) and the analysis is time consuming, all samples were run once.

Statistical analyses

We compiled a pedigree using the Groningen Zebra Finch database. Data on ancestry were available for four generations of birds, with the earliest records dating back to 2004. We used pedigree data pruned back to the 125

phenotyped individuals, plus 143 unphenotyped individuals linking the phenotyped birds. The pedigree contained 112 individuals in a full sibling relationship, 45 maternal half-siblings and 62 paternal half-siblings. Half-sibling comparison facilitated attempts to separate genetic and environmental components. We over-represented paternal half-siblings in our data collection, because females lay the eggs and may thereby potentially exert a greater environmental influence on offspring telomere length, and we were primarily interested in the genetic component of the variance. For further details of the pedigree, see Table S1 (Supplementary material).

We calculated the heritability of telomere length with an ‘animal’ model (Kruuk 2004), using a Bayesian approach (Hadfield 2010), estimating the posterior mode and 95 % credible intervals (95 % Cred. Int.) for fixed effects, variance components and heritability. In short, an ‘animal’ model uses a pedigree to calculate the proportion of the phenotypic variance that is due to additive genetic effects, by comparing the covariance due to additive genetic effects in a phenotype between relatives. We calculated heritability using the package MCMCglmm (2.15) in R 2.14.1 (Hadfield 2010; R Development Core Team 2011) with 10,000,000 iterations, a burn-in of 2500,000 and a thinning interval of 5000. Autocorrelation between sampled iterations was < 0.08. We used default priors for fixed effects, parameter expanded priors for the random variance structure (variance = 1, degree of belief = 1, prior mean = 0, prior covariance matrix = 500), and non-informative inverse-Wishart priors for the residual variance structure (variance = 1, degree of belief = 0.002). We applied several different prior distributions to confirm that our estimate of additive genetic variance was robust to prior specification.

Telomere length was normally distributed (Shapiro–Wilk $W = 0.989$, $p = 0.25$, $N = 158$). Exploratory analyses indicated that including sex and the logarithm of age (we log-transformed age because telomeres shorten faster early in life) as fixed effects improved our model fit. For individuals with an average telomere length of multiple TRF estimates, we used the logarithm of the average age at sampling. Because the data set did not have sufficient power to discriminate between the random variance components explaining environmental (birth nest and permanent environment, meaning an individual’s own common environment) and parental effects, we used a naïve model including only the pedigree component as a random effect. We therefore estimated the variance components as:

$$V_P = V_A + V_R \quad (1)$$

where V_P is the phenotypic variance, accounting for the fixed effects of sex and the logarithm of the age at which an individual was sampled; V_A is the additive genetic variance

and V_R the residual variance. We then calculated heritability as:

$$h^2 = \frac{V_A}{V_P} \quad (2)$$

We compared h^2 from the ‘animal’ model with the heritability estimate based on the intraclass correlations for sibships (Falconer and Mackay 1996), to evaluate robustness of our findings. The individual least square mean estimate of telomere length was used from a model including the logarithm of age. For the full sibling comparison, we used a general linear mixed model (GLMM) with family identity as a random effect, where a family is defined as a set of full siblings. Second, we made a half-sibling comparison in a subset of the data containing half-siblings, by building a GLMM with either mother identity or father identity as a random effect. Here we used a mean value of telomere length for each birth nest, to prevent our estimate from being biased by pseudo-replication by full siblings. This approach allowed us to compare similarity between maternal- and paternal half-siblings, testing for a trans-generational effect from mother or father. Sample sizes for all sibling relationships can be found in Table 4. Some of the individuals in the full sibling comparison were also included in the maternal ($N = 14$) and paternal ($N = 31$) half-sibling comparison. Following Falconer and Mackay (1996) equation 9.8, heritability in the full sibling data set was calculated by multiplying the correlation between full siblings by two:

$$t_{FS} = \frac{\frac{1}{2}V_A + \frac{1}{4}V_D}{V_P} \quad (3)$$

where t_{FS} is the correlation between full siblings and V_D the dominance variance. In general V_D is relatively small compared to V_A , and hence could be ignored (Falconer and Mackay 1996). This was confirmed by the similarity of our heritability estimates of full siblings and half-siblings (see “Results”). Heritability using the half-sibling data set was calculated by multiplying the correlation between half-siblings by 4, according Falconer and Mackay (1996) (equation 9.6)

$$t_{HS} = \frac{1}{4} \frac{V_A}{V_P} \quad (4)$$

where t_{HS} is the correlation between half-siblings.

As we could not include random effects in the ‘animal’ model, nor separate variance components in the intraclass correlations, we estimated variance components that may influence phenotypic similarity in telomere length with a GLMM. We ran a GLMM with sex and the logarithm of age as fixed effects, where our response was telomere length at a given age, rather than one averaged value, and included individual as random effect in the model. In

Table 2, we describe which environmental and genetic effects are embedded in each variance component. We defined phenotypic variance components as significantly different from zero if their confidence intervals (CI) did not overlap with zero.

Ethics statement

The brood size manipulations and long term foraging experiment with Zebra Finches, including blood sampling, have been approved by the animal welfare ethics committee of the University of Groningen (according to Dutch law), under license number 5150.

Results

With our ‘animal’ model, we estimated a high additive genetic variance (V_A) in telomere length, with a heritability of 0.999 (95 % Cred. Int. = 0.87–1). Few repeated measures meant that we could not partition permanent environment effects, but as the chicks were cross-fostered, these are predicted to be small. Despite our study design containing various degrees of relatedness (siblings, and maternal and paternal half-siblings, Table S1), there was not enough power to partition parental (genetic and environmental) effects. Extended models including additional variance components did not converge, due to low sample sizes and a shallow pedigree. Therefore we used a naïve ‘animal’ model, and hence our estimate of heritability may be confounded by environmental or parental effects, which may have inflated our estimate. We found no sex effect and a trend of a negative logarithm of age effect on telomere length (Table 3). Telomere length generally declines with age, but our finding that age yielded only a trend in this analysis was not unexpected, since we used a cross-sectional design, and between-individual variation is large while variation in age within the data set was intentionally small.

To test for robustness of our estimate, we calculated heritability with GLMMs based on cross-fostered full sibling and half-sibling intraclass correlations. Also with this approach we found high heritabilities, confirming the results of the ‘animal’ model (Table 4; Fig. 1). The three estimates were very similar, but CI were large, and only the estimate based on full siblings (with the largest sample size) yielded a 95 % CI that did not include zero. Based on this 95 % confidence interval of the full sibling estimate, the minimum estimate of the heritability is 0.46, assuming no V_D , while the maximum estimate exceeds 1.

Applying GLMMs including repeated telomere measurements enabled us to separate different variance components. In all models (Fig. 2; Table 5), we found an effect of the individual. In model two, we compared effects of the

Table 2 Separation of phenotypic variance components into genetic, environmental and parental effects

	Additive genetic	Permanent environment	Common environment	Maternal effects ^a	Paternal effects ^a
Individual	×	×		×	×
Birth nest	×		×	×	×
Rear nest			×		
Mother	×			×	
Father	×				×

^a Parental effects include parental environmental and non-additive genetic components

Table 3 Mode of the posterior distribution and 95 % credible intervals for the different parameters in the ‘animal’ model

	Posterior mode	95 % Credible interval
Fixed effects		
Sex	0.364	−0.164–0.705
Log age	−0.447	−0.926–0.044
Random effects		
Animal	1.827	1.222–2.378
Residual	0.003	0.000–0.382

pre-cross foster natal environment (which may include parental effects) with post-cross foster rear nest environment (94 out of 125 individuals were cross-fostered). We found no effect of rear nest identity in this or any subsequent model, indicating that pre-manipulation effects (both genetic and


environmental) were more important in determining telomere length than post-manipulation effects. Also, no effect of manipulated brood size was found in any of our models (e.g., in model 4, Fig. 2: $F_{1,55.75} = 0.18$; $p = 0.67$). Therefore, brood size was not included in our final analyses. Birth nest includes multiple potential variance components, e.g., shared environment, maternal and paternal effects. When adding parental identity (model three), the variance explained by birth nest went down to zero, with mother and father identity taking over this effect, i.e., trans-generational parental effects were important, while characteristics of that specific breeding event explained no additional variation. Mother and father identity contributed substantially and about equally to telomere length variation, as also shown by the correlational approach in the (half) siblings (Fig. 1). Together, mother and father identity explained over two times more variation than

Table 4 Heritability estimates based on the intraclass correlations of telomere length corrected for sex and age

	<i>N</i> (families)	<i>r</i> (95 % CI)	<i>h</i> ² (95 % CI)
Full siblings	94 (42)	0.59 (0.23–0.95)	1.18 (0.46–1.90)
Maternal half-siblings	27 (8)	0.34 (−0.26–0.93)	1.35 (−1.04–3.74)
Paternal half-siblings	60 (18)	0.23 (−0.07–0.53)	0.93 (−0.27–2.13)

N is the number of individuals in the analyses, with the number of natal families in brackets. The correlation (*r*), heritability (*h*²) and their 95 % confidence intervals (95 % CI) are given

Fig. 1 Correlations between full siblings (*N* = 42) (a) and half-siblings (b); closed dots maternal half-siblings (*N* = 8); and open dots paternal half-siblings (*N* = 18). Lines are linear regressions. To avoid pseudo replication due to shared maternal or paternal descent, both graphs depict a random selection of sibling pairs in cases where there were more than two phenotyped siblings, whereas in the analyses, all individuals from a family are incorporated with family as a random effect






Fig. 2 Variance components of telomere length, from four GLMMs. *Plus* depicts significant variance components for which the confidence interval (CI) does not overlap with zero and (*plus*) variance components for which the CI just overlaps zero (<1 %). *The numbers above the bars* are the log likelihoods of each model

individual identity, indicating that additive genetic and common natal environment effects were substantially more important in determining telomere length than other permanent events over the whole lifetime. Based on the log likelihood (Fig. 2) and significance of the variance components, we considered model 4 biologically most parsimonious. This implied that it is likely that the individual (which contains the additive genetic variance and a permanent environment effect), and mother and father (which contain the additive genetic variance and a parental environment effect) all contributed about equally to variation in telomere length.

Discussion

We estimated the heritability of terminal telomere length in Zebra Finches, and find it to approach 1, independent of the type of numerical approach [‘animal’ model or (half-)sibling comparison]. We found no effects of rear nest identity,

suggesting that possible environmental agents exerted their effects very early in life. Most phenotyped offspring were cross-fostered at a young age, but cross-fostering of chicks does not control for parental effects arising during laying, incubation and in the few days after hatching that they spent in their natal nest. Pre-natal environmental effects on telomere length can, for example, include endocrinology aspects of egg composition (Haussmann et al. 2012), and we cannot exclude that such effects increased the resemblance between (half-)siblings in our study. On the other hand, the mother primarily determines egg characteristics, and effects of father and mother identity on offspring telomere length were indistinguishable in our study (Table 5, model 4). This argues against a large effect of egg characteristics causing telomere length resemblance between offspring, making a quantitative genetic basis of the observed resemblance between relatives more likely. Furthermore, heritability estimates based on full siblings (1.18) and paternal half-siblings (0.93) were of similar magnitude (albeit with large CI, Table 4), lending further support to the tentative conclusion that telomere length resemblance between relatives was primarily due to genetic effects. We acknowledge, however, that larger sample sizes and/or a deeper pedigree are required to more definitely draw this conclusion.

Our study distinguishes itself from earlier reports in that we specifically measured terminal telomeres, while in previous studies (Table 1), interstitial and terminal telomeres were pooled due to the technique used to measure telomere length. This is of importance, because terminal telomeres are susceptible to ageing and predictors of survival, while interstitial telomeric repeats are, as far as we know, inert within the lifetime of an individual. It is not obvious what the effect of measurement technique will be on heritability estimates, because this depends on (1) the reliability of the measurement technique, and whether this

Table 5 Estimates for percentage of phenotypic variance explained, with 95 % CI per variance component and estimates for the fixed effects (sex coded as 1 = female, 2 = male, reference is female) with standard error and *p* value in four different GLMMs of telomere length

Model	Random effects					Fixed effects			
	Individual	Birth nest	Rear nest	Mother	Father	Sex		Log age (days)	
						Estimate (SE)	<i>p</i> value	Estimate (SE)	<i>p</i> value
1	80.1 (53.7–106.4)					–130.0 (120.5)	0.283	–448.9 (138.0)	0.002
2	36.6 (6.4–66.9)	56.4 (27.0–85.8)	–12.8 (– 31.6–6.1)			–101.9 (109.8)	0.356	–529.6 (149.5)	<0.001
3	24.7 (6.3–43.1)	–9.0 (– 23.1–5.2)		30.2 (4.4–56.0)	35.8 (5.2–66.3)	–160.2 (113.0)	0.160	–523.6 (159.3)	0.002
4	23.4 (6.1–40.8)			26.9 (–0.5–54.4)	30.7 (0.3–61.1)	–143.3 (105.8)	0.179	–553.6 (143.3)	<0.001

Significant effects are in bold, marginally significant effects are in italics

is accounted for in the analysis; and (2) contribution of the interstitial repeats versus terminal telomeres to the total variance in telomeric repeats between individuals. For example, interstitial repeats are infrequent in humans, and in this respect, the measurement technique should have little effect on telomere length heritability estimates in humans. In avian species, however, the frequency of interstitial repeats is frequently high (Foote et al. 2013), and it remains to be investigated to what extent published heritability estimates (Table 1) can be attributed to variance in interstitial repeats versus terminal telomeres. To resolve this issue, additional studies in which the heritability can specifically be assigned to either interstitial telomeric repeats and/or terminal telomeres are required.

Terminal telomeres shorten with age, and these dynamics are at least to some extent under the influence of environmental effects (e.g., Boonekamp et al. 2014). This could have increased family resemblance in our study (Rossiter 1996), because all birds were housed in similar aviaries, reducing environmental variation later in life, in particular in comparison with free-living animals. However, such homogenizing environmental effects in all likelihood did not have a large effect on our estimates, since we found no effect of rear nest, and permanent environment explained approximately one-quarter of the variance in telomere length (Table 5). Therefore, sharing the same genes and/or the pre-cross foster environment (including parental effects) were the strongest determinants of our heritability estimates. Individual differences in human telomere length persist over life (Benetos et al. 2013), and initial telomere lengths are determined in the zygote, with only a minor effect of epigenetic and/or environmental effects during life on resemblance of telomere length between relatives (Graakjaer et al. 2004). One pathway explaining the minor effect of late environment is via the telomere-elongating enzyme telomerase, which is involved in the process of maintaining telomere length in particular tissues during development. Serakinci et al. (2008) suggest that telomere dynamics in lymphocytes and mesenchymal stem cells show little random fluctuation and that telomerase possibly even further conserves the relative telomere lengths or profile between chromosome arms (Serakinci et al. 2008). It seems that specific patterns of telomere lengths are already determined in the embryo, and telomerase is an important determinant during life for resemblance of telomere lengths between relatives.

In humans (e.g. Broer et al. 2013), Kakapos (Horn et al. 2011) and King Penguins (Reichert et al. 2014), telomere length has a stronger maternal than paternal inheritance; however, in Sand Lizards (Olsson et al. 2011), stronger paternal inheritance was found, although CI overlapped (Table 1). In contrast, we found no evidence for a

difference between maternal and paternal effects in our study (Tables 4, 5). The non-human studies have relatively small sample sizes compared to the human studies, and used—except for the Great Reed Warbler study, which used an ‘animal’ model—parent-offspring regression, which is unable to separate environmental variance from genetic components. The human study included both parent-offspring regression and twin studies, which also do not separate environmental factors. Furthermore, in humans, a positive association with paternal age was demonstrated, whereas this was negative in Sand Lizards. Thus, the results on parental effects are mixed, and for non-human species in particular, there is a need for more studies.

Acknowledgments We thank Peter Korsten for feedback that improved our analyses, and an anonymous reviewer for comments that improved the manuscript. The long-term foraging experiment was supported by a Netherlands Organization for Scientific Research (NWO) Vici-grant to S.V. E.A. was funded by a grant from the Netherlands Organization for Scientific Research (ALW-NWO) to A.v.N. and S.V. H.L.D. was funded by a Natural Environment Research Council (NERC) fellowship. The experiment done in this study complies with the current laws of The Netherlands.

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Al-Attas OS, Al-Daghri NM, Alokail MS et al (2012) Circulating leukocyte telomere length is highly heritable among families of Arab descent. *BMC Med Genet* 13:38. doi:[10.1186/1471-2350-13-38](https://doi.org/10.1186/1471-2350-13-38)
- Andrew T, Aviv A, Falchi M et al (2006) Mapping genetic loci that determine leukocyte telomere length in a large sample of unselected female sibling pairs. *Am J Hum Genet* 78:480–486. doi:[10.1086/500052](https://doi.org/10.1086/500052)
- Asghar M, Bensch S, Tarka M et al (2015) Maternal and genetic factors determine early life telomere length. *Proc R Soc B Biol Sci* 282(1799):20142263
- Bauch C, Becker PH, Verhulst S, PRS B (2013) Telomere length reflects phenotypic quality and costs of reproduction in a long-lived seabird. *Proc R Soc B Biol Sci*. doi:[10.1098/rspb.2012.2540](https://doi.org/10.1098/rspb.2012.2540)
- Benetos A, Kark JD, Susser E et al (2013) Tracking and fixed ranking of leukocyte telomere length across the adult life course. *Aging Cell* 12:615–621. doi:[10.1111/ace1.12086](https://doi.org/10.1111/ace1.12086)
- Bischoff C, Graakjaer J, Petersen HC et al (2005) The heritability of telomere length among the elderly and oldest-old. *Twin Res Hum Genet* 8:433–439. doi:[10.1375/183242705774310141](https://doi.org/10.1375/183242705774310141)
- Bize P, Criscuolo F, Metcalfe NB et al (2009) Telomere dynamics rather than age predict life expectancy in the wild. *Proc R Soc B Biol Sci* 276:1679–1683. doi:[10.1098/rspb.2008.1817](https://doi.org/10.1098/rspb.2008.1817)
- Blackburn EH (1991) Structure and function of telomeres. *Nature* 350:569–573. doi:[10.1146/annurev.genet.23.1.579](https://doi.org/10.1146/annurev.genet.23.1.579)

- Blackburn EH (2005) Telomeres and telomerase: Their mechanisms of action and the effects of altering their functions. *FEBS Lett* 579:859–862
- Boonekamp JJ, Simons MJP, Hemerik L, Verhulst S (2013) Telomere length behaves as biomarker of somatic redundancy rather than biological age. *Aging Cell* 12:330–332. doi:10.1111/ace.12050
- Boonekamp JJ, Mulder GA, Salomons HM et al (2014) Nestling telomere shortening, but not telomere length, reflects developmental stress and predicts survival in wild birds. *Proc R Soc B Biol Sci*. doi:10.1098/rspb.2013.3287
- Broer L, Codd V, Nyholt DR et al (2013) Meta-analysis of telomere length in 19,713 subjects reveals high heritability, stronger maternal inheritance and a paternal age effect. *Eur J Hum Genet* 21:1163–1168. doi:10.1038/ejhg.2012.303
- De Coster G, Verhulst S, Koetsier E et al (2011) Effects of early developmental conditions on innate immunity are only evident under favourable adult conditions in Zebra Finches. *Naturwissenschaften* 98:1049–1056. doi:10.1007/s00114-011-0863-3
- Delany ME, Krupkin AB, Miller MM (2000) Organization of telomere sequences in birds: evidence for arrays of extreme length and for in vivo shortening. *Cytogenet Cell Genet* 90:139–145 (15649)
- Epel ES, Blackburn EH, Lin J et al (2004) Accelerated telomere shortening in response to life stress. *Proc Natl Acad Sci USA* 101:17312–17315. doi:10.1073/pnas.0407162101
- Falconer DS, Mackay TFC (1996) Introduction to quantitative genetics, 4th edn. Harlow, Essex
- Foote CG, Vleck D, Vleck CM (2013) Extent and variability of interstitial telomeric sequences and their effects on estimates of telomere length. *Mol Ecol Resour* 13:417–428. doi:10.1111/1755-0998.12079
- Forstmeier W, Segelbacher G, Mueller JC, Kempenaers B (2007) Genetic variation and differentiation in captive and wild Zebra Finches (*Taeniopygia guttata*). *Mol Ecol* 16:4039–4050. doi:10.1111/j.1365-294X.2007.03444.x
- Gilley D, Herbert BS, Huda N et al (2008) Factors impacting human telomere homeostasis and age-related disease. *Mech Ageing Dev* 129:27–34. doi:10.1016/j.mad.2007.10.010
- Graakjaer J, Pascoe L, Der-Sarkissian H et al (2004) The relative lengths of individual telomeres are defined in the zygote and strictly maintained during life. *Aging Cell* 3:97–102. doi:10.1111/j.1474-9728.2004.00093.x
- Hadfield J (2010) MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *J Stat Softw* 33:1–22
- Hausmann MF, Mauck RA (2008) Technical advances: new strategies for telomere-based age estimation. *Mol Ecol Resour* 8:264–274. doi:10.1111/j.1471-8286.2007.01973.x
- Hausmann MF, Vleck CM (2002) Telomere length provides a new technique for aging animals. *Oecologia* 130:325–328. doi:10.1007/s00442-001-0827-y
- Hausmann MF, Winkler DW, Vleck CM (2005) Longer telomeres associated with higher survival in birds. *Biol Lett* 1:212–214. doi:10.1098/rsbl.2005.0301
- Hausmann MF, Longenecker AS, Marchetto NM et al (2012) Embryonic exposure to corticosterone modifies the juvenile stress response, oxidative stress and telomere length. *Proc R Soc B Biol Sci* 279:1447–1456. doi:10.1098/rspb.2011.1913
- Heidinger BJ, Blount JD, Boner W et al (2012) Telomere length in early life predicts lifespan. *Proc Natl Acad Sci USA* 109:1743–1748. doi:10.1073/pnas.1113306109
- Horn T, Robertson BC, Will M et al (2011) Inheritance of telomere length in a bird. *PLoS One* 6:e17199. doi:10.1371/journal.pone.0017199
- Jeanclous E, Schork NJ, Kyvik KO et al (2000) Telomere length inversely correlates with pulse pressure and is highly familial. *Hypertension* 36:195–200
- Joeng KS, Song EJ, Lee K-J, Lee J (2004) Long lifespan in worms with long telomeric DNA. *Nat Genet* 36:607–611. doi:10.1038/ng1356
- Kilburn AE, Shea MJ, Sargent RG, Wilson JH (2001) Insertion of a telomere repeat sequence into a mammalian gene causes chromosome instability. *Mol Cell Biol* 21:126–135. doi:10.1128/MCB.21.1.126-135.2001
- Koetsier E, Verhulst S (2011) A simple technique to manipulate foraging costs in seed-eating birds. *J Exp Biol* 214:1225–1229. doi:10.1242/jeb.050336
- Kotrschal A, Ilmonen P, Penn DJ (2007) Stress impacts telomere dynamics. *Biol Lett* 3:128–130. doi:10.1098/rsbl.2006.0594
- Kruuk LEB (2004) Estimating genetic parameters in natural populations using the “animal model”. *Philos Trans R Soc Lond B Biol Sci* 359:873–890. doi:10.1098/rstb.2003.1437
- Kruuk LEB, Hadfield JD (2007) How to separate genetic and environmental causes of similarity between relatives. *J Evol Biol* 20:1890–1903. doi:10.1111/j.1420-9101.2007.01377.x
- Lansdorp PM, Verwoerd NP, van de Rijke FM et al (1996) Heterogeneity in telomere length of human chromosomes. *Hum Mol Genet* 5:685–691. doi:10.1093/hmg/5.5.685
- Meyne J, Ratliff RL, Moyzis RK (1989) Conservation of the human telomere sequence (TTAGGG)_n among vertebrates. *Proc Natl Acad Sci USA* 86:7049–7053
- Monaghan P (2014) Organismal stress, telomeres and life histories. *J Exp Biol* 217:57–66. doi:10.1242/jeb.090043
- Nawrot TS, Staessen JA, Gardner JP, Aviv A (2004) Telomere length and possible link to X chromosome. *Lancet* 363:507–510. doi:10.1016/S0140-6736(04)15535-9
- Njajou OT, Cawthon RM, Damcott CM et al (2007) Telomere length is paternally inherited and is associated with parental lifespan. *Proc Natl Acad Sci USA* 104:12135–12139. doi:10.1073/pnas.0702703104
- Nordfjäll K, Svenson U, Norrback K-F et al (2010) Large-scale parent-child comparison confirms a strong paternal influence on telomere length. *Eur J Hum Genet* 18:385–389. doi:10.1038/ejhg.2009.178
- Olovnikov AM (1973) A theory of marginotomy. The incomplete copying of template margin in enzymic synthesis of polynucleotides and biological significance of the phenomenon. *J Theor Biol* 41:181–190. doi:10.1016/0022-5193(73)90198-7
- Olsson M, Pauliny A, Wapstra E et al (2011) Sex differences in sand lizard telomere inheritance: paternal epigenetic effects increases telomere heritability and offspring survival. *PLoS One* 6:e17473. doi:10.1371/journal.pone.0017473
- R Development Core Team (2011) R: a language and environment for statistical computing. *R Found Stat Comput* 1:409. doi:10.1007/978-3-540-74686-7
- Reichert S, Rojas ER, Zahn S et al (2014) Maternal telomere length inheritance in the king penguin. *Heredity (Edinb)* 114(1):10–16. doi:10.1038/hdy.2014.60
- Rivero MT, Mosquera A, Goyanes V et al (2004) Differences in repair profiles of interstitial telomeric sites between normal and DNA double-strand break repair deficient Chinese hamster cells. *Exp Cell Res* 295:161–172. doi:10.1016/j.yexcr.2003.12.031
- Rossiter M (1996) Incidence and consequences of inherited environmental effects. *Annu Rev Ecol Syst* 27:451–476. doi:10.1146/annurev.ecolsys.27.1.451
- Salomons HM, Mulder GA, van de Zande L et al (2009) Telomere shortening and survival in free-living corvids. *Proc Biol Sci* 276:3157–3165. doi:10.1098/rspb.2009.0517

- Serakinci N, Graakjaer J, Kolvraa S (2008) Telomere stability and telomerase in mesenchymal stem cells. *Biochimie* 90:33–40. doi:[10.1016/j.biochi.2007.09.005](https://doi.org/10.1016/j.biochi.2007.09.005)
- Slagboom PE, Droog S, Boomsma DI (1994) Genetic determination of telomere size in humans: a twin study of three age groups. *Am J Hum Genet* 55:876–882
- Stansel RM, De Lange T, Griffith JD (2001) T-loop assembly in vitro involves binding of TRF2 near the 3' telomeric overhang. *EMBO J* 20:5532–5540. doi:[10.1093/emboj/20.19.5532](https://doi.org/10.1093/emboj/20.19.5532)
- Vasa-Nicotera M, Brouillette S, Mangino M et al (2005) Mapping of a major locus that determines telomere length in humans. *Am J Hum Genet* 76:147–151. doi:[10.1086/426734](https://doi.org/10.1086/426734)
- Visscher PM, Hill WG, Wray NR (2008) Heritability in the genomics era—concepts and misconceptions. *Nat Rev Genet* 9:255–266. doi:[10.1038/nrg2322](https://doi.org/10.1038/nrg2322)
- Voillemot M, Hine K, Zahn S et al (2012) Effects of brood size manipulation and common origin on phenotype and telomere length in nestling collared flycatchers. *BMC Ecol* 12:17. doi:[10.1186/1472-6785-12-17](https://doi.org/10.1186/1472-6785-12-17)
- Von Zglinicki T (2002) Oxidative stress shortens telomeres. *Trends Biochem Sci* 27:339–344
- Yang D, Xiong Y, Kim H et al (2011) Human telomeric proteins occupy selective interstitial sites. *Cell Res* 21:1013–1027. doi:[10.1038/cr.2011.39](https://doi.org/10.1038/cr.2011.39)