

University of Groningen

Self-rated health and mortality after kidney transplantation

Majerníková, Mária

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Majerníková, M. (2013). Self-rated health and mortality after kidney transplantation. Košice, Slovakia: University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Patient and graft mortality as a health outcome

Chapter 7

Self-rated health predicts mortality and graft loss after kidney transplantation: a 10-year follow-up study

Maria Majernikova, Jaroslav Rosenberger, Lucia Prihodova, Iveta Nagyova, Robert Roland,
Johan W Groothoff, Jitse P van Dijk

American Journal of Nephrology 2012; 36 (5): 459-465. DOI: 10.1159/000343884.

Abstract

Background

This study explored whether self-rated health (SRH) shortly after kidney transplantation (KT) predicts mortality and graft loss at up to 10 years follow-up.

Methods

A total of 276 patients shortly after successful KT were interviewed. SRH was measured using the first item of the SF-36 questionnaire and divided into 3 tertiles: poor, average and excellent health. Clinical data were retrieved from medical records. Cox regression was used to identify whether different levels of SRH predicted mortality and graft loss in transplant recipients. The observation period was up to 10 years.

Results

Poor SRH (HR 11.1, $p<0.001$), average SRH (HR 4.21, $p<0.05$), estimated glomerular filtration rate (HR 0.26, $p<0.05$) and age (HR 1.04, $p<0.05$) were significantly associated with mortality. Similarly, poor SRH (HR 6.4, $p<0.001$), average SRH (HR 3.6, $p<0.05$), new-onset diabetes mellitus after KT (HR 3.3, $p<0.05$) and chronic renal allograft dysfunction (HR 3.7, $p<0.00$) were significantly associated with graft loss.

Conclusion

Poor SRH shortly after transplantation indicates an increased risk of mortality and graft loss at up to 10 years follow-up. SRH could be an inexpensive and reliable indicator for starting diagnostic and/or treatment strategies. The usefulness of SRH compared to other global clinical measures predicting mortality and graft loss should also be studied.

Key words

graft loss, kidney transplantation, mortality, self-rated health

Introduction

Self-rated health (SRH) is an independent predictor of mortality in patients with end-stage renal disease (ESRD) on chronic dialysis.¹ Persons with poor SRH have a higher mortality risk compared with those with excellent SRH even after controlling for a range of demographic and clinical variables.^{1,2} Previous studies have also shown that physical, psychological and total scores of health-related quality of life were significantly correlated with increased risks of ESRD and death in patients with chronic kidney disease, independent of comorbidity factors.^{3,4}

The medical predictors of mortality are also known. A study by Desai et al. has shown that cardiovascular and inflammatory markers were predictors of mortality and might have important implications for risk stratification in the ESRD population.⁵ Other studies have demonstrated the associations between infections, graft rejections, donor and recipient factors with increased mortality and graft loss in kidney transplant recipients.⁶⁻⁹ Ravindran et al. explored the idea that new-onset diabetes mellitus after transplantation was a well-recognized complication of solid organ transplantation associated with higher patient mortality and graft loss.¹⁰

Many studies in ESRD, including those after transplantation, have shown medical factors to be associated with mortality and graft loss.⁶⁻¹⁰ Moreover, Novak et al. showed that depressive symptoms were an independent predictor of mortality censored graft loss in patients after KT.¹¹ Surprisingly, we found no studies on SRH as a predictor of mortality or graft loss in kidney transplant recipients.

The aim of this prospective observation study is to explore whether self-rated health in an early period after kidney transplantation predicts mortality and graft loss at up to 10 years follow-up.

Methods

Sample and procedure


A total of 362 consecutive patients who underwent KT between January 2001 and January 2011 at the Transplant Centre of Kosice in the eastern region of Slovakia were enrolled in the study. The baseline examination of the participants occurred between the 3rd and 12th month after successful KT. The first 3 months after KT are usually considered to be the most problematic period connected to dramatic changes, increased morbidity and even mortality.¹² Furthermore, improvement in self-perceived health most often occurs during the first 2 years after KT.¹³ Before the first 3 months after KT, 9 (2.5%) patients were not included because 3 (0.8%) of them had died and 6 (1.7%) had lost their transplanted kidney. The exclusion criteria were the presence of mental retardation, acquired cognitive impairments, severe dementia or other psychiatric diseases mentioned in the medical record. After the exclusion of 1 (0.3%) patient because of the exclusion criteria a total of 352 kidney transplant recipients after successful transplant surgery were invited to participate. Of those 34 (9.7%) refused to participate, 31 (8.8%) did not return the questionnaire and 11 (3.1%) provided questionnaire with missing data, resulting in a total of 276 patients (an effective response rate 78.4%) at the start of the study. Figure 7.1 presents more detailed information on participants.

Patients provided information about sociodemographic variables and filled in the

questionnaire. All participants were interviewed during regular outpatient clinical visits by trained personnel independent of the transplant team. Medical data were retrieved from medical records at the same time as socio-demographic data and data of SRH.

Only patients who signed informed consent prior to the study were included. The local Ethics Committee in Kosice approved the study.

Figure 7.1 Flow-chart diagram of the participants



N/n—Number, RR—response rate; KT—kidney transplantation

Measures

Sociodemographic data included age, gender and education. Age was treated as a continual variable. We categorized education into elementary (primary school, completed or not), apprenticeship (completed, or not-completed secondary school), secondary (completed, or not-completed university) and university (completed).

Self-rated health (SRH) was measured using the first question of the Short Form Health Survey (SF-36).¹⁴ The answer options 1 (poor), 2 (fair), 3 (good), 4 (very good) and 5 (excellent) were transformed into a standard scale from 0 (poor health) to 100 (excellent health) in which the higher scores indicate better health status.¹⁴ This scale was then categorized into tertiles: scores 0 to 30 (fair and poor) were categorized as 'poor health'; 31 to 60 (good) as 'average health'; and 61 to 100 (excellent and very good) as 'excellent health'.¹⁵ The validity and reliability of the first item of the SF-36 has been confirmed in patients with renal disease, including those after KT.¹⁶⁻¹⁸

Clinical data were retrieved from medical files. These included serum creatinine, previous duration of dialysis (in years), primary kidney diagnosis, source of transplanted kidney, function immediately after KT, comorbidity, the current and rejection immunosuppressive treatment, acute rejection episodes, chronic renal allograft dysfunction, uroinfection, which

included pyelonephritis, comorbidities and diagnosis of graft loss, and mortality. The estimated glomerular filtration rate (eGFR) was calculated using the CKD-EPI formula.¹⁹ Acute rejection episodes and chronic renal allograft dysfunction were diagnosed from biopsy according to the Banff 2009 update of diagnostic categories for renal allograft biopsies.²⁰

Statistical analysis

The Mann-Whitney U-test and χ^2 test were used to test the differences between respondents and non-respondents. Frequencies, means and standard deviations were calculated for the sample description. Both tests were used to identify the association between the dependent variables (mortality and graft loss), and the other variables: age, gender, education, SRH categorized into tertiles (excellent SRH - reference category), eGFR at baseline, uroinfection (pyelonephritis included), number of acute rejection episodes, chronic renal allograft dysfunction, source of transplanted kidney, cardiovascular disease – coronary artery disease, cardiac failure, myocardial infarction and categories of diabetes mellitus (no diabetes mellitus – reference category, already existing diabetes mellitus and new-onset diabetes mellitus after transplantation). Cox regression (conditional LR method) was performed in order to identify the predictors of mortality (censored for graft loss) and predictors of graft loss. The independent variables in both Cox regression models were all variables with $p < 0.1$ in Mann-Whitney U-test and χ^2 test, as appropriate. Harrell's C-statistic and Somers' D were calculated for both models (model for mortality and model for graft loss). The Statistical Package for the Social Science (SPSS Inc. Chicago, IL, USA) version 18.0.3 and STATA/SE 11.1 were used for statistical analyses.

Results

No significant differences were found between respondents and non-respondents regarding age and gender. Also, no significant differences were found between those who provided complete and incomplete data in age, gender, mortality and graft loss. The observation period was up to 10 years (mean 5.8 ± 2.7) follow-up; the mean period for poor SRH was 4.5 ± 0.2 years, for average SRH 6.5 ± 0.2 years and for excellent SRH 6.6 ± 0.4 years. The mean period between transplantation and mortality for poor SRH was 3.8 ± 1.5 years, for average SRH 6.2 ± 3.5 years and for excellent SRH 7.2 ± 2.6 years. The mean period between transplantation and graft loss for poor SRH was 4.2 ± 2.3 years, for average SRH 5.6 ± 1.5 years and for excellent SRH 7.3 ± 1.7 . During the observation period 40 patients with poor SRH (39.2%) died and 31 (30.4%) lost their transplanted kidney; 11 with average SRH (9.0%) died and 20 (16.4%) lost their graft and 2 with excellent SRH (3.8%) died and 3 (5.8%) lost their graft. Table 7.1 displays detailed information about the characteristics of the sample (N=257).

The Mann-Whitney U-test showed that age ($p < 0.001$) and eGFR ($p < 0.001$) were associated with mortality. Age ($p < 0.001$), eGFR ($p < 0.001$), and the number of acute rejection episodes ($p < 0.05$) were associated with graft loss. The χ^2 test was performed in order to identify factors associated with mortality and graft loss, and the findings were that SRH ($\chi^2 = 45.2$; $p < 0.001$) and cardiac failure ($\chi^2 = 2.7$; $p = 0.1$) were associated with mortality, while SRH ($\chi^2 = 15.9$; $p < 0.001$), chronic renal allograft dysfunction ($\chi^2 = 42.7$; $p < 0.001$), and diabetes ($\chi^2 = 4.5$; $p < 0.1$) were associated with graft loss. These associations are marked in Table 7.1 and the mentioned variables were used as independent factors in both Cox regression models (Model 1 for mortality and Model 2 for graft loss).

Table 7.1 Characteristics of the sample (N=276)

| | | N(%) or mean±SD | M | GL |
|--|---|-----------------|----|----|
| Age | | 48.27±12.34 | \$ | # |
| Gender | Male | 164(59.4%) | | |
| | Female | 112(40.6%) | | |
| Education | elementary | 48(17.4%) | | |
| | apprenticeship | 125(45.3%) | | |
| | secondary | 81(29.3%) | | |
| | university | 22(8.0%) | | |
| SRH | | 49.4±1.56 | \$ | # |
| Category of SRH | poor | 102(37.0%) | \$ | # |
| | average | 122(44.2%) | \$ | # |
| | excellent | 52(18.8%) | \$ | # |
| Duration on dialysis before KT (in years) | | 3.66±3.04 | | |
| Primary diagnosis of kidney failure | Glomerulonephritis | 98(35.5%) | | |
| | Tubulointerstitial nephritis | 65(23.6%) | | |
| | Vascular disease | 26(9.4%) | | |
| | Polycystic kidneys adult type | 25(9.1%) | | |
| | Diabetic nephropathy | 18(6.5%) | | |
| | Other/unknown | 44(15.9%) | | |
| Source of transplanted kidney | deceased donor | 262(94.9%) | | |
| | living donor | 14(5.1%) | | |
| Function immediately after KT | immediate | 157(56.88%) | | |
| | delayed | 119(43.12%) | | |
| Acute rejection episodes | | 93(33.7%) | | # |
| Type of rejection treatment | steroids | 72(27.9%) | | |
| | antithymocyte globulin | 10(3.6%) | | |
| | plasmapheresis | 6(2.2%) | | |
| | plasmapheresis+i.v.immunoglobuline | 5(1.8%) | | |
| Chronic renal allograft dysfunction | | 42(15.2%) | | # |
| Uroinfection | | 84(30.4%) | | |
| Immunosuppression treatment at baseline | CsA+P | 35(12.7%) | | |
| | CsA+AZA/CsA+AZA+P | 24(8.7%) | | |
| | CsA+MMF/CsA+MMF+P | 139(50.4%) | | |
| | Tac+MMF/Tac+MMF+P | 73(26.4%) | | |
| | SIR+MMF+P/EVER+CsA+MMF | 5(1.8%) | | |
| Estimated glomerular filtration rate (ml/min/1.73m2) at baseline | | 58.8±19.8 | \$ | # |
| Comorbidities | Coronary artery disease | 78(28.3%) | | |
| | Cardiac failure | 72(26.1%) | \$ | |
| | Myocardial infarction | 17(6.2%) | | |
| | Hypertension | 226(81.9%) | | |
| | Diabetes mellitus identified before KT | 18(6.5%) | | |
| | New onset diabetes mellitus after transplantation | 17(6.2%) | | # |
| | CKD-MBD | 160(58.0%) | | |
| | Other comorbidities;≥2 | 9(3.3%) | | |
| Diagnosis of GL | Acute rejection episodes | 5(9.3%) | | |
| | Chronic renal allograft dysfunction | 16(29.6%) | | |
| | Uroinfections | 14(25.9%) | | |
| | Others/unknown | 19(35.2%) | | |
| Diagnosis of M | Acute myocardial infarction | 25(47.2%) | | |
| | Pulmonary disease/pulmonary embolism | 7(13.2%) | | |
| | Stroke | 3(5.7%) | | |
| | Carcinoma/liver disease | 6(11.3%) | | |
| | Others/unknown | 12(22.6%) | | |

N–Number; SD–Standard deviation; KT–kidney transplantation, SRH–self-rated health, CKD-MBD–Chronic kidney disease–Mineral bone disorder. M–mortality; \$–association with mortality (p<0.1); GL–graft loss; #–association with graft loss (p<0.1)

Acute rejection episodes and cardiac failure were not predictors associated with mortality and graft loss in either Cox regression model.

For mortality Harrell's C-statistic remitted 0.88 and Somers' D 0.75. Poor (HR 11.1, 95%CI 4.22;29.04, $p<0.001$) and average (HR 4.21, 95%CI 1.3;13.71, $p<0.038$) SRH contributed significantly to this model, as did eGFR (HR 0.98, 95%CI 0.96;1.0, $p<0.021$) and age (HR 1.04, 95%CI 1.0;1.09, $p<0.03$). Regarding this, the risk of death was increased by 4% for each year of age; on the other hand, the risk of death was decreased by 2% for each 1 ml/min/1.73m² eGFR (Table 7.2). Figure 7.2 displays the differences in mortality between poor, average and excellent SRH.

For graft loss Harrell's C-statistic was 0.84 and Somers' D 0.68. Poor (HR 6.43, 95%CI 2.98;13.85, $p<0.001$) and average (HR 3.58, 95%CI 1.26;10.2, $p<0.017$) SRH contributed significantly to this model, as did chronic renal allograft dysfunction (HR 3.71, 95% CI 1.82;7.6, $p<0.001$) and new-onset diabetes mellitus after transplantation (HR 3.29, 95%CI 1.16;9.34, $p<0.015$).

In line with this, chronic renal allograft dysfunction was associated with a 3.7-fold higher and new-onset diabetes mellitus after transplantation with a 3.3-fold higher risk of graft loss (Table 7.2). Figure 7.3 displays the differences in graft loss between poor, average and excellent SRH.

Table 7.2 Final models of Cox regression containing significant predictors of mortality (Model 1) and graft loss (Model 2)

| Models | | 2Log Likelihood | B(SE) | HR | 95% CI for HR |
|---|-----------|-----------------|---------------|-----------|---------------|
| Model 1 - Mortality (N=276); Harrell's C-statistic 0.88; Somers' D 0.75 | | | | | |
| age | | 365.39 | 0.04(0.02)* | 1.04 | 1.00;1.09 |
| eGFR | | 356.59 | -0.02(0.01)* | 0.98 | 0.96;1.00 |
| SRH | poor | | 2.41(0.49)*** | 11.1 | 4.22;29.04 |
| | average | 323.62 | 1.44(0.60)* | 4.21 | 1.30;13.71 |
| | excellent | | | Reference | |
| Model 2 - Graft loss (N=276); Harrell's C-statistic 0.84; Somers' D 0.68 | | | | | |
| chronic renal allograft dysfunction | | 398.97 | 1.31(0.37)*** | 3.71 | 1.82;7.60 |
| new onset diabetes mellitus after transplantation | | 379.86 | 1.19(0.53)* | 3.29 | 1.16;9.34 |
| SRH | poor | | 1.86(0.39)*** | 6.43 | 2.98;13.85 |
| | average | 363.78 | 1.28(0.53)* | 3.58 | 1.26;10.20 |
| | excellent | | | Reference | |

*- $p<0.05$, **- $p<0.01$, ***- $p<0.001$

SRH–Self-Rated Health, eGFR–estimated Glomerular filtration rate, B(SE) – Unstandardized Coefficient B (Standard Error), HR – hazard ratio, CI – Confidence Interval

Figure 7.2: Differences in mortality between poor, average and excellent SRH over 10 years





Figure 7.3: Differences in graft loss between poor, average and excellent SRH over 10 years



Discussion

In this study we explored SRH in an early period after KT as a predictor of mortality and graft loss in kidney transplant recipients at up to 10 years follow-up. In addition to SRH, age and eGFR were predictors of patient mortality. The predictors for graft loss were SRH, chronic renal allograft dysfunction and new-onset diabetes mellitus after transplantation. In line with our findings, Kutner et al. showed that physical functioning might be used together with factors in pre-transplant recipients to prove their potential risk for morbidity, graft loss and mortality after KT.²¹

In the Cox regression model for mortality we found that patients with a higher eGFR had a significantly reduced mortality hazard ratio. On the other hand, average SRH was associated with a 4-fold higher and poor SRH with an 11-fold higher risk of mortality. In the Cox regression model for graft loss, we found that chronic renal allograft dysfunction, new-onset diabetes mellitus after KT and average SRH were associated with a 3-fold higher risk of graft loss, and poor SRH was associated with a 6-fold higher risk of graft loss. Much like our findings of the risk ratio for graft loss, many studies have explored the idea that graft rejection and new-onset diabetes mellitus increased the graft loss risk ratio.^{9,22} Surprisingly, cardiovascular disease was not associated with mortality or graft loss in our sample, which is probably due to under-diagnosis, as we relied on evidence in the medical records and did not actively evaluate patients' comorbidity.

To our knowledge, when searching the literature we found no study exploring whether there is an association of SRH with mortality and graft loss in kidney transplant recipients. Tanikella et al. in their study with 3 years follow-up explored whether liver transplanted patients with lower self-reported physical quality of life had an increased risk of mortality.²³ So far, studies on dialysis patients have established a significant connection between poorer SRH and higher risk of mortality.^{1,3,4} Spiegel et al. in their systematic review explored a close connection between poor well-being, morbidity and mortality in patients with ESRD.²⁴ These findings are in line with our results. Thus, SRH might reflect several adverse conditions and therefore may provide additional information on a patient's risk, independent of demographic, socioeconomic and clinical risk factors for mortality¹⁵ and graft loss.

Strengths and limitations

The strength of this study is the prospective follow-up for 10 years, which enabled us to explore SRH and the others factors as predictors of mortality and graft loss after KT. Moreover, all consecutive patients originating from one major transplant centre in Slovakia over a number of years were asked to participate in the study to prevent selection bias.

Missing data is a limitation of this study; on the other hand, there were no differences in age and gender between respondents and non-respondents, and no differences in age, gender, mortality and graft loss between recipients who provided complete and incomplete data. The variable observation period between minimum and maximum (1 and 10 years) is also a limitation. The SRH interviews were not conducted immediately after transplantation to prevent false findings due to perioperative stress. Therefore, patients who died or lost their transplanted kidney before the first 3 months after KT were not incorporated into the study. It could be of interest to control for a potential effect of pre-transplantation SRH, as it may predict mortality and graft loss as well. Our findings might only be used partially due to their potential lack of generalizability, as we used data from a single center sample.

Policy implications

SRH could be used as an inexpensive and swift predictor of risky kidney transplant recipients. Patients with poor SRH might undergo relevant clinical as well as laboratory assessment and/or treatment to reduce their high risk of mortality and graft loss.

Recommendations for further research

Results must be verified in a larger multicentre sample to allow for generalization. We could then verify whether SRH after KT remains a predictor of mortality and graft loss in kidney transplant recipients, or whether in a longer period after KT other variables become important. Furthermore, the pathways between psychological, physical and medical determinants associated with SRH, mortality and graft loss should be studied.

Conclusion

Poor and average SRH in an early period after transplantation indicated a higher risk of mortality and graft loss at up to 10 years follow-up. Poor SRH is associated with an 11-fold higher risk of mortality and a 6-fold higher risk of graft loss compared to excellent SRH at up to 10 years follow-up. Patients with poor SRH might undergo relevant clinical as well as laboratory assessment and/or treatment to reduce their high risk of mortality and graft loss.

References

1. Thong MSY, Kaptein AA, Benyamini Y, Krediet RT, Boeschoten EW, Dekker FW. Association between a Self-Rated Health Question and Mortality in Young and Old Dialysis Patients: A Cohort Study. *Am J Kidney Dis.* 2008; 52: 111.
2. DeSalvo KB, Bloser N, Reynolds K, He J, Muntner P. Mortality Prediction with a Single General Self-Rated Health Question. A Meta-Analysis. *J Gen Intern Med.* 2006; 21: 267.
3. Mapes DL, Lopes AA, Satayathum S, et al. Health-Related Quality of Life as a Predictor of Mortality and Hospitalization: The Dialysis Outcomes and Practice Patterns Study (DOPPS). *Kidney Int.* 2003; 64: 339.
4. Tsai YC. Quality of Life Predicts Risks of End-Stage Renal Disease and Mortality in Patients with Chronic Kidney Disease. *Nephrol Dial Transpl.* 2010; 25: 1621.
5. Nissenson A, Chertow G, Farid M, Singh I, Van Oijen M, Desai A. The Relationship between Laboratory-Based Outcome Measures and Mortality in End-Stage Renal Disease: A Systematic Review. *Hemodial Int.* 2009; 13: 347.
6. Sirivongs D, Liawnoraset W, Pongskul C, Reungjui S. Graft Survival Analysis in Kidney Transplantation: A 12-Year Experience in a Thai Medical Center. *Transpl Proc.* 2004; 36: 2034.
7. Djamali A, Samaniego M, Torrealba J, Pirsch J, Muth BL. Increase in Proteinuria >200 mg/g After Late Rejection is Associated with Poor Graft Survival. *Nephrol Dial Transpl.* 2010; 25: 1300.
8. Schold JD, Srinivas TR, Howard RJ, Jamieson IR, Meier-Kriesche HU. The Association of Candidate Mortality Rates with Kidney Transplant Outcomes and Center Performance Evaluations. *Transplantation.* 2008; 85: 1.
9. Nemati E. Prediction of Inpatient Survival and Graft Loss in Rehospitalized Kidney Recipients. *Transplant Proc.* 2007; 39: 974.
10. Ravindran V. Post Transplant Diabetes Mellitus After Renal Transplantation: The Emerging Clinical Challenge. *Yonsei Med J.* 2004; 45: 1059.
11. Novak M. Depressive Symptoms and Mortality in Patients After Kidney Transplantation: A Prospective Prevalent Cohort Study. *Psychosom Med.* 2010; 72: 527.
12. Wolfe RA, Ashby VB, Milford EL, et al. Comparison of Mortality in all Patients on Dialysis, Patients on Dialysis Awaiting Transplantation, and Recipients of a First Cadaveric Transplant. *N Engl J Med.* 1999; 341: 1725.
13. Laupacis A, Keown P, Pus N, et al. A Study of the Quality of Life and Cost-Utility of Renal Transplantation. *Kidney Int.* 1996; 50: 235.
14. Ware JE, Jr., Sherbourne CD. The MOS 36-Item Short-Form Health Survey (SF-36). I. Conceptual Framework and Item Selection. *Med Care.* 1992; 30: 473.
15. McEwen LN. Are Health-Related Quality-of-Life and Self-Rated Health Associated with Mortality? Insights from Translating Research into Action for Diabetes (TRIAD). *Primary Care Diabet.* 2009; 3: 37.
16. Wight JP, Edwards L, Brazier J, Walters S, Payne JN, Brown CB. The SF36 as an Outcome Measure of Services for End Stage Renal Failure. *Qual Health Care.* 1998; 7: 209.
17. Franke GH, Reimer J, Philipp T, Heemann U. Aspects of Quality of Life through End-Stage Renal Disease. *Qual Life Res.* 2003; 12: 103.
18. Gomez-Besteiro MI, Santiago-Perez MI, Alonso-Hernandez A, Valdes-Canedo F, Rebollo-Alvarez P. Validity and Reliability of the SF-36 Questionnaire in Patients on the Waiting List for a Kidney Transplant and Transplant Patients. *Am J Nephrol.* 2004; 24: 346.
19. Levey AS, Stevens LA. Estimating GFR using the CKD Epidemiology Collaboration (CKD-EPI) Creatinine Equation: More Accurate GFR Estimates, Lower CKD Prevalence Estimates, and Better Risk Predictions. *Am J Kidney Dis.* 2010;55(4):622-627.
20. Sis B, Mengel M, Haas M, et al. Banff '09 Meeting Report: Antibody Mediated Graft

- Deterioration and Implementation of Banff Working Groups. *J Am Soc Nephrol.* 2010; 10: 464.
21. Zhang R, Bowles T, Paintner P, Kutner N. Pretransplant physical functioning and kidney patients' risk for posttransplantation hospitalization/death: Evidence from a national cohort. *Clin J Am Soc Nephrol.* 2006; 1: 837.
 22. Cole EH, Johnston O, Rose CL, Gill JS. Impact of Acute Rejection and New-Onset Diabetes on Long-Term Transplant Graft and Patient Survival. *Clin J Am Soc Nephrol.* May 2008; 3: 814.
 23. Tanikella R, Kawut SM, Brown RS, et al. Health-Related Quality of Life and Survival in Liver Transplant Candidates. *Liver Transpl.* 2010; 16: 238.
 24. Spiegel BMR, Melmed G, Robbins S, Esrailian E. Biomarkers and Health-Related Quality of Life in End-Stage Renal Disease: A Systematic Review. *Clin J Am Soc Nephrol.* 2008; 3: 1759.