

University of Groningen

Methylated RASSF1A in malignant peripheral nerve sheath tumors identifies neurofibromatosis type 1 patients with inferior prognosis

Danielsen, Stine A.; Lind, Guro E.; Kolberg, Matthias; Holand, Maren; Bjerkehagen, Bodil; Hall, Kirsten Sundby; van den Berg, Eva; Mertens, Fredrik; Smeland, Sigbjorn; Picci, Piero

Published in:
Neuro-Oncology

DOI:
[10.1093/neuonc/nou140](https://doi.org/10.1093/neuonc/nou140)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Danielsen, S. A., Lind, G. E., Kolberg, M., Holand, M., Bjerkehagen, B., Hall, K. S., van den Berg, E., Mertens, F., Smeland, S., Picci, P., & Lothe, R. A. (2015). Methylated RASSF1A in malignant peripheral nerve sheath tumors identifies neurofibromatosis type 1 patients with inferior prognosis. *Neuro-Oncology*, 17(1), 63-69. <https://doi.org/10.1093/neuonc/nou140>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Methylated *RASSF1A* in malignant peripheral nerve sheath tumors identifies neurofibromatosis type 1 patients with inferior prognosis

Stine A. Danielsen, Guro E. Lind, Matthias Kolberg, Maren Høland, Bodil Bjerkehagen, Kirsten Sundby Hall, Eva van den Berg, Fredrik Mertens, Sigbjørn Smeland, Piero Picci, and Ragnhild A. Lothe

Department of Cancer Prevention, Institute for Cancer Research, Oslo University Hospital – The Norwegian Radium Hospital, Oslo, Norway (S.A.D., G.E.L., M.K., M.H., R.A.L.); Centre for Cancer Biomedicine, Faculty of Medicine, University of Oslo, Oslo, Norway (S.A.D., G.E.L., M.K., M.H., R.A.L.); Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway (M.H., S.S.); Department of Biosciences, Faculty of Mathematics and Natural Sciences, University of Oslo, Oslo, Norway (S.A.D., G.E.L., R.A.L.); Department of Pathology (B.B.), Division of Diagnostics and Intervention and Department of Oncology, Division of Cancer, Surgery and Transplantation, Oslo University Hospital – The Norwegian Radium Hospital, Oslo, Norway (K.S.H., S.S.); Department of Medical Genetics, University Hospital of Groningen, The Netherlands (E.v.d.B.); Department of Clinical Genetics, Skåne University Hospital, Lund, Sweden (F.M.); Laboratory of Oncologic Research of the Istituto Ortopedico Rizzoli, Bologna, Italy (P.P.)

Corresponding Author: Ragnhild A. Lothe, PhD, Department of Cancer Prevention, Institute for Cancer Research, Oslo University Hospital – The Norwegian Radium Hospital, P.O. Box 4950 Nydalen, NO-0424 Oslo, Norway (ragnhild.a.lothe@rr-research.no).

Background. Malignant peripheral nerve sheath tumor (MPNST) is a rare and highly aggressive disease with no evidence of effect from adjuvant therapy. It is further associated with the hereditary syndrome neurofibromatosis type 1 (NF1). Silencing of the tumor suppressor gene *RASSF1A* through DNA promoter hypermethylation is known to be involved in cancer development, but its impact in MPNSTs remains unsettled.

Methods. The *RASSF1A* promoter was analyzed by methylation-specific PCR in 113 specimens, including 44 NF1-associated MPNSTs, 47 sporadic MPNSTs, 21 benign neurofibromas, and 1 nonneoplastic nerve sheath control.

Results. *RASSF1A* methylation was found only in the malignant samples (60%) and identified a subgroup among patients with NF1-associated MPNST with a poor prognosis. These patients had a mean 5-year disease-specific survival of 27.3 months (95% CI: 17.2–37.4) versus 47.4 months (95% CI: 37.5–57.2) for NF1 patients with unmethylated promoters, $P = 0.014$. In multivariate Cox regression analysis, methylated *RASSF1A* remained an adverse prognostic factor independent of clinical risk factors, $P = .013$ (hazard ratio: 5.2; 95% CI: 1.4–19.4).

Conclusion. A considerable number of MPNST samples display hypermethylation of the *RASSF1A* gene promoter, and for these tumors, this is the first molecular marker that if validated can characterize a subgroup of patients with inferior prognosis, restricted to individuals with NF1.

Keywords: methylation, MPNST, NF1, *RASSF1A*, survival.

Malignant peripheral nerve sheath tumors (MPNSTs) are classified as soft tissue sarcomas and are believed to originate from the neural crest-derived Schwann cells or their precursors.¹ About half of these tumors arise in patients with the hereditary disorder neurofibromatosis type 1 (NF1) caused by germline mutations in the *NF1* gene.¹ The lifetime risk of developing MPNSTs among NF1 patients is about 10%,² whereas the general population risk is very low. MPNSTs are highly aggressive and invasive, with a resulting poor 5-year

survival rate of 20%–50%.^{3,4} The only consensus therapy for MPNST patients is surgery, whereas adjuvant chemo- and/or radiotherapy is offered to selected patients. However, the additional treatment has not improved the survival rates.⁵ Some molecular markers have been suggested to carry prognostic information,^{5–9} but none of them have so far been implemented in the clinic. However, evaluation of molecular biomarkers with prognostic value independent of high-risk clinical factors ought to be considered as routine

Received 28 January 2014; accepted 16 June 2014

© The Author(s) 2014. Published by Oxford University Press on behalf of the Society for Neuro-Oncology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Table 1. Genes reported analyzed for DNA promoter methylation in MPNSTs and neurofibromas

Gene	MPNST	Neurofibroma	Reference
<i>CASP8</i>	0/1	0/18	44
<i>DAPK</i>	0/3	0/18	44
	0/17		25
<i>FOXD3</i> ^a	10		26
<i>GSTP1</i>	0/3	0/18	44
	1/17		25
<i>MEST</i> ^a	10		26
<i>MGMT</i>	1/3	8/18	44
	3/17		25
<i>MLH1</i>	3/17		25
<i>MSH2</i> ^b	0/17		25
	10		26
<i>NF1</i> ^a	0/5	0/16	45
	0/3	0/20	46
	0/10	12/18	47
	10		26
<i>PTEN</i>	5/17		25
	11/26	3/24	48
<i>p14/CDKN2A</i> ^a	0/19		12
	1/3	0/18	44
	0/17		25
	2/16		49
	0/18		50
	10		26
<i>p15</i> ^b	0/16		49
	10		26
<i>p16/CDKN2A</i> ^a	0/19		12
	0/3	4/18	44
	0/17		25
	0/2		15
	2/16		49
	0/18		50
	10		26
<i>p73</i>	0/3	4/18	44
<i>RASSF1A</i>	3/17		25
<i>RASSF1</i> ^a	10		26
<i>RB1</i>	1/3	2/18	44
	0/17		25
<i>RUNX1</i> ^a	10		26
<i>SOX10</i> ^a	10	10	26
<i>THBS-1</i>	2/3	9/18	44
<i>TIMP-3</i>	1/3	6/18	44
<i>TP53</i>	0/3	0/18	44
<i>TSC2</i> ^a	10		26
<i>WT1</i> ^a	10		26

^aDifferentially methylated gene specifically mentioned in a genome-wide methylome analysis comparing 10 pooled MPNSTs, 10 pooled neurofibromas, and 6 pooled nonneoplastic Schwann cell samples.²⁶

^bListed as differentially methylated in the genome-wide analysis revealing 3690 genes potentially involved in MPNST development and progression.²⁶

assays guiding the selection of patients who should be offered additional treatment.

MPNSTs commonly display complex karyotypes with frequent structural and numerical alterations,^{8,10} including recurrent gains of whole or parts of chromosome arm 17q and deletions from 9p. Components of the cell cycle machinery, such as *TP53*, *CDKN2A*, and *RB1*, frequently show aberrant gene and protein expression patterns,¹¹⁻¹⁵ and growth factor receptors like *ERRB2*, *EGFR*, *KIT*, and *MET* are reported amplified.^{11,16-18} Interestingly, except for biallelic alterations of the *NF1* gene,^{19,20} few recurrent mutations have been detected in these tumors, underscored by a recent oncogene mutation survey where the 238 most frequent cancer mutations in 19 commonly activated oncogenes were analyzed in MPNST cell lines and all were found to be wild type.²¹ This emphasizes the importance of the *NF1* mutations in MPNSTs. *NF1* encodes neurofibromin, a GTPase-activating protein functioning among others as a negative regulator of Ras. Consequently, loss of *NF1* activity leads to hyperactivation of Ras and continuous signaling through the networks of both mitogen-activated protein kinase and phosphatidylinositol-3 kinase.¹ Ras is known to interact with a number of components other than *NF1*, including Ras association domain family member 1, isoform A (*RASSF1A*), which is one of the most frequently inactivated tumor suppressors identified in cancer.²² Mutations are rarely found in tumors, but reduced *RASSF1A* expression caused by DNA hypermethylation has been reported in a variety of tumor types, including cancers of the lung, liver, breast, prostate, and testicles, as well as neuroblastomas, schwannomas,^{23,24} and a few MPNSTs.^{25,26} The *RASSF1A* gene encodes a protein resembling Ras effector proteins and has been shown to play important roles in cell cycle regulation, apoptosis, and microtubule stability. The current hypothesis is that *RASSF1A*, which has no known enzyme activity, functions both as a scaffold protein and by modulating microtubule polymerization.²⁷

Epigenetic changes are frequently seen in the majority of cancer diseases, and aberrant hypermethylation of gene promoters has been established as a common and important mechanism for inactivation of tumor suppressor genes. For MPNSTs, only a limited number of DNA methylation studies have been reported, and most of them are hampered by inclusion of rather few clinical samples (overview presented in Table 1). In this study, the methylation status of *RASSF1A* was analyzed and related to the quantitative gene expression and clinical endpoints in a large European series of MPNSTs.

Materials and Methods

Specimens

Collected at 4 European centers, a total of 113 specimens from 105 patients were included in the present study. The material comprises 91 MPNSTs, 21 neurofibromas, and 1 tumor-free sciatic nerve from a chondrosarcoma patient serving as a nonneoplastic nerve sheath control (Supplementary Table S1). All specimens have been examined by at least one pathologist, and in case of uncertainty of the diagnosis, the samples were excluded from the study. The mean age at diagnosis for

patients with malignant disease was 48 years (range: 11–79) for those with sporadic tumors ($n = 47$) and 33 years (range: 14–70) for patients with NF1 ($n = 44$). The median age for patients with neurofibromas ($n = 21$) was 26 years (range: 10–75). The patients were diagnosed during 1973–2008, and samples were collected from Oslo University Hospital, Norway; Skåne University Hospital, Lund, Sweden; University Medical Center Groningen, Netherlands; and the Istituto Ortopedico Rizzoli, Bologna, Italy. The Norwegian biobank is registered at the Institute for Public Health, biobank registry no. 1781. The study was approved by the regional committee for medical and health research ethics South East, and patients were included following informed consent. Sampling, storage, and analysis of the Swedish material were approved by the Regional Ethics Committee of Lund University, following informed consent from the patients. The Dutch material was approved according to the University Medical Center Groningen, whereas the Rizzoli Institute Ethics Committee approved the Italian material, and according to institutional guidelines, adult patients or guardians for minors signed the informed consent form.

Methylation Analysis

DNA was extracted either by phenol/chloroform followed by ethanol precipitation as described by the manufacturer of the Nucleic Acid Extractor (Applied Biosystems) or by the Qiagen Allprep DNA/RNA Mini Kit according to the manufacturer's protocol. Further, the DNA was treated with sodium bisulfite using a previously described protocol^{28–30} and/or the Epitect Bisulfite Kit (Qiagen), followed by qualitative and/or quantitative methylation-specific polymerase chain reaction (MSP and/or qMSP, respectively).³¹

The MSP reactions were carried out in a total volume of 25 μ L, containing 20 ng of bisulfite-treated template, 1 \times PCR Buffer (Qiagen), 0.2 mM deoxyribonucleotide triphosphate (Roche Diagnostics), 0.8 mM of each primer, and 0.65–1 U Hot-starTaq DNA Polymerase (Qiagen). Primers were purchased from Medprobe. The sequences are listed in Supplementary Table S2 and a schematic drawing of the promoter region with primer locations is depicted in Supplementary Fig. S1. Human placental DNA (Sigma-Aldrich) treated with SssI methyltransferase (New England Biolabs) served as a positive control for the methylated reaction, whereas bisulfite-modified DNA from normal lymphocytes was used as positive control for the unmethylated reaction. Water served as a negative control for both reactions. The PCR products were separated using 2% agarose gels (BioRad) stained with ethidium bromide (Sigma-Aldrich) and visualized by UV light using a GeneGenius gel documentation system (Syngene). The results were independently scored by visual inspection by 2 of the authors, and all methylated samples were verified by a second, independent round of MSP.

The RASSF1A gene promoter was further subjected to qMSP analysis using a protocol previously described.³² Primers and probe sequences are listed in Supplementary Table S2 and illustrated in Supplementary Fig. S1. To normalize for potential variations in the amount of input DNA (30 ng of bisulfite-treated template), ALUC4 was used as an internal reference.³³ DNA methylation was calculated as percent of methylated reference

(PMR) using the following equation:

$$\left[\frac{(RASSF1A/ALUC4)^{\text{sample}}}{(RASSF1A/ALUC4)^{\text{positive control}}} \right] \times 100.$$

Samples with PMR values exceeding the highest values among the benign neurofibromas and the nonneoplastic nerve sheath control (ie, PMR > 0) were scored as methylation positive.

Gene Expression Analysis of RASSF1

Total RNA was isolated from 5 neurofibromas and 27 MPNSTs using Trizol (Invitrogen) and converted to cDNA using the High Capacity RNA-to-DNA Kit (Applied Biosystems) according to the manufacturer's protocol.

Twenty nanograms of cDNA was amplified in a 20- μ L reaction volume containing 1 \times TaqMan Universal PCR Mastermix (Applied Biosystems) and predesigned 1 \times TaqMan Gene Expression Assays (Applied Biosystems; RASSF1, Hs 00200394_m1; RASSF1 isoform A, Hs 00945257_m1) using the following PCR program: 50°C for 2 min, 95°C for 10 min, then 45 cycles of 95°C for 15 s followed by 60°C for 1 min. The quantitative gene expression of RASSF1/RASSF1A was measured in real time using the 7900 HT Sequence Detection System (TaqMan, Applied Biosystems). All samples were analyzed in triplicate, and the median value was used for further data analyses. Universal Human Reference RNA (a mixture of total RNA from 10 different cell lines; Agilent) was used to generate a standard curve, and the quantitative expression levels of RASSF1 and RASSF1A were normalized against the mean value of the endogenous controls ACTB (Hs99999903_m1) and GUSB (Hs99999908_m1).

Statistics

Statistical analyses were performed using the software package PASW version 18.0 (SPSS). Fisher's exact test was used for assessment of associations between categorical variables, while the Mann-Whitney-Wilcoxon test was used to examine association between the methylation status of RASSF1A and continuous variables. Associations to patient outcome was evaluated only for patients with available tissue from the primary tumor, and disease-specific survival was used as the endpoint where deaths of MPNST were considered events and patients who died of other causes were censored at the time of death. All MPNST cases included in the present study were monitored closely at the same institution where they were treated, thus the cause of death could be identified with great certainty. The survival curves were generated and compared using the Kaplan–Meier method with the Breslow test, and Cox regression with the Wald test was used to generate and test uni- and multivariate proportional hazards. $P \leq .05$ was considered statistically significant.

Results

Promoter Methylation of RASSF1A in MPNSTs Related to Clinicopathological Variables

In an initial (qualitative) MSP screening study of several candidate genes (not shown), the RASSF1A promoter was found

methylated in 23 of 44 MPNST samples (52%), in contrast to complete absence of methylation in 9 neurofibromas and a normal sciatic nerve (Supplementary Table S1). No associations were found between *RASSF1A* methylation and clinicopathological variables, including NF1 status, grade, localization, or size of the tumors, or patients' age, sex, or nationality.

Due to interesting results from survival analyses, we subsequently reanalyzed the *RASSF1A* gene promoter by qMSP and expanded the sample series to a total of 91 MPNSTs, 21 neurofibromas, and the sciatic nerve. Sixty percent of the MPNSTs were methylated, whereas all neurofibromas and the sciatic nerve remained unmethylated (Supplementary Table S1). The methylation frequency was slightly, but not significantly, higher among the NF1 patients compared with the non-NF1 patients (66% and 54%, respectively; $P = .3$). In concordance with the results from the qualitative analysis, no associations were found between the presence of quantitatively measured methylation of the *RASSF1A* promoter and clinicopathological variables (Supplementary Table S3).

RASSF1A Methylation Compared With Quantitative Gene Expression

For samples where RNA was obtainable ($n = 27$), quantitative gene expression measurements of 2 assays amplifying different transcripts of the *RASSF1* gene were performed. For MPNSTs, there was a strong association between promoter hypermethylation and reduced gene expression, $P = .001$ for the most specific assay, amplifying only *RASSF1A* (Fig. 1). This effect was significant both for NF1 ($n = 15$, $P = .03$) and for non-NF1 patients ($n = 12$, $P = .003$). Both MPNSTs with methylated and

unmethylated *RASSF1A* promoter had lower expression than the benign neurofibromas ($P = .0001$ and $P = .05$, respectively), suggesting that additional mechanisms are involved in silencing the *RASSF1A* gene in MPNST.

For the MPNSTs, a borderline significance was seen with the second assay, between promoter hypermethylation and reduced gene expression ($P = .06$; data not shown). In contrast to the first assay, the second assay amplified several transcript variants of the *RASSF1* gene.

Methylation of *RASSF1A* in MPNSTs Related to Survival

In the initial MSP (qualitative) analyses, follow-up data for 36 patients with primary tumors were available, and stratification




Fig. 1. The gene expression of *RASSF1A* (assay Hs 00945257_m1 detecting NM_007182.4, *RASSF1A*) is significantly downregulated in MPNSTs with promoter hypermethylation compared with tumors without *RASSF1A* methylation, and also in MPNSTs compared with neurofibromas independently of methylation.




Fig. 2. Disease-specific survival based on *RASSF1A* promoter methylation status for (A) patients with NF1-associated MPNSTs and (B) patients with sporadic disease.

according to NF1 status suggested that *RASSF1A* methylation could distinguish a subgroup of NF1 patients with worse disease-specific 5-year survival ($n = 20$, $P = .011$). This could be seen neither among patients with sporadic disease ($n = 16$, $P = .68$) nor in the unstratified MPNST patient cohort ($n = 36$, $P = .11$) (data not shown). Based on these initial findings, we subsequently expanded the sample series and used a quantitative and more standardized analysis, qMSP. For this analysis, long-term follow-up data were available for 60 specimens with primary tumor, out of the 90 MPNST patients included, and we confirmed that NF1 patients with *RASSF1A* methylated tumors ($n = 20$) had a significantly worse prognosis than did patients with unmethylated tumors ($n = 12$, $P = .014$; Fig. 2A). The mean disease-specific survival for NF1 patients with methylation was 27.3 months (95% CI: 17.2–37.4) compared with 47.4 months (95% CI: 37.5–57.2) for NF1 patients without methylated tumors. No prognostic value of *RASSF1A* could be found for patients with sporadic disease ($n = 28$, $P = .85$; Fig. 2B).

A multivariate Cox regression analysis revealed that methylation status of *RASSF1A* was the strongest predictor of disease-specific survival (hazard ratio: 5.2; 95% CI: 1.4–19.4; $P = .013$) compared with tumor size, metastasis at time of diagnosis, and tumor location ($P = .052$, $P = .054$, and $P = .076$, respectively; Table 2).

Discussion

The literature on DNA methylation in MPNSTs is limited. Studies on a handful of genes have been reported (Table 1) in addition to a recent paper where the methylomes of 10 pooled benign and 10 pooled malignant tumors were compared.²⁶ Here, we add a detailed study analyzing the promoter methylation of *RASSF1A*.

In a large European series of MPNSTs we have demonstrated *RASSF1A* to be commonly methylated in this malignancy (60%), and at a frequency considerably higher than previously reported (18%).²⁵ All normal and benign samples were

unmethylated, underscoring that *RASSF1A* promoter methylation is cancer specific. Furthermore, expression of the *RASSF1A* gene was significantly reduced in MPNST samples with promoter hypermethylation compared with unmethylated samples. This indicates that methylation causes reduced gene activity, or gene silencing, which is in agreement with reports on other cancer types.^{24,34} Notably, however, also the MPNSTs with unmethylated *RASSF1A* promoter had significantly lower gene expression than in benign tumors, indicating that additional mechanisms of gene regulation are involved. Regulation of microtubules has been suggested to be one of the essential tumor suppressor activities of *RASSF1A*, thereby ensuring and maintaining chromosomal stability.^{22,35} The impaired function of *RASSF1A* in MPNST might contribute to the increased DNA copy number variation and complex karyotypes that are key characteristics of MPNSTs.

RASSF1A methylation has been suggested to be preferentially present in the most aggressive tumors of various kinds,^{34,36} suggesting that it might serve as a marker for tumor progression and metastasis. In the current series of MPNSTs, however, presence of *RASSF1A* methylation was equally distributed among primary, relapsed, and metastatic lesions and between individuals with and without NF1. It was also independent of tumor size and site (extremities or trunk), as well as the patient's country of origin, age, and gender.

Interestingly, survival analyses showed that *RASSF1A* methylation in MPNSTs was a surrogate marker for poor prognosis among NF1 patients only and independently of clinical risk factors such as tumor size and metastasis. To the best of our knowledge, no molecular marker in MPNSTs has previously been shown to discriminate consistently between inferior and good prognosis selectively for NF1 patients. Clinical factors such as large tumor size, incomplete resection, and distant metastasis are shown to be associated with inferior prognosis,^{3,5} which was also true for the present study. However, the multivariate survival analyses identified *RASSF1A* methylation as the strongest predictor of outcome among these factors, emphasizing the clinical potential of our finding. Biomarkers that are able to predict prognosis could be valuable in decision making

Table 2. Prognostic factors for 5-year disease-specific survival of NF1 patients with MPNST

Parameter	No. of Patients	Univariate			Multivariate		
		HR	95% CI	<i>P</i>	HR	95% CI	<i>P</i>
<i>RASSF1A</i>		2.899	1.032–8.143	.04			.013
Unmethylated	12				Ref		
Methylated	20				5.235	1.413–19.397	
Tumor size, cm ^o	30	1.053	0.992–1.118	.09	1.089	0.999–1.187	.05
Tumor location		0.936	0.380–2.309	.87			.08
Nonextremities	15				Ref		
Extremities	17				0.336	0.101–1.121	
Metastasis at time of diagnosis		4.299	1.458–12.677	.008			.05
No	27				Ref		
Yes	5				3.976	0.974–16.233	

Abbreviations: HR, hazard ratio; Ref, reference category.

^oContinuous variable.

when considering treatment alternatives. In the case of MPNSTs, one could argue that NF1 patients with methylated *RASSF1A*, predicting inferior prognosis, would benefit from an extended follow-up protocol and adjuvant treatment. Although our study includes a large number of specimens, taking into account the rareness of the disease and the fact that the findings seem valid only for NF1 patients with MPNST, it results in quite small statistical categories. Thus, the prognostic value of *RASSF1A* promoter methylation needs to be validated in an independent sample series. Nevertheless, since methylation of the same gene has been suggested as a prognostic marker for various tumor types, including different sarcomas,^{37–43} the methylation status of *RASSF1A* holds great potential to differentiate patients with both NF and MPNST into groups of good and poor prognosis.

Supplementary Material

Supplementary material is available online at Neuro-Oncology (<http://neuro-oncology.oxfordjournals.org/>).

Funding

This work was supported by grants from the Norwegian Cancer Society (PR-2006-0442 to R.A.L., financing S.A.D. as postdoctoral fellow, PR-2008-0163 to G.E.L., PR-51260-2012 to R.A.L., financing M.K. as a staff scientist) and the Faculty of Medicine, University of Oslo (to R.A.L., financing M.H. as a Ph.D student).

Acknowledgments

We are grateful to Hilde Honne and Trude Holmeide Ågesen for excellent technical assistance.

Conflict of interest statement. The authors declare that they have no conflict of interest.

References

- Carroll SL. Molecular mechanisms promoting the pathogenesis of Schwann cell neoplasms. *Acta Neuropathol.* 2012;123(3):321–348.
- Evans DG, Huson SM, Birch JM. Malignant peripheral nerve sheath tumours in inherited disease. *Clin Sarcoma Res.* 2012;2(1):17–12.
- Anghileri M, Miceli R, Fiore M, et al. Malignant peripheral nerve sheath tumors: prognostic factors and survival in a series of patients treated at a single institution. *Cancer.* 2006;107(5):1065–1074.
- Kolberg M, Holand M, Agesen TH, et al. Survival meta-analyses for >1800 malignant peripheral nerve sheath tumor patients with and without neurofibromatosis type 1. *Neuro Oncol.* 2013;15(2):135–147.
- Zou C, Smith KD, Liu J, et al. Clinical, pathological, and molecular variables predictive of malignant peripheral nerve sheath tumor outcome. *Ann Surg.* 2009;249(6):1014–1022.
- Skotheim RI, Kallioniemi A, Bjerkhagen B, et al. Topoisomerase-II alpha is upregulated in malignant peripheral nerve sheath tumors and associated with clinical outcome. *J Clin Oncol.* 2003;21(24):4586–4591.
- Brekke HR, Kolberg M, Skotheim RI, et al. Identification of p53 as a strong predictor of survival for patients with malignant peripheral nerve sheath tumors. *Neuro-oncol.* 2009;11:514–528.
- Brekke HR, Ribeiro FR, Kolberg M, et al. Genomic changes in chromosomes 10, 16, and X in malignant peripheral nerve sheath tumors identify a high-risk patient group. *J Clin Oncol.* 2010;28(9):1573–1582.
- Torres KE, Zhu QS, Bill K, et al. Activated MET is a molecular prognosticator and potential therapeutic target for malignant peripheral nerve sheath tumors. *Clin Cancer Res.* 2011;17(12):3943–3955.
- Upadhyaya M, Spurlock G, Thomas L, et al. Microarray-based copy number analysis of neurofibromatosis type-1 (NF1)-associated malignant peripheral nerve sheath tumors reveals a role for Rho-GTPase pathway genes in NF1 tumorigenesis. *Hum Mutat.* 2012;33(4):763–776.
- Rahrman EP, Watson AL, Keng VW, et al. Forward genetic screen for malignant peripheral nerve sheath tumor formation identifies new genes and pathways driving tumorigenesis. *Nat Genet.* 2013;45(7):756–766.
- Aagesen TH, Florenes VA, Molenaar WM, et al. Expression patterns of cell cycle components in sporadic and neurofibromatosis type 1-related malignant peripheral nerve sheath tumors. *J Neuropathol Exp Neurol.* 2005;64(1):74–81.
- Berner JM, Sorlie T, Mertens F, et al. Chromosome band 9p21 is frequently altered in malignant peripheral nerve sheath tumors: studies of CDKN2A and other genes of the pRB pathway. *Genes Chromosomes Cancer.* 1999;26(2):151–160.
- Kourea HP, Cordon-Cardo C, Dudas M, et al. Expression of p27(kip) and other cell cycle regulators in malignant peripheral nerve sheath tumors and neurofibromas: the emerging role of p27(kip) in malignant transformation of neurofibromas. *Am J Pathol.* 1999;155(6):1885–1891.
- Nielsen GP, Stemmer-Rachamimov AO, Ino Y, et al. Malignant transformation of neurofibromas in neurofibromatosis 1 is associated with CDKN2A/p16 inactivation. *Am J Pathol.* 1999;155(6):1879–1884.
- Perrone F, Da RL, Orsenigo M, et al. PDGFRA, PDGFRB, EGFR, and downstream signaling activation in malignant peripheral nerve sheath tumor. *Neuro Oncol.* 2009;11(6):725–736.
- Holtkamp N, Malzer E, Zietsch J, et al. EGFR and erbB2 in malignant peripheral nerve sheath tumors and implications for targeted therapy. *Neuro Oncol.* 2008;10(6):946–957.
- Holtkamp N, Okuducu AF, Mucha J, et al. Mutation and expression of PDGFRA and KIT in malignant peripheral nerve sheath tumors, and its implications for imatinib sensitivity. *Carcinogenesis.* 2006;27(3):664–671.
- Upadhyaya M, Kluwe L, Spurlock G, et al. Germline and somatic NF1 gene mutation spectrum in NF1-associated malignant peripheral nerve sheath tumors (MPNSTs). *Hum Mutat.* 2008;29(1):74–82.
- Bottillo I, Ahlquist T, Brekke H, et al. Germline and somatic NF1 mutations in sporadic and NF1-associated malignant peripheral nerve sheath tumours. *J Pathol.* 2008;217(5):693–701.
- Sun D, Tainsky MA, Haddad R. Oncogene mutation survey in MPNST cell lines enhances the dominant role of hyperactive Ras in NF1 associated pro-survival and malignancy. *Transl Oncogenomics.* 2012;5:1–7; doi:10.4137/TOG.S8830.
- Donninger H, Barnoud T, Nelson N, et al. *RASSF1A* and the rs2073498 cancer associated SNP. *Front Oncol.* 2011;1(54):11–17.

23. Agathangelou A, Cooper WN, Latif F. Role of the Ras-association domain family 1 tumor suppressor gene in human cancers. *Cancer Res.* 2005;65(9):3497–3508.
24. Donninger H, Vos MD, Clark GJ. The RASSF1A tumor suppressor. *J Cell Sci.* 2007;120(18):3163–3172.
25. Kawaguchi KI, Oda Y, Saito T, et al. DNA hypermethylation status of multiple genes in soft tissue sarcomas. *Mod Pathol.* 2006;19(1):106–114.
26. Feber A, Wilson GA, Zhang L, et al. Comparative methylome analysis of benign and malignant peripheral nerve sheath tumors. *Genome Res.* 2011;21(4):515–524.
27. Richter AM, Pfeifer GP, Dammann RH. The RASSF proteins in cancer; from epigenetic silencing to functional characterization. *Biochim Biophys Acta.* 2009;1796(2):114–128.
28. Lind GE, Kleivi K, Meling GI, et al. ADAMTS1, CRABP1, and NR3C1 identified as epigenetically deregulated genes in colorectal tumorigenesis. *Cell Oncol.* 2006;28(5–6):259–272.
29. Grunau C, Clark SJ, Rosenthal A. Bisulfite genomic sequencing: systematic investigation of critical experimental parameters. *Nucleic Acids Res.* 2001;29(13):e65.
30. Fraga MF, Esteller M. DNA methylation: a profile of methods and applications. *Biotechniques.* 2002;33(3):632–649.
31. Herman JG, Graff JR, Myohanen S, et al. Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. *PNAS.* 1996;93(18):9821–9826.
32. Lind GE, Raiborg C, Danielsen SA, et al. SPG20, a novel biomarker for early detection of colorectal cancer, encodes a regulator of cytokinesis. *Oncogene.* 2011;30(37):3967–3978.
33. Weisenberger DJ, Campan M, Long TI, et al. Analysis of repetitive element DNA methylation by MethyLight. *Nucleic Acids Res.* 2005;33(21):6823–6836.
34. Pfeifer GP, Dammann R. Methylation of the tumor suppressor gene RASSF1A in human tumors. *Biochemistry (Mosc).* 2005;70(5):576–583.
35. Vos MD, Martinez A, Elam C, et al. A role for the RASSF1A tumor suppressor in the regulation of tubulin polymerization and genomic stability. *Cancer Res.* 2004;64(12):4244–4250.
36. van der Weyden L, Adams DJ. The Ras-association domain family (RASSF) members and their role in human tumourigenesis. *Biochim Biophys Acta.* 2007;1776(1):58–85.
37. Honda S, Haruta M, Sugawara W, et al. The methylation status of RASSF1A promoter predicts responsiveness to chemotherapy and eventual cure in hepatoblastoma patients. *Int J Cancer.* 2008;123(5):1117–1125.
38. Maat W, van der Velden PA, Out-Luiting C, et al. Epigenetic inactivation of RASSF1a in uveal melanoma. *Invest Ophthalmol Vis Sci.* 2007;48(2):486–490.
39. Martins AT, Monteiro P, Ramalho-Carvalho J, et al. High RASSF1A promoter methylation levels are predictive of poor prognosis in fine-needle aspirate washings of breast cancer lesions. *Breast Cancer Res Treat.* 2011;129(1):1–9.
40. Wang J, Wang B, Chen X, et al. The prognostic value of RASSF1A promoter hypermethylation in non-small cell lung carcinoma: a systematic review and meta-analysis. *Carcinogenesis.* 2011;32(3):411–416.
41. Seidel C, Bartel F, Rastetter M, et al. Alterations of cancer-related genes in soft tissue sarcomas: hypermethylation of RASSF1A is frequently detected in leiomyosarcoma and associated with poor prognosis in sarcoma. *Int J Cancer.* 2005;114(3):442–447.
42. Tomizawa Y, Kohno T, Kondo H, et al. Clinicopathological significance of epigenetic inactivation of RASSF1A at 3p21.3 in stage I lung adenocarcinoma. *Clin Cancer Res.* 2002;8(7):2362–2368.
43. Kawai Y, Sakano S, Suehiro Y, et al. Methylation level of the RASSF1A promoter is an independent prognostic factor for clear-cell renal cell carcinoma. *Ann Oncol.* 2010;21(8):1612–1617.
44. Gonzalez-Gomez P, Bella MJ, Arjona D, et al. Aberrant CpG island methylation in neurofibromas and neurofibrosarcomas. *Oncol Rep.* 2003;10(5):1519–1523.
45. Harder A, Rosche M, Reuss DE, et al. Methylation analysis of the neurofibromatosis type 1 (NF1) promoter in peripheral nerve sheath tumours. *Eur J Cancer.* 2004;40(18):2820–2828.
46. Lujten M, Redeker S, van Noesel MM, et al. Microsatellite instability and promoter methylation as possible causes of NF1 gene inactivation in neurofibromas. *Eur J Hum Genet.* 2000;8(12):939–945.
47. Fishbein L, Eady B, Sanek N, et al. Analysis of somatic NF1 promoter methylation in plexiform neurofibromas and Schwann cells. *Cancer Genet Cytogenet.* 2005;157(2):181–186.
48. Bradtmöller M, Hartmann C, Zietsch J, et al. Impaired PTEN expression in human malignant peripheral nerve sheath tumours. *PLoS ONE.* 2012;7(11):e47595.
49. Perrone F, Tabano S, Colombo F, et al. p15INK4b, p14ARF, and p16INK4a inactivation in sporadic and neurofibromatosis type 1-related malignant peripheral nerve sheath tumors. *Clin Cancer Res.* 2003;9(11):4132–4138.
50. Endo M, Kobayashi C, Setsu N, et al. Prognostic significance of p14ARF, p15INK4b, and p16INK4a inactivation in malignant peripheral nerve sheath tumors. *Clin Cancer Res.* 2011;17(11):3771–3782.