

University of Groningen

## Male patients affected by mosaic PCDH19 mutations

de Lange, I. M.; Rump, P.; Neuteboom, R. F.; Augustijn, P. B.; Hodges, K.; Kistemaker, A. I.; Brouwer, O. F.; Mancini, G. M. S.; Newman, H. A.; Vos, Y. J.

*Published in:*  
Neurogenetics

*DOI:*  
[10.1007/s10048-017-0517-5](https://doi.org/10.1007/s10048-017-0517-5)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2017

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

de Lange, I. M., Rump, P., Neuteboom, R. F., Augustijn, P. B., Hodges, K., Kistemaker, A. I., Brouwer, O. F., Mancini, G. M. S., Newman, H. A., Vos, Y. J., Helbig, K. L., Peeters-Scholte, C., Kriek, M., Knoers, N. V., Lindhout, D., Koeleman, B. P. C., van Kempen, M. J. A., & Brilstra, E. H. (2017). Male patients affected by mosaic PCDH19 mutations: five new cases. *Neurogenetics*, *18*(3), 147-153.  
<https://doi.org/10.1007/s10048-017-0517-5>

### **Copyright**


Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Male patients affected by mosaic *PCDH19* mutations: five new cases

I. M. de Lange<sup>1</sup>  · P. Rump<sup>2</sup> · R. F. Neuteboom<sup>3</sup> · P. B. Augustijn<sup>4</sup> · K. Hodges<sup>5</sup> · A. I. Kistemaker<sup>6</sup> · O. F. Brouwer<sup>7</sup> · G. M. S. Mancini<sup>8</sup> · H. A. Newman<sup>9</sup> · Y. J. Vos<sup>2</sup> · K. L. Helbig<sup>9</sup> · C. Peeters-Scholte<sup>10</sup> · M. Kriek<sup>11</sup> · N. V. Knoers<sup>1</sup> · D. Lindhout<sup>1,12</sup> · B. P. C. Koeleman<sup>1</sup> · M. J. A. van Kempen<sup>1</sup> · E. H. Brilstra<sup>1</sup>

Received: 3 May 2017 / Accepted: 16 June 2017 / Published online: 1 July 2017  
© The Author(s) 2017. This article is an open access publication

**Abstract** Pathogenic variants in the *PCDH19* gene are associated with epilepsy, intellectual disability (ID) and behavioural disturbances. Only heterozygous females and mosaic males are affected, likely due to a disease mechanism named cellular interference. Until now, only four affected mosaic male patients have been described in literature. Here, we report five additional male patients, of which four are older than the oldest patient reported so far. All reported patients were selected for genetic testing because of developmental delay and/or epilepsy. Custom-targeted next generation sequencing gene panels for epilepsy genes were used. Clinical data were collected from medical records. All patients were mosaic in blood for likely pathogenic variants in the *PCDH19* gene. In most, clinical features were very similar to the female phenotype, with normal development before seizure onset, which occurred between 5 and 10 months of

age, clustering of seizures and sensitivity to fever. Four out of five patients had mild to severe ID and behavioural problems. We reaffirm the similarity between male and female *PCDH19*-related phenotypes, now also in a later phase of the disorder (ages 10–14 years).

**Keywords** *PCDH19* · Mosaicism · Epilepsy · Intellectual disability

## Introduction

Pathogenic variants in *PCDH19* are associated with early onset, clustered epileptic seizures often provoked by fever, intellectual disability (ID) that can be present in variable degrees and behavioural disturbances such as autistic features,

**Electronic supplementary material** The online version of this article (doi:10.1007/s10048-017-0517-5) contains supplementary material, which is available to authorized users.

✉ I. M. de Lange  
i.m.delange-2@umcutrecht.nl

<sup>1</sup> Department of Genetics, Center for Molecular Medicine, University Medical Center Utrecht, Lundlaan 6, 3584CG, Utrecht, The Netherlands

<sup>2</sup> Department of Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

<sup>3</sup> Department of Neurology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

<sup>4</sup> Department of Child Epileptology, Stichting Epilepsie Instellingen Nederland (S.E.I.N.), Heemstede, The Netherlands

<sup>5</sup> Adult and Child Neurology, Miller Children's Hospital, Long Beach, CA, USA

<sup>6</sup> Department of Paediatrics, Tjongerschans Hospital, Heerenveen, The Netherlands

<sup>7</sup> Department of Neurology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

<sup>8</sup> Department of Clinical Genetics, Erasmus University Medical Center, Rotterdam, The Netherlands

<sup>9</sup> Division of Clinical Diagnostics, Ambry Genetics, Aliso Viejo, CA, USA

<sup>10</sup> Department of Neurology, Leiden University Medical Center, Leiden, The Netherlands

<sup>11</sup> Department of Clinical Genetics, Leiden University Medical Center, Leiden, The Netherlands

<sup>12</sup> Department of Research, Stichting Epilepsie Instellingen Nederland (S.E.I.N.), Zwolle, The Netherlands

attention deficit, hyperactivity and aggression. The clinical features may resemble those of Dravet syndrome (phenotype MIM # 300088) [1–5].

*PCDH19* is located at Xq22.1 and codes for protocadherin-19, a transmembrane protein involved in neuronal organization and migration and cell-cell and cell-matrix adhesion [6–10]. It is highly expressed in the central nervous system [1]. *PCDH19*-related epilepsy shows a remarkable inheritance pattern: generally, females carrying heterozygous pathogenic variants are affected, whereas hemizygous male carriers are asymptomatic or only show psychiatric or behavioural symptoms [1, 2, 11, 12]. However, in 2009, the first affected male mosaic for a *PCDH19* pathogenic variant was described [2]. This finding gave rise to a theory of cellular interference as disease mechanism: disease occurs when two different cell populations exist (cells expressing the normal *PCDH19* protein and cells expressing a mutant form of the protein), as is true for heterozygous and mosaic pathogenic variants, but not for hemizygous pathogenic variants in males. These non-homogeneous cell populations are likely to disrupt cell-cell interactions, leading to disease [2].

Only four affected mosaic male patients have been described in literature until now, the oldest being 7 years old [2, 13, 14]. Here, we report five additional male patients with mosaic *PCDH19* mutations, of ages 2, 10, 13, 13 and 14 years old. We compared the male and female phenotypes.

## Methods

### Patients and molecular analysis

All five patients were selected for diagnostic genetic testing because of developmental delay and/or epilepsy. Genetic testing on DNA from lymphocytes was performed using a custom targeted next generation sequencing (NGS) gene panel for epilepsy genes (see online resource 1 for details). Mosaicism of *PCDH19* variants was determined based on the simultaneous presence of a variant allele and the reference allele, as *PCDH19* is located on the X-chromosome and all patients were male. The percentage of mosaicism was based on the percentage of reads that showed the alternate allele. *PCDH19* mosaic male patients from different centres were collected through personal communication between authors, meaning no structured cohort was tested. Detailed clinical data were collected from medical records. The parents of all patients gave informed consent for the publication of clinical data.

### Literature search

A literature study was carried out in the PubMed database to identify previously described patients with *PCDH19* pathogenic variants.

## Results

### Molecular analysis

All five patients carried a *PCDH19* variant in various degrees of mosaicism (patient A: c.1864G > C, p.Gly622Arg, 60%; patient B: c.840C > G, p.Tyr280\*, 22%; patient C: c.462C > G, p.Tyr154\*, 65%; patient D: c.1682C > G, p.Pro561Arg, 78%; patient E: c.799G > T p.Glu267\*, 20%, RefSeq NM\_001184880). According to American College of Medical Genetics and Genomics (ACMG) criteria, the variant of patient A was classified as likely pathogenic, and the variants of patient B–D were classified as pathogenic. The variants of patient B, C and E lead to a premature stopcodon; the variants of patient A and D were both predicted probably damaging and damaging by PolyPhen and SIFT, respectively. None of the variants was present in control databases (Database of Single Nucleotide Polymorphisms (dbSNP), NHLBI Exome Sequencing Project (ESP) and the the Exome Aggregation Consortium (ExAC) database). The Pro561Arg variant has been previously described in two affected female siblings with ID, microcephaly and seizures and was paternally transmitted [15]; the other variants are novel. All variants were confirmed de novo, except for the variant in patient D, whose parents were not tested. No other variants that could explain the phenotypes were found.

Clinical characteristics of the newly reported and previously described male patients with de novo *PCDH19* pathogenic variants are summarized and compared to those of previously published females in Table 1. See online resource for extensive clinical descriptions (online resource 2). Overall, all patients had normal development before seizure onset, occurring between 5 and 10 months of age, except for patient E. This patient had a delay in speech development, like his father, before onset of seizures and seizure onset occurred later, at 31 months. First seizure types were generalized clonic or clusters of focal or complex partial seizures. Later seizures were mainly complex partial seizures and primary or secondary generalized clonic and tonic-clonic seizures. In all five patients, seizures tended to cluster and could be provoked by fever. Four out of five patients had mild to severe ID, autism and additional behavioural problems.

## Discussion

We here report five male patients with mosaic *PCDH19* likely pathogenic variants, which raises the total number of described male patients to nine [2, 13, 14]. Four of the currently described male patients are the oldest reported so far (ages 10, 13 twice and 14 years old), which gives the opportunity to investigate whether *PCDH19*-related phenotypes evolve the same way in male and female patients. Our current findings

**Table 1** Clinical description of male and female patients carrying *PCDH19* likely pathogenic variants

Patient	A	B	C	D	E	F [2]	G [13]	H [14]	I [14]	Female patients <sup>d</sup>
Age at inclusion (years)	10	13	14	24 months	13	7	6	4	3,5	1–54
Variant <sup>a</sup>	c.1864G > C, p.Gly622Arg	c.840C > G, p.Tyr280*	c.462C > G, p.Tyr154*	c.1682C > G, p.Pro561Arg	c.799G > T, p.Glu267*	del <i>PCDH19</i>	c.605C > A, p.Ser202*	c.918C > G, p.Tyr306*	c.1352 C > T, p.Pro451Leu	<i>PCDH19</i> deletions, duplications, nonsense and missense variants
% of mosaicism	60% (blood)	22% (blood)	65% (blood)	78% (blood)	20% (blood)	100% (lymphocytes), 47% (fibroblasts)	50% (blood), 70% (buccal cells), 100% (urine sediment)	10% (lymphocytes, saliva, hair)	90% (lymphocytes, urine)	–
Technique used (number of alternate alleles/total read depth at base position)	NGS gene panel (92/153 reads)	NGS gene panel (77/380 reads)	NGS gene panel (38/59 reads)	NGS gene panel (351/450 reads)	NGS gene panel (19/93 reads)	Detection by microarray, estimation of % mosaicism by FISH (100 cells)	Detection by exome sequencing, estimation of % mosaicism by Sanger	NGS gene panel (157 reads)	NGS gene panel (135 reads)	Various
Sex	Male	Male	Male	Male	Male	Male	Male	Male	Male	Female
Exam at birth	Meconium stained	Normal	Normal	Umbilical cord around neck, quick recovery	Normal	?	Normal	?	?	Normal
Development prior to sz onset	Normal	Normal	Normal	Normal	Speech delay	Normal	Normal	Normal	Normal	Normal
Sz onset age (months)	5	10	10	7	31	12	9	9	10	3–70 (most <12)
Sz type at onset	Generalized, clonic (fs)	Generalized clonic	Cluster of CP seizures	Clusters of focal sz, generalizing to GTC and tonic seizures	Clusters of focal sz, generalizing to one hemisphere	GTC (fs, prolonged, repetitive)	Focal myoclonic, tonic-clonic, FSsG	Atetonic hypotonic seizure with hypopnea (40 min)	24 h cluster of febrile sz with fixed gaze, loss of contact, upper limb hypertonia, and jerks (30–40 s.)	Febrile, febrile SE, afebrile; GTC, generalized tonic-clonic; hemiconvulsion; focal, FSsG; CP
Later sz types	CP, secondary generalized. Focal with affective symptoms, secondary generalized. Primary generalized (fever related)	Generalized clonic	CP, secondary tonic clonic, tonic status	Focal, febrile, GTC, tonic, CP	CP, febrile	Hemiclonic, GTC, myoclonic jerks	Focal myoclonic, tonic-clonic, rapid secondary generalization	FS, focal tonic-vibratory	FS and afebrile seizure clusters, tonic. Often fearful screaming at start	FS or afebrile; generalized clonic, tonic, tonic-clonic or atonic; hemiclonic; focal, CP, FSsG, myoclonic; absences; SE
Clusters of sz	+	+	+	+	+	+	?	+	+	+
Focal sz with affective symptoms	+	–	+	–	+	?	?	?	+	+
Fever sensitivity	+	+	+	+	+	+	?	+	+	+
AEDs used and response <sup>b</sup>	VPA: -, LTG: -, CBZ:-	VPA: +	VPA:?, LEV: +/-, LZP (during clusters): -	LEV: +/-, TPM: -, OXC: +, VPA: +, PHB: +, LSM: -, Diazepam: -	VPA: +, CBZ: +/-, OXC: +/-	LEV: +/-, ZNS: +, multiple AEDs: -	LEV: +/-, ZNS: +, multiple AEDs: -	VPA: +	PHB:?, VPA: +	LEV, VPA, CLB, CLN, TPM, STP, LTG, PHB, CBZ, OXC

**Table 1** (continued)

Patient	A	B	C	D	E	F [2]	G [13]	H [14]	I [14]	Female patients <sup>d</sup>
Current AEDs <sup>b</sup>	LEV, OXC, CLB, VPA TPM	LEV, OXC, CLB, VPA	LEV	OXC, VPA, PHB, OXC LEV	OXC, OXC	?	ZNS	–	PHB, VPA	ZNS, NTZ, VGB, KBR, LZP: different responses Many different AED's, some cases no AED's
Sz outcome	1 cluster of sz per year (5–10 sz/cluster)	Last sz at age 10 years	Ongoing sz	Ongoing sz	Ongoing sz during febrile illness	Persistence of febrile sz in spite of treatment	Seizure free for 20 months	Seizure free age 14–42 months (no AEDs), since then 1 cluster of sz	4–5 clusters per year	Often seizures less frequent or seizure free at certain age (4–36 years) Mostly normal
EEG at onset	7–8 months: normal. 10–11 months: asymmetrical background activity, non-specific high voltage delta-activity occipital right > left.	Frequent generalized epileptic discharges	2 years: normal	Multiple focal discharges, right centroparietal, secondary generalization. Mild diffuse background slowing	Focal epileptic discharges left parietotemporal with generalization to one hemisphere (ictal)	?	Slower rhythm for age, mild diffuse disturbance. Infrequent right frontal, and rare left temporal sharp wave discharges, suggestive of epileptiform activity	Rare right frontotemporal sharp waves	Normal	
EEG at follow-up	3,5 years: normal	–	3 years: No epileptic activity, generalized and focal slow activity	–	32 months: interictal EEG normal	?	?	?	Bilateral centroparietal onset of seizures	Normal, interictal spikes, generalized poly-spike waves, slow waves, slow background
Last EEG	6 years: normal background activity, diffuse fast activity mainly frontal. No epileptic activity.	–	4 years: No epileptic activity, generalized and focal slow activity	–	+	?	?	?	Normal interictal	–
ID <sup>c</sup>	++ (estimated: slowed PMD, special education)	+(IQ 66 at 5 years)	++ to +++	– (only delayed speech development, but no true ID yet, based on clinical evaluation)	+(IQ 55 at 9 years)	++ to +++	+/- (IQ = 76)	+/- (GDQ 78 at 46 months, 72 at 52 months)	– (GDQ 101 at 30 months and 103 at 40 months)	– to +++
Developmental stagnation/- regression	–	–	–	–	–	?	?	–	–	Regression in some cases
Language (words/- sentences)	Delayed: words, sentences	Not delayed. First words at 12 months. Two-word	Sentences, stereotyped phrases	Mildly delayed speech development	Mildly delayed speech development	delay: words-sentences	?	?	?	Sometimes normal, often delayed, words-- sentences, in rare cases absent

**Table 1** (continued)

Patient	A	B	C	D	E	F [2]	G [13]	H [14]	I [14]	Female patients <sup>d</sup>
Behavioural/psychiatric disturbances	Autism, aggression, behavioural problems, ADHD	Autism, mood disorder	Autism spectrum disorder, anxiety	–	Behaviour problems resembling autism spectrum disorder, short attention span	Behavioural problems, autistic features	Irritability, aggression, rigidity, poor sleep, ADHD, anxiety, OCD, ODD	Compulsive and stereotyped behaviours	–	Prominent behavioural problems in most cases (autism, attention deficit, hyperactivity, aggression, emotional lability, impulsiveness, anxiety, jealousy, obsession, depression, psychogenic nonepileptic sz) Mostly normal; hypotonia, dyspraxia, ataxia or motor delay in some
Neurological examination	Crouched gait	Hypotonia	Motor delay and balance problems	Balance problems (medication induced), improving	Reduced coordination	Motor delay, ataxia	?	?	?	Usually normal (mild atrophy/cortical dysplasia is rarely reported)
MRI images (age)	Expanded perivascular spaces (8 years)	Normal (7 years)	Widened peripheral subarachnoid spaces	Normal (7 months)	Normal (2.5 years)	?	Normal	?	Normal (10 months)	
Additional comments	–	Pes plano valgus, obesity (BMI 23.3; +2.3 SD)	–	Hand food mouth disease	–	–	dysmorphic features: plagiocephaly occipital/parietal area, cupped ears, intradigital webbing of phalanges. Severe myopia	–	–	

*SZ* seizure(s), *AED* anti-epileptic drug, *PMD* psychomotor development, *GTC* generalized tonic-clonic, *CP* partial complex, *FS* fever sensitive, *SE* status epilepticus, *F5sG* focal seizure with secondary generalization, *CBZ* carbamazepine, *CLB* clobazam, *CLN* clonazepam, *KBR* potassium bromide, *LEV* levetiracetam, *LSM* lacosamide, *LTG* lamotrigine, *LZP* lorazepam, *NTZ* nitrazepam, *OXC* oxcarbazepine, *PHB* phenobarbital, *PHT* phenytoin, *STP* stiripentol, *TPM* topiramate, *VGB* vigabatrin, *VPA* valproic acid, *ZNS* zonisamide, *GDOQ* Griffiths Developmental Quotient

<sup>a</sup> RefSeq NM\_001184880

<sup>b</sup> “–” not effective, “+/-” slight effect, “+” good effect

<sup>c</sup> “–” absent, “+/-” borderline, “+” mild, “++” moderate, “+++” severe

<sup>d</sup> [2–5, 11, 12, 15–29]

confirm previously reported observations of similar clinical features in male and female patients, also for older children [2, 13, 14]. Focal seizures with affective symptoms (fearful screaming) are very common in female patients and become more prevalent with an increasing age [26]. This distinctive seizure type is also reported in three male patients [14, current study]. These similarities suggest that there are no differences in clinical consequences between phenotypes caused by postzygotic *PCDH19* pathogenic variants in mosaic males, and phenotypes caused by heterozygous *PCDH19* pathogenic variants in females. This lends further support to the hypothesis that cellular interference is the main disease mechanism, as proposed by Depienne et al. [2] The increasing number of affected mosaic male patients undermines the theory of a compensating effect by the nonparalogous *PCDH11Y* gene in male patients, as proposed by Dibbens et al. [1]. It is highly unlikely that this gene would only compensate for a complete absence of *PCDH19* in hemizygous affected males, but not for a partial loss of *PCDH19* in mosaic males.

In female patients, seizure frequency often diminishes around puberty [3, 5, 11, 28–30], possibly due to hormonal changes [30]. Our 13-year old patient has been seizure free since the age of 10 years old; our 10-year-old patient only has one cluster of seizures per year. However, our other 13- and 14-year-old patients still have ongoing seizures. Although four of our male patients are the oldest reported until now, none of them has reached adolescence yet, and the numbers are still small, making it hard to draw definite conclusions. Nevertheless, since some of the few reported male patients already show a reduction in seizure frequency with increasing age, it seems unlikely that a declining seizure frequency is exclusively occurring in females and is related to female specific hormones.

We hypothesised that males with a mutated allele percentage around 50% in the brain, who would have an inherent high level of cellular interference, may show a more severe clinical picture than males with a lower or higher percentage of mosaicism. High or low percentages of mosaicism would resemble skewed X-inactivation in female patients, which has also been suggested to lead to a milder phenotype [5, 15], although no clear correlation has been shown [25, 29]. Indeed, in our cohort, the three patients with the lowest and highest percentages of mosaicism in blood are the least severely affected (patients B, D and E), and two previously described patients, one with borderline ID and one with normal intelligence showed percentages of mosaicism of 10 and 90%, respectively (patients H and I). However, patient G shows an equal number of mutated and wild-type alleles in blood and is also mildly affected, and patient F shows no mosaicism in blood cells at all, only in fibroblasts, but has moderate to severe ID and ongoing seizures. It is thus not possible to predict the phenotype based on the percentage of mosaicism in blood, most likely because it does not necessarily equal the percentage of mosaicism in the brain.

The number of identified male patients mosaic for *PCDH19* pathogenic variants increases [2, 13, 14], probably due to our improving abilities to detect mosaicism in general by using NGS techniques. It is now clear that *PCDH19* pathogenic variants can cause epilepsy both in males and females and that mosaicism for *PCDH19* pathogenic variants in males might be more common than previously thought. Because our five described patients were gathered through personal communication between authors from different diagnostic centres, no structured cohort with clearly defined inclusion criteria was tested, which makes it difficult to estimate the frequency of mosaic pathogenic *PCDH19* variants in male patients. Since the *PCDH19* and Dravet syndrome phenotypes show many similarities, testing a cohort of *SCN1A*-negative, male Dravet syndrome patients for mosaic *PCDH19* pathogenic variants could give more insight in the true incidence. In this cohort and in males with clinical features characteristic of a *PCDH19*-related disorder, NGS techniques with high coverage should be used to look for *PCDH19* pathogenic variants, as traditional Sanger Sequencing is not sensitive enough to reliably detect mosaicism. This overview helps create more knowledge about the disease course in male patients, which is extremely relevant for counselling those affected and their families. Reporting on more (older) patients in the future is essential for establishing a good understanding of prognosis in male patients.

#### Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflicts of interest. HAN and KLH are employed by Ambry Genetics; *PCDH19* sequencing is among its commercially available tests. This study was supported by the “Stichting Vrienden WKZ” (project 1614054) on behalf of Stichting Panta Rhei. Funders had no involvement in the study design, the collection, analysis, and interpretation of data, the writing of the report and in the decision to submit the paper for publication.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

#### References

1. Dibbens LM, Tarpey PS, Hynes K et al (2008) X-linked protocadherin 19 mutations cause female-limited epilepsy and cognitive impairment. *Nat Genet* 40:776–781. doi:10.1038/ng.149.X-linked
2. Depienne C, Bouteiller D, Keren B et al (2009) Sporadic infantile epileptic encephalopathy caused by mutations in *PCDH19* resembles Dravet syndrome but mainly affects females. *PLoS Genet* 5: e1000381. doi:10.1371/journal.pgen.1000381
3. Specchio N, Marini C, Terracciano A et al (2011) Spectrum of phenotypes in female patients with epilepsy due to protocadherin

- 19 mutations. *Epilepsia* 52:1251–1257. doi:10.1111/j.1528-1167.2011.03063.x
4. Camacho A, Simón R, Sanz R et al (2012) Cognitive and behavioral profile in females with epilepsy with PCDH19 mutation: two novel mutations and review of the literature. *Epilepsy Behav* 24:134–137. doi:10.1016/j.yebeh.2012.02.023
  5. Depienne C, Leguern E (2012) PCDH19-related infantile epileptic encephalopathy: an unusual X-linked inheritance disorder. *Hum Mutat* 33:627–634. doi:10.1002/humu.22029
  6. Kim SY, Chung HS, Sun W, Kim H (2007) Spatiotemporal expression pattern of non-clustered protocadherin family members in the developing rat brain. *Neuroscience* 147:996–1021. doi:10.1016/j.neuroscience.2007.03.052
  7. Emond MR, Biswas S, Blevins CJ, Jontes JD (2011) A complex of protocadherin-19 and n-cadherin mediates a novel mechanism of cell adhesion. *J Cell Biol* 195:1115–1121. doi:10.1083/jcb.201108115
  8. Compagnucci C, Petrini S, Higurashi N et al (2015) Characterizing PCDH19 in human induced pluripotent stem cells (iPSCs) and iPSC-derived developing neurons: emerging role of a protein involved in controlling polarity during neurogenesis. *Oncotarget* 6:26804–26813. doi:10.18632/oncotarget.5757
  9. Cooper SR, Emond MR, Duy PQ et al (2015) Protocadherins control the modular assembly of neuronal columns in the zebrafish optic tectum. *J Cell Biol* 211:807–814. doi:10.1083/jcb.201507108
  10. Pederick DT, Homan CC, Jaehne EJ et al (2016) Pcdh19 loss-of-function increases neuronal migration in vitro but is dispensable for brain development in mice. *Sci Rep* 6:26765. doi:10.1038/srep26765
  11. Scheffler IE, Turner SJ, Dibbens LM et al (2008) Epilepsy and mental retardation limited to females: an under-recognized disorder. *Brain* 131:918–927. doi:10.1093/brain/awm338
  12. Van Harssel JJT, Weckhuysen S, Van Kempen MJA et al (2013) Clinical and genetic aspects of PCDH19-related epilepsy syndromes and the possible role of PCDH19 mutations in males with autism spectrum disorders. *Neurogenetics* 14:23–34. doi:10.1007/s10048-013-0353-1
  13. Thiffault I, Farrow E, Smith L et al (2016) PCDH19-related epileptic encephalopathy in a male mosaic for a truncating variant. *Am J Med Genet Part A* 170:1585–1589. doi:10.1002/ajmg.a.37617
  14. Terracciano A, Trivisano M, Cusmai R et al (2016) PCDH19-related epilepsy in two mosaic male patients. *Epilepsia* 57:e51–e55. doi:10.1111/epi.13295
  15. Depienne C, Trouillard O, Bouteiller D et al (2011) Mutations and deletions in PCDH19 account for various familial or isolated epilepsies in females. *Hum Mutat* 32:1959–1975. doi:10.1002/humu.21373
  16. Cappelletti S, Specchio N, Moavero R et al (2015) Cognitive development in females with PCDH19 gene-related epilepsy. *Epilepsy Behav* 42:36–40. doi:10.1016/j.yebeh.2014.10.019
  17. Dibbens LM, Kneen R, Bayly MA et al (2011) Recurrence risk of epilepsy and mental retardation in females due to parental mosaicism of PCDH19 mutations. *Neurology* 76:1514–1519. doi:10.1212/WNL.0b013e318217e7b6
  18. Dimova PS, Kirov A, Todorova A et al (2012) A novel PCDH19 mutation inherited from an unaffected mother. *Pediatr Neurol* 46:397–400. doi:10.1016/j.pediatrneurol.2012.03.004
  19. Gagliardi M, Annesi G, Sesta M et al (2015) PCDH19 mutations in female patients from southern Italy. *Seizure* 24:118–120. doi:10.1016/j.seizure.2014.08.010
  20. Higurashi N, Shi X, Yasumoto S et al (2012) PCDH19 mutation in Japanese females with epilepsy. *Epilepsy Res* 99:28–37. doi:10.1016/j.eplepsyres.2011.10.014
  21. Higurashi N, Nakamura M, Sugai M et al (2013) PCDH19-related female-limited epilepsy: further details regarding early clinical features and therapeutic efficacy. *Epilepsy Res* 106:191–199. doi:10.1016/j.eplepsyres.2013.04.005
  22. Hynes K, Tarpey P, Dibbens LM et al (2010) Epilepsy and mental retardation limited to females with PCDH19 mutations can present de novo or in single generation families. *J Med Genet* 47:211–216. doi:10.1136/jmg.2009.068817
  23. Jamal SM, Basran RK, Newton S et al (2010) Novel de novo PCDH19 mutations in three unrelated females with epilepsy female restricted mental retardation syndrome. *Am J Med Genet Part A* 152(A):2475–2481. doi:10.1002/ajmg.a.33611
  24. Kwong AKY, Ho ACC, Fung CW, Wong VCN (2015) Analysis of mutations in 7 genes associated with neuronal excitability and synaptic transmission in a cohort of children with non-syndromic infantile epileptic encephalopathy. *PLoS One* 10:e0126446. doi:10.1371/journal.pone.0126446
  25. Leonardi E, Sartori S, Vecchi M et al (2014) Identification of four novel PCDH19 mutations and prediction of their functional impact. *Ann Hum Genet* 78:389–398. doi:10.1111/ahg.12082
  26. Marini C, Darra F, Specchio N et al (2012) Focal seizures with affective symptoms are a major feature of PCDH19 gene-related epilepsy. *Epilepsia* 53:2111–2119. doi:10.1111/j.1528-1167.2012.03649.x
  27. Vincent AK, Noor A, Janson A et al (2012) Identification of genomic deletions spanning the PCDH19 gene in two unrelated girls with intellectual disability and seizures. *Clin Genet* 82:540–545. doi:10.1111/j.1399-0004.2011.01812.x
  28. Liua A, Xu X, Yang X et al (2016) The clinical Spectrum of female epilepsy patients with PCDH19 mutations in a Chinese population. *Clin Genet*. doi:10.1111/cge.12846
  29. Marini C, Mei D, Parmeggiani L et al (2010) Protocadherin 19 mutations in girls with infantile-onset epilepsy. *Neurology* 75:646–653. doi:10.1212/WNL.0b013e3181ed9e67
  30. Tan C, Shard C, Ranieri E et al (2015) Mutations of protocadherin 19 in female epilepsy (PCDH19-FE) lead to allopregnanolone deficiency. *Hum Mol Genet* 24:5250–5259. doi:10.1093/hmg/ddv245