

University of Groningen

## Complete Genome Sequences of Two Methicillin-Resistant *Staphylococcus haemolyticus* Isolates of Multilocus Sequence Type 25, First Detected by Shotgun Metagenomics

Couto, Natacha; Chlebowicz, Monika A; Raangs, Erwin C; Friedrich, Alex W; Rossen, John W

*Published in:*  
Genome Announcements

*DOI:*  
[10.1128/genomeA.00036-18](https://doi.org/10.1128/genomeA.00036-18)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2018

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Couto, N., Chlebowicz, M. A., Raangs, E. C., Friedrich, A. W., & Rossen, J. W. (2018). Complete Genome Sequences of Two Methicillin-Resistant *Staphylococcus haemolyticus* Isolates of Multilocus Sequence Type 25, First Detected by Shotgun Metagenomics. *Genome Announcements*, 6(14), [e00036-18]. <https://doi.org/10.1128/genomeA.00036-18>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*



# Complete Genome Sequences of Two Methicillin-Resistant *Staphylococcus haemolyticus* Isolates of Multilocus Sequence Type 25, First Detected by Shotgun Metagenomics

 Natacha Couto,<sup>a</sup> Monika A. Chlebowicz,<sup>a</sup> Erwin C. Raangs,<sup>a</sup> Alex W. Friedrich,<sup>a</sup> John W. Rossen<sup>a</sup>

<sup>a</sup>University of Groningen, University Medical Center Groningen, Department of Medical Microbiology, Groningen, Netherlands

**ABSTRACT** The emergence of nosocomial infections by multidrug-resistant *Staphylococcus haemolyticus* isolates has been reported in several European countries. Here, we report the first two complete genome sequences of *S. haemolyticus* sequence type 25 (ST25) isolates 83131A and 83131B. Both isolates were isolated from the same clinical sample and were first identified through shotgun metagenomics.

A recent worldwide increase in the number of multidrug-resistant (MDR) *Staphylococcus haemolyticus* isolates causing hospital infections has been associated with an unusual genome plasticity generated by a large number of insertion sequences (1). In particular, isolates belonging to multilocus sequence type 25 (ST25), initially detected in India (2), are known to cause keratitis and belong to clonal complex 1 (CC1). So far, ST25 isolates have been isolated in several European countries (3).

Here, we report the complete genome sequences of two coexisting ST25 *S. haemolyticus* isolates from the same clinical sample (peritoneal fluid), originally identified through shotgun metagenomics (N. Couto, L. Schuele, E.C. Raangs, et al., submitted for publication).

DNA extraction was performed with the DNeasy UltraClean microbial kit (Qiagen, Hilden, Germany). DNA concentration was measured with a Qubit 2.0 fluorometer (Life Technologies, Thermo Fisher Scientific, Waltham, MA). Both strains were sequenced using the MiSeq platform (Illumina, San Diego, CA), generating  $2 \times 250$ -bp reads, and the MinION instrument (Oxford Nanopore, Oxford, UK) in order to close the first genomes of this lineage. The Illumina and MinION reads were assembled using Unicycler v0.4.1 (4). Genome annotation was performed at the GenBank submission step using the Prokaryotic Genome Automatic Annotation Pipeline (PGAAP) from NCBI (5).

The genome of the ST25 *S. haemolyticus* 83131A isolate has a circular chromosome with 2,593,983 bp and a GC content of 32.8%, 2,449 coding sequences (CDS), 60 tRNAs, and 19 rRNAs. The genome of the ST25 *S. haemolyticus* 83131B isolate has a circular chromosome with 2,593,418 bp and a GC content of 32.8%, 2,437 CDS, 60 tRNAs, and 19 rRNAs. The 83131A isolate harbors one circular plasmid of 16,882 bp (p83131A-A), and the 83131B strain harbors two circular plasmids of 16,882 bp (p83131A-B) and 2,366 bp (p83131B).

The hybrid assembly yielded a contig with the entire genome, which was different from that obtained by using Illumina alone. This is due to the resulting short-read length, which cannot span across repetitive regions, such as transposons and insertion sequences, and which creates gaps in assemblies. Thus, our results reflect the suitability of MinION plus Illumina sequencing for full characterization of plasmids and other mobile genetic elements, which are chimeric elements that are impossible to elucidate by using short-read sequencing approaches alone (5). Through whole-genome MLST (wgMLST) analysis using Ridom SeqSphere+ v4.1.9, we were able to determine that

**Received** 16 January 2018 **Accepted** 6 February 2018 **Published** 5 April 2018

**Citation** Couto N, Chlebowicz MA, Raangs EC, Friedrich AW, Rossen JW. 2018. Complete genome sequences of two methicillin-resistant *SStaphylococcus haemolyticus* isolates of multilocus sequence type 25, first detected by shotgun metagenomics. *Genome Announc* 6:e00036-18. <https://doi.org/10.1128/genomeA.00036-18>.

**Copyright** © 2018 Couto et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Natacha Couto, [n.monge.gomes.do.couto@umcg.nl](mailto:n.monge.gomes.do.couto@umcg.nl).

the two isolates differed in 7 out of 2,321 alleles. In the NCBI database, there were only three closed full-genome sequences of *S. haemolyticus* available. The most closely related genome was from an *S. haemolyticus* ST42 strain (GenBank accession number CP011116) isolated during a nosocomial outbreak in Brazil (6), which differed in 514 alleles. The observed high genomic diversity among *S. haemolyticus* genomes shows the need for more full-genome sequences. Genomic comparisons often provide information about similarities and differences of isolates of the same species that belong to different lineages and therefore help identify factors that contribute to the emergence of successful clones.

**Accession number(s).** The GenBank genome accession numbers for the sequences described in this study are [CP024809](#) and [CP025396](#) (genome sequences for isolates 83131A and 83131B, respectively), [CP024810](#) (p83131A-A), [CP025397](#) (p83131A-B), and [CP025398](#) (p83131B).

## ACKNOWLEDGMENTS

This work was partially supported by the INTERREG VA (project 202085)-funded project EurHealth-1Health, part of a Dutch-German cross-border network supported by the European Commission, the Dutch Ministry of Health, Welfare and Sport (VWS), the Ministry of Economy, Innovation, Digitalisation and Energy of the German Federal State of North Rhine-Westphalia, and the German Federal State of Lower Saxony.

## REFERENCES

1. Czekaj T, Ciszewski M, Szewczyk M. 2015. *Staphylococcus haemolyticus*—an emerging threat in the twilight of the antibiotics age. *Microbiology* 161:2061–2068. <https://doi.org/10.1099/mic.0.000178>.
2. Panda S, Jena S, Sharma S, Dhawan B, Nath G, Singh DV. 2016. Identification of novel sequence types among *Staphylococcus haemolyticus* isolated from variety of infections in India. *PLoS One* 11:e0166193. <https://doi.org/10.1371/journal.pone.0166193>.
3. Cavanagh JP, Hjerde E, Holden MTG, Kahlke T, Klingenberg C, Flægstad T, Parkhill J, Bentley SD, Sollid JUE. 2014. Whole-genome sequencing reveals clonal expansion of multiresistant *Staphylococcus haemolyticus* in European hospitals. *J Antimicrob Chemother* 69:2920–2927. <https://doi.org/10.1093/jac/dku271>.
4. Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
5. Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity GM, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of standard operating procedures (SOPs) for (meta)genomic annotation. *OMICS* 12:137–141. <https://doi.org/10.1089/omi.2008.0017>.
6. de Almeida LM, Pires C, Cerdeira LT, de Oliveira TGM, McCulloch JA, Perez-Chaparro PJ, Sacramento AG, Brito AC, da Silva JL, de Araújo MR, Lincopan N, Martin MJ, Gilmore MS, Mamizuka EM. 2015. Complete genome sequence of linezolid-susceptible *Staphylococcus haemolyticus* Sh29/312/L2, a clonal derivative of a linezolid-resistant clinical strain. *Genome Announc* 3:e00494-15. <https://doi.org/10.1128/genomeA.00494-15>.