

University of Groningen

Social cognition and treatment in psychosis

van Donkersgoed, Rozanne Janna Margaretha

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van Donkersgoed, R. J. M. (2018). Social cognition and treatment in psychosis. [Groningen]: Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

REFERENCES

REFERENCES

- Aan het Rot, M., & Hogenelst, K. (2014). The influence of affective empathy and autism spectrum traits on empathic accuracy. *PLoS One*, 9(6), e98436.
- Addington, J., Penn, D., Woods, S.W., Addington, D., & Perkins, D.O. (2008). Facial affect recognition in individuals at clinical high risk for psychosis. *Br J Psychiatry*, 192(1), 67-8.
- Addington, J., Piskulic, D., Perkins, D., Woods, S.W., Liu, L., & Penn, D.L. (2012). Affect recognition in people at clinical high risk of psychosis. *Schizophren Res*, 1040, 87-92.
- Addington, J., Saeedi, H., & Addington, D. Facial affect recognition: a mediator between cognitive and social functioning in psychosis? *Schizophr Res*. 2006; 85(1-3): 142-50.
- Adolphs, R., Tranel, D., Damasio, A.R. (1998). The human amygdala in social judgment. *Nature*, 393, 470-474.
- Aleman, A., Kahn, R.S., & Selten, J.P. (2003). Sex differences in the risk of schizophrenia: evidence from meta-analysis. *Arch Gen Psychiatry*, 60(6), 565-71.
- Allott, K.A., Schäfer, M.R., Thompson, A., Nelson, B., Bendall, S., Bartholomeusz, C.F., Yuen, H.P., McGorry, P.D., Schlögelhofer, M., Bechdolf, A., & Amminger, G.P. (2014). Emotion recognition as a predictor of transition to a psychotic disorder in ultra-high risk participants. *Schizophren Res*, 153, 25-31.
- Amador, X.F., & David, A.S. (2004). *Insight and Psychosis: Awareness of Illness in Schizophrenia and Related Disorders* (2nd edition). New York: Oxford University Press.
- Amador, X.F., Strauss, D.H., Yale, S.A., Flaum, M.M., Endicott, J., & Gorman, J.M., (1993). Assessment of insight in psychosis. *Am J Psychiatry*, 150, 873- 879.
- Amminger, G.P., Schäfer, M.R., Klier, C.M., Schlögelhofer, M., Mossaheb, N., Thompson, A., Bechdolf, A., Allott, K., McGorry, P.D., & Nelson, B. (2012). Facial and vocal affect perception in people at ultra-high risk of psychosis, first-episode schizophrenia and healthy controls. *Early Interv Psychiatry*, 6(4), 450-4.
- Amminger, G.P., Schäfer, M.R., Papageorgiou, K., Klier, C.M., Cotton, S.M., Harrigan, S.M., Mackinnon, A., McGorry, P.D., & Berger, G.E. (2010). Long-chain omega-3 fatty acids for indicated prevention of psychotic disorders: a randomized, placebo-controlled trial. *Arch Gen Psychiatry*, 67(2), 146-154.
- Amminger, G.P., Schäfer, M.R., Papageorgiou, K., Klier, C.M., Schlögelhofer, M., Mossaheb, N., Wernneck-Rohrer, S., Nelson, B., & McGorry, P.D. (2012). Emotion recognition in individuals at clinical high-risk for schizophrenia. *Schizophren Bull*, 38(5), 1030-9.
- An, S.K., Kang, J.I., Park, J.Y., Kim, K.R., Lee, S.Y., & Lee, E. (2010). Attribution bias in ultra-high risk for psychosis and first-episode schizophrenia. *Schizophr Res*, 118(1-3), 54-61.
- An der Heiden, W., & H(ä)fner, H. (2000). The epidemiology of onset and course of schizophrenia. *Eur Arch Psychiatry*, 250(6), 292-303.
- Ascher-Svanum, H., & Whitesel, J. (1999). A randomized controlled study of two styles of group patient education about schizophrenia. *Psychiatr Serv*, 50, 926-930.
- Auquier, P., Lançon, C., Rouillon, F., & Lader, M. (2007). Mortality in schizophrenia. *Pharmacoepidemiol Drug Saf*, 16(12), 1308-12.
- Baldwin, S.A., Wampold, B.E., & Imel, Z.E. (2007). Untangling the alliance-outcome correlation: exploring the relative importance of therapist and patient variability in the alliance. *Journal of Consulting and Clinical Psychology*, 75(6), 842-852.
- Barbato, M., Liu, L., Penn, D.L., Keefe, R.S., Perkins, D.O., & Woods, S.W., & Addington, J. (2013). Social cognition as a mediator between neurocognition and functional outcome in individuals at clinical high risk for psychosis. *Schizophr Res*, 150(2-3), 542-6.
- Bargenquast, R., & Schweitzer, R.D. (2013). Enhancing sense of recovery and self-reflectivity in people with schizophrenia: a pilot study of Metacognitive Narrative Psychotherapy. *Psychol Psychother*, 87(3), 338-56.
- Barnes, T.R.E., & Paton, C. (2011). Do antidepressants improve negative symptoms in schizophrenia? *British Medical Journal*, 10, 342-349.
- Baron-Cohen, S., O'Riordan, M., Stone, V., Jones, R., & Plaisted, K., (1999). Recognition of faux pas by normally developing children and children with Asperger syndrome or high-functioning autism. *J Autism Dev Disord*, 29(5), 407-418.

- Baron-Cohen, S., Wheelwright, S., Hill, J., Raste, Y., & Plumb, I. (2001). The Reading the Mind in the Eyes Test revised version: A study with normal adults, and adults with Asperger syndrome or high-functioning autism. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 42, 241–251.
- Bartholomeusz, C.F., Whittle, S.L., Plioussis, E., Allott, K., Rice, S., Schäfer, M.R., Pantelis, C., & Amminger, G.P. (2014). Relationship between amygdala volume and emotion recognition in adolescents at ultra-high risk for psychosis. *Psychiatry Research: Neuroimaging*, 224, 159–167.
- Bateman, A.W., & Fonagy, P. (2011). *Handbook of mentalizing in mental health practice*. Washington DC: American Psychiatric Publishing.
- Beck, A.T., Baruch, E., Balter, J.M., Steer, R.A., & Warman, D.M. (2004). A new instrument for measuring insight: the Beck Cognitive Insight Scale. *Schizophr Res*, 68, 319–329.
- Bellugi, U., Adolphs, R., Cassady, C., & Chiles, M. (1999). Towards the neural basis for hypersociability in a genetic syndrome. *Neuroreport*, 10, 1653–1657.
- Biedermann, F., Frajo-Apor, B., & Hofer, A. (2012). Theory of mind and its relevance in schizophrenia. *Curr Opin Psychiatry*, 25(2), 71–75.
- Birchwood, M., Smith, J., Drury, V., Healy, J., Macmillan, F., & Slade, M.A. (1994). Self report Insight Scale for psychosis: reliability, validity and sensitivity to change. *Acta Psychiatr Scand*, 89, 62–67.
- Blakemore, S.J. (2008). The social brain in adolescence. *Nat Rev Neurosci*, 9(4), 267–77.
- Blair, R.J. (2005). Responding to the emotions of others: dissociating forms of empathy through the study of typical and psychiatric populations. *Conscious Cogn*, 14(4), 698–718.
- Bonfils, K.A., Lysaker, P.H., Minor, K.S., & Salyers, M.P. (2016). Affective empathy in schizophrenia: a meta-analysis. *Schizophr Res*, 175(1–3), 109–17.
- Bora, E., Eryavuz, A., Kayahan, B., Sungu, C., & Veznedaroglu, B. (2006). Social functioning, theory of mind and neurocognition in outpatients with schizophrenia; mental state decoding may be a better predictor of social functioning than mental state reasoning. *Psychiatry Res*, 145, 95–103.
- Bora, E., Lin, A., Wood, S.J., Yung, A.R., McGorry, P.D., & Pantelis, C. (2014). Cognitive deficits in youth with familial and clinical high risk to psychosis: a systematic review and meta-analysis. *Acta Psychiatr Scand*, 130(1), 1–15.
- Bora, E., & Pantelis, C. (2013). Theory of mind impairments in first-episode psychosis, individuals at ultra-high risk for psychosis and in first-degree relatives of schizophrenia: systematic review and meta-analysis. *Schizophr Res*, 144(1–3), 31–6.
- Bouwman, C., De Jong, K., Timman, R., Zijlstra-Vlasveld, M., Van der Feltz-Cornelis, C., Tan Swan, S., Hakkaart, & Van Roijen, L. (2013). Feasibility, reliability and validity of a questionnaire on health care consumption and productivity loss in patients with a psychiatric disorder (TiC-P). *BMC Health Serv Res*, 13, 127.
- Boyd Ritsher, J., Otilingam, P.G., & Grajales, M. (2003). Internalized stigma of mental illness: psychometric properties of a new measure. *Psychiatry Research*, 121(1), 31–49.
- Brent, B. (2009). Mentalization based psychodynamic psychotherapy for psychosis. *J Clin Psychol*, 65(8), 803–814.
- Bromley, E., & Brekke, J.S. (2010). Assessing function and functional outcome in schizophrenia. In: *Behavioral Neurobiology of Schizophrenia and its Treatment*. Springer Berlin Heidelberg, p. 3–21.
- Brüne, M. (2005). Theory of mind in schizophrenia: a review of the literature. *Schizophr Bull*, 31(1), 21–42.
- Brüne, M., Abdel-Hamid, M., Lehmkämoer, C., & Sonntag, C. (2007). Mental state attribution, neurocognitive functioning and psychopathology: what predicts poor social competence in schizophrenia best? *Schizophr Res*, 92, 151–159.
- Brüne, M., Ozgürdal, S., Ansgor, N., Von Reventlow, H.G., Peters, S., Nicolas, V., Tegenthoff, M., Juckel, C., & Lissek, S. (2011). An fMRI study of "theory of mind" in at-risk states of psychosis: comparison with manifest schizophrenia and healthy controls. *Neuroimage*, 55(1), 329–37.
- Buchanan, R.W. (2007). Persistent negative symptoms in schizophrenia: An overview. *Schizophrenia Bulletin*, 33(4), 1013–1022.

REFERENCES

- Buck, K.D., & Lysaker, P.H. (2009). Addressing metacognitive capacity in the psychotherapy for schizophrenia: a case study. *Clin Case Studies*, 8(6), 463-472.
- Buck, B., Minor, K.S., & Lysaker, P.H. (2015). Differential lexical correlates of social cognition and metacognition in schizophrenia; a study of spontaneously-generated life narratives. *Compr Psychiatry*, 58, 138-45.
- Buckley, P.F., & Stahl, S.M. (2007). Pharmacological treatment of negative symptoms of schizophrenia: therapeutic opportunity or cul-de-sac? *Acta Psychiatrica Scandinavica*, 115(2), 93-100.
- Cannon, T.D., Cadenhead, K., Cornblatt, B., Woods, S.W., Addington, J., Walker, E., Seidman, L.J., Perkins, D., Tsuang, M., McGlashan, T., & Heinssen, R. (2008). Prediction of psychosis in youth at high clinical risk: a multisite longitudinal study in North America. *Arch Gen Psychiatr*, 65, 28-37.
- Carroll, A., Fattah, S., Clyde, Z., Coffey, I., Owens, D.G.C., & Johnstone, E.C., (1998). Correlates of insight and insight change in schizophrenia. *Schizophr Res*, 35, 247-253.
- Cartwright-Hatton, S., & Wells, A. (1997). Beliefs about worry and intrusions, the meta-cognitions questionnaire and its correlates. *J Anxiety Disord*, 11(3), 279-296.
- Chan, S.H., Lee, S.W., & Chan, I.W., (2007). TRIP: a psycho-educational programme in Hong Kong for people with schizophrenia. *Occup Ther Int*, 14(2), 86-98.
- Chan, S.W., Yip, B., Tso, S., Cheng, B., & Tame, W., (2009). Evaluation of a psychoeducation program for Chinese clients with schizophrenia and their family caregivers. *Patient Educ Couns*, 75, 67-76.
- Charlton, C., Rasbash, J., Browne, W. J., Healy, M., & Cameron, B. (2017). MLwiN. Bristol: Centre for Multilevel Modelling, University of Bristol.
- Choi-Kain, L.W., & Gunderson, J.G. (2008). Mentalization: ontogeny, assessment and application in the treatment of borderline personality disorder. *Am J Psychiatr*, 165, 1127-1135.
- Cohen, J. (1977). *Statistical power analysis for the behavioral sciences*. New York: Academic Press.
- Cohen, S. (2004). Social Relationships and Health. *American Psychologist*, 59, 676-84.
- Combs, D.R., Chapman, D., Waguspack, J., Basso, M.R., & Penn, D.L. (2011). Attention shaping as a means to improve emotion perception deficits in outpatients with schizophrenia and impaired controls. *Schizophr Res*, 127(1-3), 151-6.
- Combs, D.R., & Gouvier, W.D. (2004). The role of attention in affect perception: an examination of Mirsky's four factor model of attention in chronic schizophrenia. *Schizophr Bull*, 30(4), 727-38.
- Corrigan, P.W., Garman, A., & Nelson, D. (1996). Situational feature recognition in schizophrenic outpatients. *Psychiatry Res*, 62(3), 251-7.
- Cunningham-Owens, D.G., Carroll, A., Fattah, S., Clyde, Z., Coffey, E., & Johnstone, E.C., (2001). A randomized, controlled trial of a brief interventional package for schizophrenic outpatients. *Acta Psychiatr Scand*, 103, 362-369.
- Chiliza, B., Asmal, L., & Emsley, R. (2012). Early intervention in schizophrenia in developing countries: focus on duration of untreated psychosis and remission as a treatment goal. *Int Rev Psychiatry*, 24(5), 483-8.
- Chung, Y.S., Kang, D., Shin, N.Y., Yoo, S.Y., & Kwon, J.S. (2008). Deficit of theory of mind in individuals at ultra-high-risk for schizophrenia. *Schizophren Res*, 99, 111-118.
- Comparelli, A., Corigliano, V., De Carolis, A., Mancinelli, I., Trovini, G., Ottavi, G., Dehning, J., Tatarelli, R., Brugnoli, R., & Girardi, P. (2013). Emotion recognition impairment is present early and is stable throughout the course of schizophrenia. *Schizophr Res*, 143(1), 65-9.
- Corcoran, R., Cahill, C., & Frith, C.D. (1997). The appreciation of visual jokes in people with schizophrenia: a study of mentalizing ability. *Schizophr Res*, 24(3), 319-327.
- Corcoran, C.M., Keilp, J.G., Kayser, J., Klim, C., Butler, P.D., Bruder, G.E., Gur, R.C., & Javitt, D.C. (2015). Emotion recognition deficits as predictors of transition in individuals at clinical high risk for schizophrenia: a neurodevelopmental perspective. *Psychol Med*, 4, 1-15.
- Corigliano, V., De Carolis, A., Trovini, G., Dehning, J., Di Pietro, S., Curto, M., Donato, N., De Pisa, E., Girardi, P., & Comparelli, A. (2014).

- Neurocognition in schizophrenia: from prodrome to multi-episode illness. *Psychiatry Res.* 220(1-2), 129-34.
- Couture, S.M., Penn, D.L., Addington, J., Woods, S.W., & Perkins, D.O. (2008). Assessment of social judgments and complex mental states in the early phases of psychosis. *Schizophr Res.* 100(1-3), 237-41.
- Dam, J. (2006). Insight in schizophrenia: a review. *Nord. J. Psychiatry*, 60, 114-120.
- David, A.S. 1990. Insight and psychosis. *Br. J. Psychiatry*, 156, 789-809.
- David, A.S. (2004). The clinical importance of insight: an overview. In: Amador, X.F., & David, A.S. (Eds.), *Insight and Psychosis*. Oxford University Press, Oxford.
- David, A.S., Ster, I.C., & Zavarei, H. (2012). Effect of video self versus other observation on insight in psychotic disorders. *J Nerv Ment Dis.* 200(4), 358-361.
- Davidoff, S.A., Forester, B.P., Chaemi, S.N., & Bodkin, J.A. (1998). The effect of video self-observation on development of insight in psychotic disorders. *J Nerv Ment Dis.* 186(11), 697-700.
- Davis, M.H. (1983). Measuring individual differences in empathy: Evidence for a multidimensional approach. *Journal of personality and social psychology*, 44(1), 113.
- Davis, M.H. (1980). A multidimensional approach to individual differences in empathy. *JSAS Catalog of Selected Documents in Psychol.* 10, 85.
- Davis, L.W., Eicher, A.C., & Lysaker P.H. (2011). Metacognition as a predictor of therapeutic alliance over 26 weeks of psychotherapy in schizophrenia. *Schizophr Res.* 129, 85-90.
- de Jong, S., van Donkersgoed, R.J.M., Aleman, A., van der Gaag, M., Wunderink, L., Arends, J., Lysaker, P.H., & Pijnenborg, M. (2016). Practical Implications of Metacognitively Oriented Psychotherapy in Psychosis. *The Journal of Nervous and Mental Disease*, 204(9), 713-716.
- De Jong, S., van Donkersgoed, R., Pijnenborg, G.H.M., & Lysaker, P.H. (2016). Metacognitive reflection and insight therapy (MERIT) with a patient with severe symptoms of disorganization. *Journal of Clinical Psychology*, 72, 164-174.
- DeVylder, J.E., Ben-David, S., Kimhy, D., & Corcoran, C.M. (2013). Attributional style among youth at clinical risk for psychosis. *Early Interv Psychiatry*, 7(1), 84-8.
- Devlin, H.C., Zaki, J., Ong, D.C., & Gruber, J. (2014). Not as good as you think? Trait positive emotion is associated with increased self-reported empathy but decreased empathic performance. *Plos One*, 9(10), e110470.
- Dimaggio, G., & Lysaker P.H. *Metacognition and severe adult mental disorders: From basic research to treatment*. London: Bruner Routledge; 2010.
- Dimaggio, G., & Lysaker, P.H. (2015). Metacognition and mentalizing in the psychotherapy of patients with psychosis and personality disorders. *Journal of Clinical Psychology: In Session*, 71(2), 117-124.
- Dimaggio, G., Lysaker, P.H., Carcione, A., Nicolò, G., & Semerari, A. (2008). Know yourself and you shall know the other... to a certain extent: Multiple paths of influence of self-reflection on mindreading. *Conscious and Cognition*, 17, 778-789.
- Dimaggio, G., Semerari, A., Carcione, A., Nicolò, G., & Procacci, M. (2007). *Psychotherapy of Personality Disorders*. London: Bruner Routledge.
- Drake, R.J., Dunn, G., Tarrier, N., Bentall, R.P., Haddock, G., & Lewis, S.W. (2007). Insight as a predictor of the outcome of first-episode nonaffective psychosis in a prospective cohort study in England. *J Clin Psychiatry*, 68, 81-86.
- Drury, V., Birchwood, M., & Cochrane, R. (2000). Cognitive therapy and recovery from acute psychosis: a controlled trial. 3. Five-year follow-up. *Br J Psychiatry*, 177, 8-14.
- Eack, S.M., Mermon, D.E., Montrose, D.M., Miewald, J., Gur, R.E., Gur, R.C., Sweeney, J.A., & Keshavan, M.S. (2010). Social cognition deficits among individuals at familial high risk for schizophrenia. *Schizophr Bull.* 36(6), 1081-8.
- Edwards, J., Pattison, P.E., Jackson, H.J., & Wales, R.J. (2001). Facial affect and affective prosody recognition in first-episode schizophrenia. *Schizophr Res.* 48(2-3), 235-53.
- Egger, M., Smith, G.D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *Brit Med J.* 315, 639-34.

REFERENCES

- Eisenberg, N., & Miller, P.A. (1987). The relation of empathy to prosocial and related behaviors. *Psychol Bull*, 101(1), 91-119.
- Ekman, P., & Friesen, W.V. (1976). *Pictures of Facial Affect*. Palo Alto, CA: Consulting Psychologists Press.
- Elowe, J., & Conus, P. (2016). Much ado about everything: A literature review of insight in first episode psychosis and schizophrenia. *Eur Psychiatry*, 39, 73-79.
- Elliott, R., Bohart, A.C., Watson, J.C., & Greenberg, L.S. (2011). Empathy. *Psychotherapy*, 48, 43-49.
- Emmerson, L.C., Granholm, E., Link, P.C., McQuaid, J.R., & Jeste, D.V. (2009). Insight and treatment outcome with cognitive behavioral social skills training for older people with schizophrenia. *J Rehabil Res Dev*, 46(8), 1053-1058.
- Fava, G.A., & Kellner, R. (1993). Staging: a neglected dimension in psychiatric classification. *Acta Psychiatr Scand*, 87, 225-30.
- Favrod, J., Maire, A., Bardy, S., Pernier, S., & Bonsack, C. (2011). Improving insight into delusions: a pilot study of metacognitive training for patients with schizophrenia. *J Adv Nurs*, 67(2), 401-407.
- Feingold, D., Hasson-Ohayon, I., Laukka, P., Vishne, T., Dembinsky, Y., & Kravets, S. (2016). Emotion recognition deficits among persons with schizophrenia: beyond stimulus complexity level and presentation modality. *Psychiatry Research*, 240, 60-65.
- Ferguson, C.J. (2009). An effect size primer: a guide for clinicians and researchers. *Professional Psychol: Res and Pract*, 40(5), 535-538.
- Flavell, J.H. (1979). Metacognition and cognitive monitoring: a new area of cognitive developmental inquiry. *Am Psychol*, 34(10), 906-911.
- Fonagy, P., Gergely, G., Jurist, E.L., & Target, M. (2002). *Affect Regulation, Mentalization and the Development of the Self*. New York: Other Press.
- Fowler, D., Hodgekins, J., Painter, M., Reilly, T., Crane, C., Macmillan, I., Mugford, M., Croudace, T., & Jones, P.B. (2009). Cognitive Behaviour therapy for improving social recovery in psychosis: a report from the ISREP MRC Trial Platform study (Improving Social Recovery in Early Psychosis). *Psychol Med*, 39, 1-10.
- Frith, C.D. (1992). *The cognitive neuropsychology of schizophrenia*. Sussex, England: Lawrence Erlbaum Associates.
- Frith, C.D., & Frith, U. (2008). Implicit and explicit processes in social cognition. *Neuron*, 60(3), 503-10.
- Fung, K.M.T., Tsang, H.W.H., & Cheung, W. (2011). Randomized controlled trial of the self-stigma reduction program amongst individuals with schizophrenia. *Psychiatry Res*, 189, 208-214.
- Fusar-Poli, P., Bonoldi, I., Yung, A.R., Borgwardt, S., Kempton, M.J., Valmaggia, L., Barale, F., Caverzasi, E., & McGuire, P. (2012). Predicting psychosis: meta-analysis of transition outcomes in individuals at high clinical risk. *Arch Gen Psychiatry*, 69(3), 220-9.
- Fusar-Poli, P., & Borgwardt, S. (2012). Predictive power of attenuated psychosis syndrome: is it really low? The case of mild cognitive impairment. *Schizophr Res*, 135(1-3), 192-3.
- Fusar-Poli, P., Deste, G., Smieskova, R., Barlati, S., Yung, A.R., Howes, O., Stieglitz, R.D., Vita, A., McGuire, P., & Borgwardt, S. (2012). Cognitive functioning in prodromal psychosis: a meta-analysis. *Arch Gen Psychiatry*, 69(6), 562-71.
- Fusar-Poli, P., Papanastasiou, E., Stahl, D., Rocchetti, M., Carpenter, W., Shergill, S., & McGuire, P. (2015). Treatments of negative symptoms in schizophrenia: Meta-analysis of 168 randomized placebo-controlled trials. *Schizophrenia Bulletin*, 41(4), 892-899.
- Garety, P., Fowler, D., Kuipers, E., Freeman, D., Dunn, G., Bebbington, P., Hadley, C., & Jones, S. (1997). London-East Anglia randomized controlled trial for psychosis. II: Predictors of outcome. *Br J Psychiatry*, 171, 420-426.
- Gez, D.C., & Cannon, T.D. (2011). Prediction of conversion to psychosis: review and future directions. *Rev Bras Psiquiatr*, 33(2), s129-s142.
- Gez, D.C., Karlsgodt, K.H., Van Erp, T.G., Bearden, C.E., Lieberman, M.D., Belger, A., Perkins, D.O., Olvet, D.M., Cornblatt, B.A., Constable, T., Woods, S.W., Addington, J., Cadenhead, K.S., McGlashan, T.H., Seidman, L.J., Tsuang, M.T., Walker, E.F., Cannon, T.D. & NAPLS Consortium (2012). Altered age-related

- trajectories of amygdala-prefrontal circuitry in adolescents at clinical high risk for psychosis: a preliminary study. *Schizophr Res*, 134(1), 1-9.
- Gesn, P.R., & Ickes, W. (1999). The development of meaning contexts for empathic accuracy: channels and sequence effects. *Journal of Personality and Social Psychology*, 77, 746-761.
- Gill, K.E., Cressman, V., Poe, S.L., Steinfeld, S., Ben-David, S., Keilp, J.G., Moore, H., Turkstra, L.S., & Corcoran, C. (2014). Social inference in individuals at clinical high risk for psychosis. *Early Intervention in Psychiatry*, 10(1), 77-80.
- Giuliano, A.J., Li, H., Mesholam-Gately, R.I., Sorenson, S.M., Woodberry, K.A., & Seidman, L.J. (2012). Neurocognition in the psychosis risk syndrome: a quantitative and qualitative review. *Curr Pharm Des*, 18(4), 399-415.
- Gould, R.A., Mueser, K.T., Bolton, E., Mays, V., & Goff, D. (2001). Cognitive therapy for psychosis in schizophrenia: An effect size analysis. *Schizophrenia Research*, 48(2-3), 335-342.
- Granholm, E., McQuaid, J.R., McClure, F.S., Auslander, L.A., Perivoliotis, D., Pedrelli, P., Patterson, T., & Jeste, D.V. (2005). A randomised controlled trial of cognitive behavioral social skills training for middle-aged and older outpatients with chronic schizophrenia. *Am. J. Psychiatry*, 162, 520-529.
- Granholm, E., McQuaid, J.R., McClure, F.S., Link, P.C., Perivoliotis, D., Gottlieb, J.D., Patterson, T.L., & Jeste, D.V. (2007). Randomised controlled trial of cognitive behavioral social skills training for middle-aged and older outpatients with chronic schizophrenia: 12-month follow-up. *J. Clin. Psychiatry*, 68, 730-737.
- Gray, R., Leese, M., Bindman, J., Becker, T., Burti, L., David, A., Gournay, K., Kikkert, M., Koeter, M., Puschner, B., Schene, A., Thornicroft, G., & Tanselle, M. (2006). Adherence therapy for people with schizophrenia. *Br. J. Psychiatry*, 189, 508-514.
- Green, M.F., Bearden, C.E., Cannon, T.D., Fiske, A.P., Helleman, G.S., Horan, W.P., Kez, K., Kern, R.S., Lee, J., Sergi, M.J., Subotnik, K.L., Sugar, C.A., Ventura, J., Yee, C.M., & Nuechterlein, K.H. (2012). Social cognition in schizophrenia, Part 1: performance across phase of illness. *Schizophr Bull*, 38(4), 854-64.
- Gross, G. (1989). The 'basic' symptoms of schizophrenia. *Br J Psychiatry*, 155(7), 21-25.
- Green, M.F., & Harvey, P.D. (2014). Cognition in schizophrenia: Past, present, and future. *Schizophren Res Cogn*, 1(1), e1-e9.
- Gross, J.J., John, O.P., & Richards, J. (1995). *Berkeley expressivity questionnaire*. Edwin Mellen Press.
- Guastella, A.J., Hermens, D.F., Van Zwieten, A., Naismith, S.L., Lee, R.S., Cacciotti-Sajja, C., Scott, E.M., & Hickie, I.B. (2013). Social cognitive performance as a marker of positive psychotic symptoms in young people seeking help for mental health problems. *Schizophr Res*, 149(1-3), 77-82.
- Guo, X., Zhai, J., Zhening, L., Maosheng, F., Wang, B., Wang, C., Bin, H., Sun, X., Luxian, L., Zheng, L., Cui, M., He, X., Guo, T., Xie, S., Wu, R., Xue, Z., Chen, J., Tuamley, E.W., Jin, H., & Zhao, J. (2010). Effect of antipsychotic medication alone vs combined with psychosocial intervention on outcomes of early-stage schizophrenia. *Arch. Gen. Psychiatry*, 67(9), 895-904.
- Haddock, G., Lewis, S., Bentall, R., Dunn, G., Drake, R., & Tarrier, N. (2006). Influence of age on outcome of psychological treatments in first-episode psychosis. *Br. J. Psychiatry*, 188, 250-254.
- Hamm, J.A., Renard, S.B., Fogley, R.L., Leonhardt, B.L., Dimaggio, G., Buck, K.D., & Lysaker, P.H. (2012). Metacognition and social cognition in schizophrenia: stability and relationship to concurrent and prospective symptom assessments. *Journal of Clinical Psychology*, 68(12), 1303-12.
- Happé, F.G. (1994). An advanced test of theory of mind: understanding of story characters' thoughts and feelings by able autistic, mentally handicapped, and normal children and adults. *J Autism Dev Disord*, 24, 129-154.
- Hasson-Ohayon, I., Kravetz, S., & Lysaker, P.H. (2017). The special challenges of psychotherapy with persons with psychosis: intersubjective metacognitive model of agreement and shared meaning. *Clin Psychol Psychother*, 24(2), 428-440.
- Harvey, P.O., Zaki, J., Lee, J., Ochsner, K., & Green, M.F. (2013). Neural substrates of empathic accuracy in people with schizophrenia. *Schizophr Bull*, 39(3), 617-28.
- Haro, J.M., Kamath, S.A., Ochoa, S., Novick, D., Rele, K., Fargas, A., Rodriguez, M.J., Rele, R.,

REFERENCES

- Orta, J., Kharbeng, A., Araya, S., Gervin, M., Alonso, J., Mavreas, V., Lavrentzou, E., Liotos, N., Gregor, K., Jones, P.B., & SOHO Study Group. (2003). The Clinical Global Impression-Schizophrenia scale: a simple instrument to measure the diversity of symptoms present in schizophrenia. *Acta Psychiatrica Scandinavica, Suppl 107*(416), 16-23.
- Hayward, P., David, A.S., Green, N., Rabe-Hesketh, S., Haworth, E., Thompson, N., Shaffer, S., Hastilow, S., & Wykes, T. (2009). Promoting therapeutic alliance in clozapine users: an exploratory randomized controlled trial. *Clin. Schizophr. Rel. Psychoses, 3*, 127-132.
- Healey, K.M., Penn, D.L., Perkins, D., Woods, S.W., & Addington, J. (2013). Theory of mind and social judgements in people at clinical high risk of psychosis. *Schizophren Res, 150*, 498-504.
- Hendryx, M., Dyck, D.G., McBride, D., & Whitbeck, J. (2001). A test of the reliability and validity of the Multnomah Community Ability Scale. *Community Mental Health Journal, 37*(2), 157-168.
- Henry, C.S., & Ghaemi, N. (2004). Insight in psychosis: a systematic review of treatment interventions. *Bipolar Disord. 37*, 194-199.
- Hillis, J.D., Leonhardt, B.L., Vohs, J.L., Buck, K.D., Salvatore, G., Popolo, R., Dimaggio, C., & Lysaker, P.H. (2015). Metacognitive reflective and insight therapy for people in early phase of a schizophrenia spectrum disorder. *Journal of Clinical Psychology, 71*(2), 125-135.
- Hoekert, M., Kahn, R.S., Pijnenborg, M., & Aleman, A. (2007). Impaired recognition and expression of emotional prosody in schizophrenia: review and meta-analysis. *Schizophr Res, 96*(1-3): 135-45.
- Horan, W.P., Reise, S.P., Kern, R.S., Lee, J., Penn, D.L., & Green, M.F. (2015). Structure and correlates of self-reported empathy in schizophrenia. *J Psychiatr Res, 66-67*, 60-6.
- Houbova, M., Prasko, J., Matousek, S., Latalova, K., Marackova, M., Vrbova, K., Grambal, A., Slepceky, M., & Zatkova, M. (2016). Comparison of self-stigma and quality of life in patients with depressive disorders and schizophrenia spectrum disorders - a cross-sectional study. *Neuropsychiatr Dis Treat, 12*, 3021-3030.
- Hunter, R., & Barry, S. (2012). Negative symptoms and psychosocial functioning in schizophrenia: Neglected but important targets for treatment. *European Psychiatry, 27*(6), 432-436.
- Hur, J.W., Byun, M.S., Shin, N.Y., Shin, Y.S., Kim, S.N., Jang, J.H., & Kwon, J.S. (2013). General intellectual functioning as a buffer against theory-of-mind deficits in individuals at ultra-high risk for psychosis. *Schizophren Res, 149*(1-3), 83-7.
- Ickes, W., Stinson, L., Bissonnette, V., & Garcia, S. (1990). Naturalistic social cognition: empathic accuracy in mixed-sex dyads. *Journal of Personality and Social Psychology, 59*, 730-742.
- Ihnen, G.H., Penn, D.L., Corrigan, P.W., & Martin J. (1998). Social perception and social skill in schizophrenia. *Psychiatry Res, 80*(3), 275-86.
- Instituut, T. (2012). MULTIDISCIPLINAIRE RICHTLIJN SCHIZOFRENIE 2012.
- Ising, H.K., Smit, F., Veling, W., Rietdijk, J., Dragt, S., Klaassen, R.M., Savelsberg, N.S., Boonstra, N., Nieman, D.H., Linszen, D.H., Wunderink, L., & Van der Gaag, M. (2014). Cost-effectiveness of preventing first-episode psychosis in ultra-high-risk subjects: multi-centre randomized controlled trial. *Psychol Med, 21*, 1-12.
- Jong-Hoon, K., Jun-Hung, A., & Lee, J. (2011). Insight change and its relationship to subjective well-being during acute atypical antipsychotic treatment in schizophrenia. *J. Clin. Pharm. Ther, 36*, 687-694.
- Kay, S.R., Fiszbein, A., & Opler, L.A. (1987). The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull, 13*(2), 261 - 276.
- Kazdin, A.E. (2010). *Research Design in Clinical Psychology (Fourth)*. Boston: Allyn & Bacon.
- Kazdin, A.E., & Bass, D. (1989). Power to detect differences between alternative treatments in comparative psychotherapy outcome research. *J Consult Clin Psychol, 57*(1), 138-147.
- Kemp, R., & David, A.T. (1996). Insight and compliance. In: Blackwell B. *Treatment Compliance and the Therapeutic Alliance*. Gordon & Breach, Newark.
- Kemp, R., Kirov, G., Everitt, B., Hayward, P., & David, A. (1998). A randomised controlled trial of compliance therapy: 18-month follow-up. *Br. J. Psychiatry, 172*, 413-419.

- Kerr, S.L., & Neale, J.M. (1993). Emotion perception in schizophrenia: specific deficit or further evidence of generalized poor performance. *Journal of Abnormal Psychology*, 102(2), 312-318.
- Keyser, C., & Gazzola, V. (2006). Towards a unifying neural theory of social cognition. *Prog Brain Res*, 156, 379-401.
- Kim, H.S., Shin, N.Y., Choi, J.S., Jung, M.H., Jang, J.H., Kang, D.H., & Kwon, J.S. (2010). Processing of facial configuration in individuals at ultra-high risk for schizophrenia. *Schizophrenia Res*, 118(1-3), 81-7.
- Kim, H.S., Shin, N.Y., Jang, J.H., Kim, E., Shim, G., Park, H.Y., Hong, K.S., & Kwon, J.S. (2011). Social cognition and neurocognition as predictors of conversion to psychosis in individuals at ultra-high risk. *Schizophrenia Res*, 130(1-3), 170-5.
- Kinderman, P., & Bentall, R.P. (1996). A new measure of causal locus: the internal, personal and situational attributions questionnaire. *Personality and Individual Differences*, 20(2), 261-4.
- Klosterkötter, J., Schultze-Lutter, F., Gross, G., Huber, G., & Steinmeyer, E.M. (1997). Early self-experienced neuropsychological deficits and subsequent schizophrenic diseases: an 8-year average follow-up prospective study. *Acta Psychiatrica Scand*, 95(5), 396-404.
- Kohler, C.G., Richard, J.A., Brensinger, C.M., Borgmann-Winter, K.E., Conroy, C.G., Moberg, P.J., Gur, R.C., Gur, R.E., & Colkins, M.E. (2014). Facial emotion perception differs in young persons at genetic and clinical high-risk for psychosis. *Psychiatry Res*, 216(2), 206-12.
- Kohler, C.G., Walker, J.B., Martin, E.A., Healey, K.M., & Moberg, P.J. (2010). Facial emotion perception in schizophrenia: a meta-analytic review. *Schizophrenia Bull*, 36(5), 1009-19.
- Kraepelin, E. (1919). *Dementia praecox and paraphrenia*. New York: Barclay RM.
- Kukla, M., Lysaker, P.H., & Salyers, M.P. (2013). Do persons with schizophrenia who have better metacognitive capacity also have a stronger subjective experience of recovery? *Psychiatry Res*, 209(3), 381-385.
- Kukla, M., Whitesel, F., & Lysaker, P.H. (2015). An integrative psychotherapy approach to foster community engagement and rehabilitation in schizophrenia: A case study illustration. *Journal of Clinical Psychology: In Session*, 72(2), 152-163.
- Kundu, P.S., Sinha, V.K., Paul, S.E., & Desarkar, P. (2013). Current social functioning in adult-onset schizophrenia and its relation with positive symptoms. *Ind Psychiatry*, 22(1), 65-8.
- Kurtz, M.M., Moberg, J.P., Ragland, J.D., Gur, R.C., & Gur R.E. (2005). Symptoms versus neurocognitive test performance as predictors of psychosocial status in schizophrenia: a 1- and 4-year prospective study. *Schizophrenia Bulletin*, 31, 167-174.
- Langdon, R., Ward, P.B., & Coltheart, M. (2010). Reasoning anomalies associated with delusions in schizophrenia. *Schizophrenia Bull*, 36(2), 321-30.
- Lavoie, M.A., Plana, I., Bédard Lacroix, J., Godmaire-Duhaine, F., Jackson, P.L., & Achim, A.M. (2013). Social cognition in first-degree relatives of people with schizophrenia: a meta-analysis. *Psychiatry Res*, 209(2), 129-35.
- Lawrence, K., Campbell, R., & Skuse, D. (2015). Age, gender and puberty influence the development of facial emotion recognition. *Front Psychol*, 6, 761.
- Lee, S.Y., Bang, M., Kim, K.R., Lee, M.K., Park, J.Y., Song, Y.Y., Kang, J.I., Lee, E., & An S.K. (2015). Impaired facial emotion recognition in individuals at ultra-high risk for psychosis and with first-episode schizophrenia and their associations with neurocognitive deficits and self-reported schizotypy. *Schizophrenia Res*, 165(1), 60-5.
- Lee, T.Y., Hong, S.B., Shin, N.Y., & Kwon, J.S. (2015). Social cognitive functioning in prodromal psychosis: a meta-analysis. *Schizophrenia Res*, 164(1-3), 28-34.
- Lee, D.A., Randall, F., Beattie, G., & Bentall, R.P. (2004). Delusional discourse: an investigation comparing the spontaneous causal attributions of paranoid and non-paranoid individuals. *Psychol Psychother*, 77(4), 525-40.
- Lee, J., Zaki, J., Harvey, P.O., Ochsner, K., & Green, M.F. (2011). Schizophrenia patients are impaired in empathic accuracy. *Psychol Med*, 41(11), 2297-304.
- Leitman, D.I., Foxe, J.J., Butler, P.D., Saperstein, A., Revheim, N., & Javitt, D.C. (2005). Sensory contributions to impaired prosodic processing in schizophrenia. *Biol Psychiatry*, 58(1), 56-61.

REFERENCES

- Levenson, R.W., & Ruef, A.M. (1992). Empathy: a physiological substrate. *J Pers Soc Psychol*, 63(2), 234-46.
- Lieburg, E.J., Kneftering, H., Klien, H.C., Kortekaas, R., & Aleman, A. (2012). Antipsychotic medication and prefrontal cortex activation: a review of neuroimaging findings. *Eur. Neuropsychopharmacol*, 22(6), 387-400.
- Lincoln, T.M., Lullmann, E., & Rief, W. (2007). Correlates and long-term consequences of poor insight in patients with schizophrenia: A systematic review. *Schizophr. Bull*, 33, 1324-1342.
- Lipsey, M.W., & Wilson, D.B. (2000). *Practical Meta-Analysis*. SAGE publications, Thousand Oaks, California.
- Luther, L., Firmin, R.L., Vohs, J.L., Buck, K.D., Rand, K.L., & Lysaker P.H. (2016). Intrinsic motivation as a mediator between metacognition deficits and impaired functioning in psychosis. *British Journal of Clinical Psychology*, 55(3), 332-47.
- Lysaker, P.H., Buck, K.D., Carcione, A., Procacci, M., Salvatore, G., Nicola, G., & Dimaggio, G. (2010). Addressing metacognitive capacity for self-reflection in the psychotherapy for schizophrenia: a conceptual model of the key tasks and processes. *Psychol Psychother*, 84(1), 58-69.
- Lysaker, P.H., Buck, K.D., & Ringer, J. (2007). The recovery of metacognitive capacity in schizophrenia across 32 months of individual psychotherapy: a case study. *Psychotherapy Res*, 17(6), 713-720.
- Lysaker, P.H., Bob, P., Ondrej, P., Hamm, J., Kukula, M., Vohs, J., Popolo, R., Salvatore, G., & Dimaggio, G. (2013). Synthetic metacognition, as a link between brain and behavior in schizophrenia. *Transl Neurosci*, 4(3), 368-377.
- Lysaker, P.H., Carcione, A., Dimaggio, G., Johannesen, J.K., Nicolò, G., Procacci, M., & Semerari, A. (2005). Metacognition amidst narratives of self and illness in schizophrenia: associations with insight, neurocognition, symptom and function. *Acta Psychiatrica Scandinavica*, 112, 64-71.
- Lysaker, P.H., Clements, C.A., Plascak-Hallberg, C.D., Knipscheer, S.J., & Wright, D.E. (2002). Insight and personal narratives of illness in schizophrenia. *Psychiatry*, 65(3), 197-206.
- Lysaker, P.H., Davis, L.W., Eckert, G.J., Strasburger, A.M., Hunter, N.L., & Buck, K.D. (2005). Changes in narrative structure and content in schizophrenia in long term individual psychotherapy: a single case study. *Clin Psychol Psychother*, 12(5), 406-416.
- Lysaker, P.H., & Dimaggio, G. (2014). Metacognitive capacities for reflection in schizophrenia: Implications for developing treatments. *Schizophrenia Bulletin*, 40(3), 487-491.
- Lysaker, P.H., Dimaggio, G., Carcione, A., Procacci, M., Buck, K.D., Davis, L.W., & Nicolò, G. (2010). Metacognition and schizophrenia: the capacity for self-reflexivity as a predictor for prospective assessments of work performance over six months. *Schizophrenia Res*, 122(1-3), 124-130.
- Lysaker, P.H., Dimaggio, G., Daroyanni, P., Buck, K.D., LaRocco, V.A., Carcione, A., & Nicolò, G. (2010). Assessing metacognition in schizophrenia with the Metacognition Assessment Scale: Associations with the Social Cognition and Object Relations Scale. *Psychology and Psychotherapy*, 83, 303-15.
- Lysaker, P.H., Erickson, M., Ringer, J., Buck, K.D., Semerari, A., Carcione, A., & Dimaggio, G. (2011). Metacognition in schizophrenia: the relationship of mastery to coping, insight, self-esteem, social anxiety, and various facets of neurocognition. *Br J Clin Psychol*, 50(4), 412-424.
- Lysaker, P.H., Kukla, M., Dubreuca, J., Gumley, A., McLeod, H., Vohs, J.L., Buck, K.D., Minor, K.S., Luther, L., Leonhardt, B.L., Belanger, E.A., Popolo, R., & Dimaggio, G. (2015). Metacognitive deficits predict future levels of negative symptoms in schizophrenia controlling for neurocognition, affect recognition, and self-expectation of goal attainment. *Schizophrenia Research*, 168, 267-272.
- Lysaker, P.H., McCormick, B.P., Sneathen, G., Buck, K.D., Hamm, J.A., Grant, M., Nicolò, G., & Dimaggio, G. (2011). Metacognition and social function in schizophrenia: associations of mastery with functional skills competence. *Schizophr Res*, 131, 214-218.
- Lysaker, P.H., Shea, A.M., Buck, K.D., Dimaggio, G., Nicolò, G., Procacci, M., Salvatore, G., & Rand, K.L. (2010). Metacognition as a mediator of the effects of impairments in neurocognition on social function in schizophrenia spectrum disorders. *Acta Psychiatr Scand*, 122(5), 405-413.

- Lysaker, P.H., Vohs, J., Hamm, J.A., Kukla, M., Minor, K.S., de Jong, S., van Donkersgoed, R., Pijnenborg, M.H., Kent, J.S., Matthews, S.C., Ringer, J.M., Leonhardt, B.L., Francis, M.M., Buck, K.D., & Dimaggio, G. (2014). Deficits in metacognitive capacity distinguish patients with schizophrenia from those with prolonged medical adversity. *Journal of Psychiatric Research*, 55, 126-32.
- MacBeth A, & Gumley, A. (2008). Premorbid adjustment, symptom development and quality of life in first episode psychosis: a systematic review and critical reappraisal. *Acta Psychiatr Scand*, 117(2), 85-99.
- Macbeth, A., Gumley, A., Schwannauer, M., Carcione, A., Fisher, R., McLeod, H.J., & Dimaggio, G. (2013). Metacognition, symptoms and premorbid functioning in a first episode psychosis sample. *Compr Psychiatry*, 55(2), 268-73.
- Macpherson, R., Jerrom, B., & Hughes, A. (1996). A controlled study of education about drug treatment in schizophrenia. *Br. J. Psychiatry*, 168, 709-717.
- Mairs, H., Lovell, K., Campbell, M., & Keeley, P. (2011). Development and pilot investigation of behavioral activation for negative symptoms. *Behavior Modification*, 35(5), 486-506.
- Marengo, J. (1994). Classifying the courses of schizophrenia. *Schizophr Bull*, 20(3), 519-36.
- Marjoram, D., Job, D.E., Whalley, H.C., Gountouna, V.E., McIntosh, A.M., Simonotto, E., Cunningham-Owens, D., Johnstone, E.C., & Laurie, S. (2006). A visual joke fMRI investigation into Theory of Mind and enhanced risk of schizophrenia. *Neuroimage*, 31(4), 1850-8.
- Marshall, M., Lewis, S., Lockwood, A., Drake, R., Jones, P., & Croudace, T. (2005). Association between duration of untreated psychosis and outcome in cohorts of first-episode patients: a systematic review. *Arch Gen Psychiatry*, 62(9), 975-83.
- Mazza, M., Pollice, R., Pacitti, F., Pino, M.C., Mariano, M., Tripaldi, S., Casacchia, M., & Roncone, R. (2012). New evidence in theory of mind deficits in subjects with chronic schizophrenia and first episode: correlation with symptoms, neurocognition and social function. *Riv Psichiatr*, 47(4), 327-36.
- McDonald, S., Flanagan, S., Rollins, S., & Kinch, J. (2003). TASIT: A new clinical tool for assessing social perception after traumatic brain injury. *Journal of Head Trauma Rehabilitation*, 18(3), 219-238.
- McEvoy, J.P., Apperson, J., Appelbaum, P.S., Ortillip, P., Brecosky, J., Hammill, K., Celler, J.L., & Roth, L. (1989). Insight in schizophrenia: its relationship to acute psychopathology. *J. Nerv. Ment. Dis*, 17, 43-47.
- McEvoy, J.P., Johnson, J., Perkins, D., Lieberman, J.A., Hamer, R.M., Keefe, R.S.E., Tohen, M., Glick, I.D., & Sharma, T. (2006). Insight in first-episode psychosis. *Psychol. Med*, 36, 1385-1393.
- McGlashan, T.H., Miller, T.J., & Woods, S.W. (2001). Pre-onset detection and intervention research in schizophrenia psychoses: current estimates of benefit and risk. *Schizophr Bull*, 27(4), 563-70.
- McCorry, P.D., Purcell, R., & Hickie, I.B., Yung, A.R., Pantelis, C., & Jackson, H.J. (2007) Clinical staging: A heuristic model for psychiatry and youth mental health. *Medical Journal of Australia*, 187, S40-S42.
- McGrath, J., Saha, S., Chant, D., & Welham, J. (2008). Schizophrenia: a concise overview of incidence, prevalence, and mortality. *Epidemiologic Reviews*, 30, 67-76.
- McGurk, S.R., Moriarty, P.J., Harvey, P.D., Parrella, M., White, L., & Davis, K.L. (2000). The longitudinal relationship of clinical symptoms, cognitive functioning, and adaptive life in geriatric schizophrenia. *Schizophrenia Research*, 42(1), 47-55?
- Mehta, U.M., Thirthalli, J., Naveen Kumar, C., Keshav Kumar, J., Keshavan, M.S., & Gangadhar, B.N. (2013). Schizophrenia patients experience substantial social cognition deficits across multiple domains in remission. *Asian J Psychiatr*, 6(4), 324-9.
- Merinder, L.B., Viuff, A.G., Laugesen, H.D., Clemmensen, K., Mifselt, S., & Espensen, B. (1999). Patient and relative education in community psychiatry: a randomized controlled trial regarding its effectiveness. *Soc. Psychiatry Psychiatr. Epidemiol*, 34, 287-294.
- Michaels, T.M., Horan, W.P., Ginger, E.J., Martinovich, Z., Pinkham, A.E., & Smith, M.J. (2014). Cognitive empathy contributes to poor social functioning in schizophrenia: Evidence from a new self-report measure of cognitive and affective empathy. *Psychiatry Res.*, 220(3), 803-10.

REFERENCES

- Milev, P., Ho, B.C., Arndt, S., & Andreasen, N.C. (2005). Predictive values of neurocognition and negative symptoms on functional outcome in schizophrenia: a longitudinal first-episode study with 7-year follow-up. *American Journal of Psychiatry*, 162, 495-506.
- Minor, K.S., Marggraf, M.P., Davis, B.J., Luther, L., Vohs, J.L., Buck, K.D., & Lysaker, P.H. (2015). Conceptual disorganization weakens links in cognitive pathways: Disentangling neurocognition, social cognition, and metacognition in schizophrenia. *Schizophr Res*, 169(1-3), 153-8.
- Mintz, A.R., Dobson, K.S., & Romney, D.M. (2003). Insight in schizophrenia: a meta-analysis. *Schizophr. Res*, 61(1), 75-88.
- Möller, H.J. (2007). Clinical evaluation of negative symptoms in schizophrenia. *European Psychiatry*, 22(6), 380-386.
- Morris, S.B. (2008). Estimating effect sizes from pretest-posttest-control group designs. *Organ. Res. Methods*, 11(2), 364-386.
- Morrison, A.P., French, P., Walford, L., Lewis, S.W., Kilcommons, A., Green, J., Parker, S., & Bentall, R.P. (2004). Cognitive therapy for the prevention of psychosis in people at ultra-high risk: randomised controlled trial. *Br J Psychiatry*, 185, 291-7.
- Moritz, S., Kerstan, A., Veckenstedt, R., Randjibar, S., Vitzthum, F., Schmidt, C., & Woodward, T.S. (2011). Further evidence for the efficacy of a metacognitive group training in schizophrenia. *Behaviour Research and Therapy*, 49(3), 151-7.
- Moritz, S., & Woodward, T. (2007). Metacognitive training for schizophrenia patients (MCT): a pilot study on feasibility, treatment adherence and subjective efficacy. *German J of Psychiatry*, 10, 69-78.
- National Institute of Mental Health (NIMH) (2017). Research Domain Criteria (RDoC).
- Nasrallah, H., Morosini, P., & Gagnon, D. D. (2008). Reliability, validity and ability to detect change of the Personal and Social Performance scale in patients with stable schizophrenia. *Psychiatry Research*, 161(2), 213-24.
- Nelson, B., Yuen, H.P., Wood, S.J., Lin, A., Spiliotacopoulos, D., Bruxner, A., Broussard, C., Simmons, M., Foley, D.L., Brewer, W.J., Francey, S.M., Amminger, G.P., Thompson, A., McGorry, P.D., & Yung, A.R. (2013). Long-term follow-up of a group at ultra high risk ("prodromal") for psychosis: the PACE 400 study. *JAMA Psychiatry*, 70, 793-802.
- Nicolò, G., Dimaggio, G., Popolo, R., Carcione, A., Procacci, M., Hamm, J., Buck, K.D., Pompili, E., Buccione, I., Lagrotteria, B., & Lysaker, P.H. (2012). Associations of metacognition with symptoms, insight, and neurocognition in clinically stable outpatients with schizophrenia. *J Nerv Ment Dis*, 200(7), 644-647.
- Nieman, D.H., Ruhrmann, S., Dragt, S., Soen, F., Van Tricht, M.J., Koelman, J.H., Bour, L.J., Velthorst, E., Becker, H.E., Weiser, M., Linszen, D.H., & de Haan, L. (2014). Psychosis prediction: stratification of risk estimation with information-processing and premorbid functioning variables. *Schizophr Bull*, 40(6), 1482-90.
- O'Donnell, C., Donohoe, G., Sharkey, L., Owens, N., Migone, M., Harries, R., Kinsella, A., Larkin, C., & O'Callaghan, E. (2003). Compliance therapy: a randomised controlled trial in schizophrenia. *Br. Med. J*, 327, 834-836.
- Ofir-Eyal, S., Hasson-Ohayon, & I., Kravetz, S. (2014). Affective and cognitive empathy and social quality of life in schizophrenia: A comparison between a parallel process model and an integrative meditation model. *Psychiatry Research*, 220(1-2), 51-7.
- Ohmuro, N., Matsumoto, K., Katsura, M., Sakuma, A., Izuka, K., & Kikuchi, T. (2012). Association of deficits in theory of mind and functioning in at-risk mental states and first-episode psychosis. IEPA conference, San Francisco.
- Ostrom, J. (1984). The sovereignty of social cognition. In: Wyer R.S. and Srull T. K., *Handbook of Social Cognition*, vol. 1. Hillsdale: Erlbaum, pp. 1-38.
- Owen, M.J., Sawa, A., & Mortensen, P.B. (2016). Schizophrenia. *Lancet*, 388(10039), 86-97.
- Patterson, T.L., Moscona, S., McKibbin, C.L., Davidson, K., & Jeste, D.V. (2001). Social skills performance assessment among older patients with schizophrenia. *Schizophr Res*, 48(2-3), 351-60.
- Penn, D.L., Meyer, P.S., Evans, E., Wirth, R.J., Cai, K., & Burchinal, M. (2009). Randomized controlled trial of group cognitive-behavioral therapy vs. enhanced supportive therapy for auditory hallucinations. *Schizophr. Res*, 109, 52-57.

- Perkins, D.O., Gu, H., Boteva, K., & Lieberman, J.A. (2005). Relationship between duration of untreated psychosis and outcome in first-episode schizophrenia: a critical review and meta-analysis. *Am J Psychiatry*, 162(10), 1785-804.
- Peters, E., Landau, S., McCrone, P., Cooke, M., Fisher, P., Steel, C., Evans, R., Carswell, K., Dawson, K., Williams, S., Howard, A., & Kuipers, E. (2010). A randomised controlled trial of cognitive behaviour therapy for psychosis in a routine clinical service. *Acta Psychiatr. Scand*, 122, 302-318.
- Petkova, E., Lu, F., Kantrowitz, J., Sanchez, J.L., Lehrfeld, J., Scaramello, N., Silipo, G., DiCostanza, J., Ross, M., Su, Z., Javitt, D.C., & Butler, P.D. (2014). Auditory tasks for assessment of sensory function and affective prosody in schizophrenia. *Compr Psychiatry*, 55(8), 1862-74.
- Picco, L., Pang, S., Lau, Y.W., Jeyagurunathan, A., Satghare, P., Abidin, E., Vaingankar, J.A., Lim, S., Poh, C.L., Chong, S.A., & Subramaniam, M. (2016). Internalized stigma among psychiatric outpatients: Associations with quality of life, functioning, hope and self-esteem. *Psychiatry Res*, 246, 500-506.
- Pickard, A.S., Kohlmann, T., & Janssen, M.F. (2007). Evaluating equivalency between response systems application of the Rasch model to a 3-level EQ-5D. *Med Care*, 45, 812-819.
- Pijnenborg, G.H., Timmerman, M.E., Derks, E.M., Fleischacker, W.W., Kahn, R.S., & Aleman, A. (2015). Differential effects of antipsychotic drugs on insight in first episode schizophrenia: Data from the European First-Episode Schizophrenia Trial (EUFEST). *Eur Neuropsychopharmacol*, 25(6), 808-16.
- Pijnenborg, G.H.M., Vander Gaag, M., Bockting, C., & Aleman, A. (2011). REFLEX: a social-cognitive group treatment to improve insight in schizophrenia: study protocol of a multi-centre RCT. *BMC Psychiatry*, 11, 161.
- Pinkham, A.E., Penn, D.L., Green, M.F., Buck, B., Healy, K., & Harvey, P.D. (2014). The Social Cognition Psychometric Evaluation Study: Results of the Expert Survey and RAND Panel. *Schizophren Bull*, 40(40), 813-823.
- Pinkham, A.E., Penn, D.L., Perkins, D.O., Graham, K.A., & Siegel, M. (2007). Emotion perception and social skill over the course of psychosis: a comparison of individuals "at-risk" for psychosis and individuals with early and chronic schizophrenia spectrum illness. *Cogn Neuropsychiatry*, 12(3), 198-212.
- Potkin, S., Bugarski-Kirola, D., Edgar, C., & Luo, S. (2012). PRM30 evaluating readiness for work in patients with schizophrenia: "The Readiness for Work Questionnaire" (WoRQ) Value Health, 15(7), A650.
- Premkumar, P., Peters, E.R., Fannon, D., Anilkumar, A.P., Kuipers, E., & Kumari, V. (2011). Coping styles predict responsiveness to cognitive behaviour therapy in psychosis. *Psychiatry Res*, 187(3), 354-362.
- Priebe, S., Huxley, P., Knight, S., & Evans, S. (1999). Application and results of the manchester short assessment of quality of life (MANSA) *Int J Soc Psychiatry*, 45(1), 7-12.
- Rabash, J., Browne, W., Goldstein, H., Yang, M., Plewis, I., Healy, M., Woodhouse, G., Draper, D., Langford, I., & Lewis, T. (2000). A user's guide to MLwiN. Centre for Multilevel Modelling: Institute of Education University of London.
- Rathod, S., Kingdon, D., Smith, P., & Turkington, D. (2005). Insight into schizophrenia: the effects of cognitive behavioural therapy on the components of insight and association with sociodemographics-data on a previously published randomised controlled trial. *Schizophr. Res*, 74, 211-219.
- Reitan, R.M., & Wolfson, D. (1985). *The Halstead-Reitan Neuropsychological Test Battery: Therapy and clinical interpretation*. Tucson, Neuropsychological Press.
- Reniers, R.L., Corcoran, R., Drake, R., Shryane, N.M., & Völlm, B.A. (2011). The QCAE: a Questionnaire of Cognitive and Affective Empathy. *J Pers Assess*, 93(1), 84-95.
- Riggs, S.E., Grant, P.M., Perivoliotis, D., & Beck, A.T. (2012). Assessment of cognitive insight: A qualitative review. *Schizophrenia Bulletin*, 38(2), 338-350.
- Ritsher, J.B., Otilingam, P.C., & Grajales, M. (2003). Internalized stigma of mental illness: psychometric properties of a new measure. *Psychiatry Res*, 121(1), 31-49.
- Robertson, B.R., Prestia, D., Twamley, E.W., Patterson, T.L., Bowie, C.R., & Harvey, P.D. (2014). Social competence versus negative symptoms as predictors of real world social functioning in schizophrenia. *Schizophr Res*, 160(1-3), 136-41.

REFERENCES

- Ruhrmann, S., Schultze-Lutter, F., & Klosterkötter, J. (2003). Early detection and intervention in the initial prodromal phase of schizophrenia. *Pharmacopsychiatry*, 36(3), 162-7.
- Ruhrmann, S., Schultze-Lutter, F., Salokangas, R.K., Heinimaa, M., Linszen, D., Dingemans, P., Birchwood, M., Patterson, P., Juckel, G., Heinz, A., Morrison, A., Lewis, S., von Reventlow, H.G., & Klosterkötter, J. (2010). Prediction of psychosis in adolescents and young adults at high risk: results from the prospective European prediction of psychosis study. *Arch Gen Psychiatry*, 67(3), 241-51.
- Rush, A.J., Trivedi, M.H., Ibrahim, H.M., Carmody, T.J., Arnow, B., Klein, D.N., Markowitz, J.C., Ninan, P.T., Kornstein, S., Manber, R., Thase, M.E., Kocsis, J.H., & Keller, M.B. (2003). The 16-item quick inventory of depressive symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): a psychometric evaluation in patients with chronic major depression. *Biol Psychiatry*, 54(5), 573-583.
- Savva, G.N., Vella, L., Armstrong, C.C., Penn, D.L., & Twamley, E.W. (2013). Deficits in domains of social cognition in schizophrenia: a meta-analysis of the empirical evidence. *Schizophr Bull*, 39(5), 979-92.
- Salvatore, G., Lysaker, P.H., Gumley, A., Popolo, R., Mari, J., & Dimaggio, G. (2012). Out of illness experience: metacognition-oriented therapy for promoting self-awareness in individuals with psychosis. *Am J Psychother*, 66(1), 85-106.
- Salvatore, G., Procacci, M., & Popolo, R. (2009). Adapted metacognitive interpersonal therapy for improving adherence to intersubjective contexts in a person with schizophrenia. *Clinical Case Studies*, 8, 473-488.
- Salvatore, G., Russo, B., Russo, M., Popolo, R., & Dimaggio, G. (2012). Metacognition-Oriented Therapy for Psychosis: The Case of a Woman with Delusional Disorder and Paranoid Personality Disorder. *Journal of Psychotherapy Integration*, 22(4), 314-329.
- Schenkel, L.S., Spaulding, W.D., & Silverstein, S.M. (2005). Poor premorbid social functioning and theory of mind deficit in schizophrenia: evidence of reduced context processing? *J Psychiatr Res*, 39(5), 499-508.
- Schmand, B.A., Bakker, D., Saan, R.J., & Louman, J. (1991). De Nederlandse Leestest voor Volwassenen: een maat voor het premorbid intelligentieniveau. The Dutch Adult Reading Test: a measure of premorbid intelligence. *Tijdschr Gerontol Geriatr*, 22(10), 15-19.
- Schmidt, S.J., Mueller, D.R., & Roder, V. (2011). Social cognition as a mediator variable between neurocognition and functional outcome in schizophrenia: empirical review and new results by structural equation modeling. *Schizophr Bull*, 37 Suppl 2, S41-54.
- Schultze-Lutter, F., Michel, C., Schmidt, S.J., Schimmelmann, B.G., Maric, N.P., Salokangas, R.K., Riecher-Rössler, A., Van der Gaag, M., Nordentoft, M., Raballo, A., Menezhelli, A., Marshall, M., Morrison, A., Ruhrmann, S., & Klosterkötter, J. (2015). EPA guidance on the early detection of clinical high risk states of psychoses. *Eur Psychiatry*, 30(3), 405-16.
- Schultze-Lutter, F., Ruhrmann, S., Berning, J., Maier, W., & Klosterkötter, J. (2010). Basic symptoms and ultrahigh risk criteria: symptom development in the initial prodromal state. *Schizophr bull*, 36(1), 182-91.
- Seidman, L.J., Giuliano, A.J., Meyer, E.C., Addington, J., Cadenhead, K.S., Cannon, T.D., McClashan, T.H., Perkins, D.O., Tsuang, M.T., Walker, E.F., Woods, S.W., Bearden, C.E., Christensen, B.K., Hawkins, K., Heaton, R., Keefe, R.S., Heinssen, R., Cornblatt, B.A. & North American Prodrome Longitudinal Study (NAPLS) Group. (2010). Neuropsychology of the prodrome to psychosis in the NAPLS consortium: relationship to family history and conversion to psychosis. *Arch Gen Psychiatry*, 67(6), 578-588.
- Seiferth, N.Y., Pauly, K., Habel, U., Kellermann, T., Shah, N.J., Ruhrmann, S., Klosterkötter, J., Schneider, F., & Kircher, T. (2008). Increased neural response related to neutral faces in individuals at risk for psychosis. *Neuroimage*, 40(1), 289-97.
- Semerari, A., Carcione, A., Dimaggio, G., Falcone, M., Nicolò, G., Procacci, M., & Alleva, G. (2003). How to evaluate metacognitive functioning in psychotherapy? The metacognition assessment scale and its applications. *Clin Psychol Psychother*, 10(4), 238-261.
- Semerari, A., Carcione, A., Dimaggio, G., Nicolò, G., & Procacci, M. (2007). Understanding minds, different functions and different disorders? The contribution of psychotherapeutic research. *Psychotherapy Research*, 17, 106-119.
- Sergi, M.J., Fiske, A.P., Horan, W.P., Kern, R.S., Kee, K.S., Subotnik, K.L., Nuechterlein, K.H., & Green, M.F. (2009). Development of a measure of

- relationship perception in schizophrenia. *Psychiatry Res*, 166(1), 54-62.
- Shad, M.U., Muddasani, S., Prasad, K., Sweeney, J.A., & Keshavana, M.S. (2004). Insight and prefrontal cortex in first-episode schizophrenia. *NeuroImage*, 22(3), 1315-1320.
- Shamay-Tsoory, S.G. (2011). The neural bases for empathy. *Neuroscientist*, 17(1), 18-24.
- Sheehan, D.V., Lecrubier, Y., Sheehan, K.H., Amorim, P., Janavs, J., Weiller, E., Herqueta, T., Baker, R., & Dunbar, G.C. (1998). The Mini-international Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry*, 59(20), 22-33.
- Singer, T. (2006). The neuronal basis and ontogeny of empathy and mind reading: review of literature and implications for future research. *Neurosci Biobehav Rev*, 30(6), 855-63.
- Smith, M.J., Horan, W.P., Karpouzian, T.M., Abram, S.V., Cobia, D.J., & Csernansky, J.G. (2012). Self-reported empathy deficits are uniquely associated with poor functioning in schizophrenia. *Schizophr Res*, 137(1-3), 196-202.
- Smith, M.J., Schroeder, M.P., Abram, S.V., Goldman, M.B., Parrish, T.B., Wang, X., Derrnfl, B., Habel, U., Decety, J., Reilly, J.L., Csernansky, J.G., & Breiter, H.C. (2015). Alterations in brain activation during cognitive empathy are related to social functioning in schizophrenia. *Schizophr Bull*, 41(1), 211-22.
- Snijders, T.A.B., & Bosker, R.J. (2012). *Multilevel Analysis: An Introduction to Basic and Advanced Multilevel Modeling*. London: Sage Publishers.
- Sprong, M., Schothorst, P., Vos, E., Hox, J., & Van Engeland, H. (2007). Theory of mind in schizophrenia: meta-analysis. *Br J Psychiatry*, 191, 5-13.
- Stanford, A.D., Messinger, J., Malaspina, D., & Corcoran, C.M. (2011). Theory of Mind in patients at clinical high risk for psychosis. *Schizophr Res*, 131(1-3), 11-7.
- Staring, A.B.P., Van Der Gaag, M., Van Den Berge, M., Duivenvoorden, H.J., & Mulder, C.L. (2009). Stigma moderates the associations of insight with depressed mood, low self-esteem and low quality of life in patients with schizophrenia spectrum disorders. *Schizophr. Res*, 115, 363-369.
- Stefanopoulou, E., Lafuente, A.R., Saez Fonseca, J.A., & Huxley, A. (2009). Insight, global functioning and psychopathology amongst in-patient clients with schizophrenia. *Psychiatr. Q*, 80, 155-165.
- Stone, V.E., Baron-Cohen, S., & Knight, R.T. (1998). Frontal lobe contributions to theory of mind. *J Cogn Neurosci*, 10, 640-656.
- Szily, E., & Kéri, S. (2013). Delusion Proneness and emotion appraisal in individuals with high psychosis vulnerability. *Clin Psychol Psychother*, 20, 166-170.
- The EuroQol group (1990). EuroQol - a new facility for the measurement of health related quality of life. *Health Policy*, 16, 199-208.
- Thompson, A.D., Bartholomeusz, C., & Yung, A.R. (2011). Social cognition deficits and the 'ultra high risk' for psychosis population: a review of literature. *Early Interv Psychiatry*, 5(3), 192-202.
- Thompson, A., Papas, A., Bartholomeusz, C., Allott, K., Amminger, G.P., Nelson, B., Wood, S., & Yung, A. (2012). Social cognition in clinical "at risk" for psychosis and first episode psychosis populations. *Schizophr Res*, 141(2-3), 204-9.
- Thompson, A., Papas, A., Bartholomeusz, C., Nelson, B., & Yung, A. (2013). Externalized attributional bias in the Ultra High Risk (UHR) for psychosis population. *Psychiatry Res*, 206(2-3), 200-5.
- Tombaugh, T.N. (2004). Trailmaking test A and B: normative data stratified by age and education. *Arch Clin Neuropsychol*, 19(2), 203-214.
- Tsapakis, E.M., Dimopoulou, T., & Tarazi, F.I. (2015). Clinical management of negative symptoms of schizophrenia: An update. *Pharmacology and Therapeutics*, 153, 135-147.
- Turkington, D., Kingdon, D., & Turner, T. (2002). Effectiveness of a brief cognitive-behavioral therapy intervention in the treatment of schizophrenia. *Br. J. Psychiatry*, 180, 523-527.
- Turkington, D., Kingdon, D., Rathod, S., Hammond, K., Pelton, J., & Metha, R. (2006). Outcomes of an effectiveness trial of cognitive behavioural intervention by mental health nurses in schizophrenia. *Br. J. Psychiatry*, 189, 36-40.

REFERENCES

- Umberson, D., & Montez, J.K. (2010). Social relationships and health: a flashpoint for health policy. *J Health Soc Behav*, 51 Suppl, S54-66.
- Van der Gaag, M., Schutz, C., Ten Napel, A., Landa, Y., Delespaul, P., Bak, M., Tschacher, W., & De Hert, M. (2013). Development of the Davos Assessment of Cognitive Biases Scale (DACOBS) *Schizophr Res*, 144(1-3), 63-71.
- Van Donkersgoed, R. J. M., de Jong, S., & Pijnenborg, G. H. M. (2016). Metacognitive Reflection and Insight Therapy (MERIT) with a Patient with Persistent Negative Symptoms. *Journal of Contemporary Psychotherapy*, 46(4), 245-253.
- Van Donkersgoed, R.J., De Jong, S., Van der Gaag, M., Aleman, A., Lysaker, P.H., Wunderink, L., & Pijnenborg, G.H. (2014). A manual-based individual therapy to improve metacognition in schizophrenia: protocol of a multi-center RCT. *BMC Psychiatry*, 14(1), 27.
- Van Os, J., & Guloksuz, S. (2017). A critique of the "ultra-high risk" and "transition" paradigm. *World Psychiatry*, 16(2), 200-206.
- Van Os, J., & Kapur, S. (2009). Schizophrenia. *Lancet*, 374(9690), 635-45.
- Van Rijn, S., Aleman, A., de Sonneville, L., Sprong, M., Ziermans, T., Schothorst, P., van Engeland, H., & Swaab, H. (2011). Misattribution of facial expressions of emotion in adolescents at increased risk of psychosis: the role of inhibitory control. *Psychol Med*, 41(3), 499-508.
- Van Veluw, S.J., & Chance, S.A. (2014). Differentiating between self and others: an ALE meta-analysis of fMRI studies of self-recognition and theory of mind. *Brain Imaging and Behavior*, 8(1), 24-38.
- Vauth, R., Rüschi, N., Wirtz, M., & Corrigan, P.W. (2004). Does social cognition influence the relation between neurocognitive deficits and vocational functioning in schizophrenia? *Psychiatry Res*, 128(2), 155-65.
- Velthorst, E., Nieman, D.H., Becker, H.E., van de Fliert, R., Dingemans, P.M., Klaassen, R., de Haan, L., van Amelsvoort, T., & Linszen, D.H. (2009). Baseline differences in clinical symptomatology between ultra high risk subjects with and without a transition to psychosis. *Schizophr Res*, 109, 60-5.
- Wechsler, D. (1995). *Manual for the Wechsler Adult Intelligence Scale*. Psychological Corp, Oxford, England.
- Wells, A. (2009). *Metacognitive Therapy for Anxiety and Depression*. New York: The Guilford Press.
- Wiffen, B.R.D., Rabinowitz, J., Fleischhacker, W.W., & David, A.S. (2010). Insight: demographic differences and associations with one-year outcome in schizophrenia and schizoaffective disorder. *Clin Schizophr Relat Psychoses*, 4(3), 169-175.
- Wilson, D.B. (2010). *Professional Development Course on Meta-analysis*. Available: <http://mason.gmu.edu/~dwilsonb/ma.html>.
- Wölwer, W., Combs, D.R., Frommann, N., & Penn, D.L. (2010). Treatment approaches with a special focus on social cognition: overview and empirical results. In: *Neurocognition and Social Cognition in Schizophrenia Patients: Basic Concepts and Treatment*, Karger Publishers, Basel, Switzerland, pp. 61-78.
- Wood, S.J., Yung, A.R., McGorry, P.D., & Pantalis, C. (2011). Neuroimaging and treatment evidence for clinical staging in psychotic disorders: from the at-risk mental state to chronic schizophrenia. *Biol Psychiatry*, 70(7), 619-25.
- Wunderink, L., Nieboer, R.M., Wiersma, D., Sytma, S., & Nienhuis, F.J. (2013). Recovery in remitted first-episode psychosis at 7 years of follow-up of an early dose reduction/discontinuation or maintenance treatment strategy: long-term follow-up of a 2-year randomized clinical trial. *JAMA Psychiatry*, 70(9), 913-20.
- Wykes, T., Steel, C., Everitt, B., & Tarrier, N. (2008). Cognitive behavior therapy for schizophrenia: Effect sizes, clinical models, and methodological rigor. *Schizophrenia Bulletin*, 34(3), 523-537.
- Xiang, Y., Weng, Y., Li, W., Gao, L., Chen, G., Xie, L., Chang, Y., Tang, W., & Ungvari, G.S. (2007). Efficacy of the Community Re-Entry Module for patients with schizophrenia in Beijing, China: outcome at 2-year follow-up. *Br. J. Psychiatry*, 190, 49-56.
- Yong, E., Barbato, M., Penn, D.L., Keefe, R.S., Woods, S.W., Perkins, D.O., & Addington, J. (2014). Exploratory analysis of social cognition and neurocognition in individuals at clinical high risk for psychosis. *Psychiatry Res*, 218(1-2), 39-43.
- Young A., Perrett D., Calder A., Sprengelmeyer R., & Ekman P. (2002). *Facial Expressions of*

Emotion: Stimuli and Tests (FEEST). Thames Valley Test Company, Bury St. Edmunds.

Yung, A.R., & McGorry, P.D. (2007). Prediction of psychosis: setting the stage. *Br J Psychiatry, Suppl 51*, S1-8.

Zaki, J., Bolger, N., & Ochsner, K. (2008). It takes two: the interpersonal nature of empathic accuracy. *Psychol Sci, 19*(4), 399-404.

Zaki, J., & Ochsner, K. (2009). The need for a cognitive neuroscience of naturalistic social cognition. *Ann N Y Acad Sci, 1167*, 16-30.

Zaki, J., Bolger, N., & Ochsner, K. (2009). Unpacking the informational bases of empathic accuracy. *Emotion, 9*, 478-487.

Zaki, J., Weber, J., Bolger, N., & Ochsner, K. (2009). The neural bases of empathic accuracy. *Proceedings of the National Academy of Sciences USA, 106*, 11382-11387.

Zaki, J., Hennigan, K., Weber, J., & Ochsner, K. (2010). Social cognitive conflict resolution: contributions of domain-general and domain-specific neural systems. *Journal of Neuroscience, 30*, 8481-8488.

REFERENCES