

University of Groningen

## Complete Coding Sequences of Five Dengue Virus Type 2 Clinical Isolates from Venezuela Obtained through Shotgun Metagenomics

Lizarazo, Erley; Couto, Natacha; Vincenti-Gonzalez, Maria; Raangs, Erwin C; Jaenisch, Thomas; Friedrich, Alex W; Tami, Adriana; Rossen, John W

*Published in:*  
Genome Announcements

*DOI:*  
[10.1128/genomeA.00545-18](https://doi.org/10.1128/genomeA.00545-18)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2018

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Lizarazo, E., Couto, N., Vincenti-Gonzalez, M., Raangs, E. C., Jaenisch, T., Friedrich, A. W., ... Rossen, J. W. (2018). Complete Coding Sequences of Five Dengue Virus Type 2 Clinical Isolates from Venezuela Obtained through Shotgun Metagenomics. *Genome Announcements*, 6(25), [UNSP e00545-18]. <https://doi.org/10.1128/genomeA.00545-18>


### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*



# Complete Coding Sequences of Five Dengue Virus Type 2 Clinical Isolates from Venezuela Obtained through Shotgun Metagenomics

Erley Lizarazo,<sup>a</sup>  Natacha Couto,<sup>a</sup> Maria Vincenti-Gonzalez,<sup>a</sup> Erwin C. Raangs,<sup>a</sup> Thomas Jaenisch,<sup>b</sup> Alex W. Friedrich,<sup>a</sup> Adriana Tami,<sup>a,c</sup> John W. Rossen<sup>a</sup>

<sup>a</sup>Department of Medical Microbiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

<sup>b</sup>University of Heidelberg, Heidelberg, Germany

<sup>c</sup>Departamento de Parasitología, Facultad de Ciencias de la Salud, Universidad de Carabobo, Valencia, Venezuela

**ABSTRACT** Dengue is a disease endemic in Latin American countries, like Venezuela, and has become one of the most important public health problems. We report five complete coding sequences of dengue virus serotype 2 (DENV-2) isolated from DENV-infected patients in Venezuela. Phylogenetic analysis placed the isolates within the American/Asian genotype.

Dengue virus (DENV) infection continues to be one of the most prevalent arboviral diseases in tropical and subtropical regions, with an estimated burden of 390 million cases/year worldwide (1). Infection by any of the four DENV serotypes (DENV-1 to DENV-4) can lead to a wide spectrum of clinical outcomes, ranging from asymptomatic cases, mild disease as a flu-like syndrome (dengue without warning signs), to a more severe form of disease (dengue with warning signs or severe dengue) (1–3). Severe disease is frequently associated with the Asian DENV-2 and DENV-3 genotypes with secondary infections (4). To date, there is no fully successful vaccine or specific treatment for DENV infection (5, 6).

We report five complete coding sequences of dengue virus serotype 2 (DENV-2) isolated from DENV-infected patients in Venezuela in 2015, of whom two patients had dengue with warning signs. The study was approved by the ethics review committee of the Biomedical Research Institute, Carabobo University [Aval Bioetico numbers CBIIB(UC)-014 and CBIIB-(UC)-2013-1], Maracay, Venezuela, and the Ethics, Bioethics and Biodiversity Committee of the National Foundation for Science, Technology and Innovation, Caracas, Venezuela. For sequencing, RNA isolation was performed with the QIAamp viral RNA isolation kit (Qiagen, Hilden, Germany). Libraries were prepared with the TruSeq V2 RNA (Illumina, San Diego, CA, USA), which includes a cDNA synthesis step. Sequencing was performed on a MiSeq instrument with the MiSeq reagent kit version 2 (Illumina) that generated 150-bp paired-end reads. The sequences were assembled and analyzed using the CLC Genomics Workbench version 10.1.1 software (Qiagen). Genome annotation was performed using the plugin MetaGeneMark version 1.4.

The complete open reading frame (ORF) of the DENV-2 polyprotein was obtained in all samples with a coverage of greater than 177-fold. The lengths of the genomes sequenced were 10,694 nucleotides (nt), 10,619 nt, 10,712 nt, 10,711 nt, and 10,704 nt, with the lengths of the 5'/3' untranslated regions being 87/431 nt, 87/356 nt, 84/452 nt, 86/449 nt, and 77/451 nt, respectively. These are slightly shorter than the 96/451 nt reported for the DENV-2 reference genome (GenBank accession number NC\_001474).

Received 21 May 2018 Accepted 21 May 2018 Published 21 June 2018

**Citation** Lizarazo E, Couto N, Vincenti-Gonzalez M, Raangs EC, Jaenisch T, Friedrich AW, Tami A, Rossen JW. 2018. Complete coding sequences of five dengue virus type 2 clinical isolates from Venezuela obtained through shotgun metagenomics. *Genome Announc* 6:e00545-18. <https://doi.org/10.1128/genomeA.00545-18>.

**Copyright** © 2018 Lizarazo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Natacha Couto, [n.monge.gomes.do.couto@umcg.nl](mailto:n.monge.gomes.do.couto@umcg.nl).

The phylogenetic analysis based on the complete ORF using the maximum likelihood method revealed that the isolates belong to the American/Asian genotype. Strains clustered in two different subpopulations, sharing a common ancestor within the Venezuelan clade.

Molecular surveillance to monitor circulating DENV-2 strains in Venezuela is lacking nowadays, and any information about current or past circulating strains has not been reported since 2008 (7). The replacement of less virulent strains by the (re)emergence of more virulent strains or *in situ* recombination has happened before in Venezuela; consequently, such potential changes should be monitored to reveal evolutionary trends (8). The sequences of DENV-2 described in this work will help follow up on the molecular epidemiology of DENV in Venezuela. In addition, these genome sequences add to the knowledge of the current DENV-2 diversity and endemicity, which are important for developing future accurate vaccines.

**Accession number(s).** The complete coding sequences of the five DENV-2 strains described here have been deposited in GenBank under the accession numbers [MH069495](#) (DENV-2/VE/IDAMS/921099), [MH069496](#) (DENV-2/VE/IDAMS/921096), [MH069497](#) (DENV-2/VE/IDAMS/921095), [MH069498](#) (DENV-2/VE/IDAMS/910121), and [MH069499](#) (DENV-2/VE/IDAMS/910105).

## ACKNOWLEDGMENTS

This study was partially supported by the National Science, Technology and Innovation Funds (FONACIT), project number 2011000303 under contract number 201100129, and by the International Research Consortium on Dengue Risk Assessment, Management and Surveillance (IDAMS), funded by FP7-HEALTH-2011, grant number 281803, during data and sample collection in Venezuela.

Erley Lizarazo and Maria Vincenti-Gonzalez were funded by the Abel Tasman Talent Program.

## REFERENCES

- Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, Drake JM, Brownstein JS, Hoen AG, Sankoh O, Myers MF, George DB, Jaenisch T, Wint GRW, Simmons CP, Scott TW, Farrar JJ, Hay SI. 2013. The global distribution and burden of dengue. *Nature* 496:504–507. <https://doi.org/10.1038/nature12060>.
- Weaver SC, Vasilakis N. 2009. Molecular evolution of dengue viruses: contributions of phylogenetics to understanding the history and epidemiology of the preeminent arboviral disease. *Infect Genet Evol* 9:523–540. <https://doi.org/10.1016/j.meegid.2009.02.003>.
- WHO/TDR. 2012. Handbook for clinical management of dengue. World Health Organization, Geneva, Switzerland.
- Leitmeyer KC, Vaughn DW, Watts DM, Salas R, Villalobos I, de Chacon Ramos C, Rico-Hesse R. 1999. Dengue virus structural differences that correlate with pathogenesis. *J Virol* 73:4738–4747.
- The Lancet Infectious Diseases. 2018. The dengue vaccine dilemma. *Lancet Infect Dis* 18:123. [https://doi.org/10.1016/S1473-3099\(18\)30023-9](https://doi.org/10.1016/S1473-3099(18)30023-9).
- Yoon I-K, Thomas SJ. 2018. Encouraging results but questions remain for dengue vaccine. *Lancet Infect Dis* 18:125–126. [https://doi.org/10.1016/S1473-3099\(17\)30634-5](https://doi.org/10.1016/S1473-3099(17)30634-5).
- Rodriguez-Roche R, Blanc H, Bordería AV, Díaz G, Henningsson R, Gonzalez D, Santana E, Alvarez M, Castro O, Fontes M, Vignuzzi M, Guzman MG. 2016. Increasing clinical severity during a dengue virus type 3 Cuban epidemic: deep sequencing of evolving viral populations. *J Virol* 90:4320–4333. <https://doi.org/10.1128/JVI.02647-15>.
- Ali A, Ali I. 2015. The complete genome phylogeny of geographically distinct dengue virus serotype 2 isolates (1944–2013) supports further groupings within the cosmopolitan genotype. *PLoS One* 10:e0138900. <https://doi.org/10.1371/journal.pone.0138900>.