

University of Groningen

Physical activity in hard-to-reach physically disabled people

Krops, Leonie

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Krops, L. (2018). *Physical activity in hard-to-reach physically disabled people: Development, implementation and effectiveness of a community-based intervention*. [Groningen]: Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

References

1. Adler RH. Engel's biopsychosocial model is still relevant today. *J. Psychosom. Res.* 2009;67:607–11.
2. Engel GL. The biopsychosocial model and the education of health professionals. *Gen. Hosp. Psychiatry.* 1979;1:156–65.
3. Huber M, Knottnerus JA, Green L, Van Der Horst H, Jadad AR, Kromhout D, et al. How should we define health? *BMJ.* 2011;343.
4. Hays RD, Hahn H, Marshall G. Use of the SF-36 and other health-related quality of life measures to assess persons with disabilities. *Arch. Phys. Med. Rehabil.* 2002;83:S4-s9.
5. Gellert GA. The importance of quality of life research for health care reform in the USA and the future of public health. *Qual. Life Res.* 1993;2:357–61.
6. Lee IM, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT, et al. Effect of physical inactivity on major non-communicable diseases worldwide: An analysis of burden of disease and life expectancy. *Lancet.* 2012;380:219–29.
7. Sallis JF, Bull F, Guthold R, Heath GW, Inoue S, Kelly P, et al. Progress in physical activity over the Olympic quadrennium. *Lancet.* 2016;388:1325–36.
8. Warburton DE, Bredin SS. Health benefits of physical activity: a systematic review of current systematic reviews. *Curr Opin Cardiol.* 2017;32:541–56.
9. Liffiton JA, Horton S, Baker J, Weir PL. Successful aging: How does physical activity influence engagement with life? *Eur. Rev. Aging Phys. Act.* 2012;9:103–8.
10. Pinto Pereira SM, Geoffroy MC, Power C. Depressive Symptoms and Physical Activity During 3 Decades in Adult Life. *JAMA Psychiatry.* 2014;71:1373.
11. Marlier M, Van Dyck D, Cardon G, De Bourdeaudhuij I, Babiak K, Willem A. Interrelation of sport participation, physical activity, social capital and mental health in disadvantaged communities: A SEM-analysis. *PLoS One.* 2015;10:e0140196.
12. Cowan RE. Exercise Is Medicine Initiative: Physical Activity as a Vital Sign and Prescription in Adult Rehabilitation Practice. *Arch. Phys. Med. Rehabil.* 2016;97:S232–7.
13. Morris JN. Exercise in the prevention of coronary heart disease: today's best buy in public health. *Med. Sci. Sports Exerc.* 1994;26:807–14.
14. Das P, Horton R. Physical activity—time to take it seriously and regularly. *Lancet.* 2016;388:1254–5.
15. Nederlandse Gezondheidsraad. Beweegrichtlijnen 2017. [Dutch Health Council. Physical activity recommendations 2017]. 2017.
16. Pratt M, Sarmiento OL, Montes F, Ogilvie D, Marcus BH, Perez LG, et al. The implications of megatrends in information and communication technology and transportation for changes in global physical activity. *Lancet.* 2012;380:282–93.
17. Ding D, Lawson KD, Kolbe-Alexander TL, Finkelstein EA, Katzmarzyk PT, van Mechelen W, et al. The economic burden of physical inactivity: a global analysis of major non-communicable diseases. *Lancet.* 2016;388:1311–24.

18. Andersen LB, Mota J, Di Pietro L. Update on the global pandemic of physical inactivity. *Lancet*. 2016;388:1255–6.
19. World Health Organisation. *World Report On Disability*. Geneva, Switzerland: WHO Press; 2011.
20. von Heijden A, van den Dool R, van Lindert C, Breedveld K. (On)beperkt sportief 2013. Monitor sport- en beweegdeelname van mensen met een handicap [(Un)limited exercise 2013. Monitoring sport- and exercise participation in handicapped people]. Utrecht, the Netherlands: Mulier Instituut / Arko Sports Media; 2013.
21. Van den Brink C, Savelkoul M. Gezondheidsmonitor GGD'en, CBS en RIVM [Health monitor Joint Health Duties, Statistics Netherlands and National Institute for Health and Environment]. *Volksgezond. Toekomst Verkenning, Natl. Kompas Volksgezond*. 2013.
22. Seeman TE, Merkin SS, Crimmins EM, Karlamangla AS. Disability trends among older Americans: National Health and Nutrition Examination surveys, 1988-1994 and 1999-2004. *Am. J. Public Health*. 2010;100:100–7.
23. Heath GW, Fentem PH. Physical activity among persons with disabilities--a public health perspective. *Exerc. Sport Sci. Rev*. 1997;25:195–234.
24. Rosenberg DE, Bombardier CH, Hoffman JM, Belza B. Physical Activity Among Persons Aging with Mobility Disabilities: Shaping a Research Agenda. *J. Aging Res*. 2011;2011:1–16.
25. Van Den Berg-Emons RJ, Bussmann JB, Stam HJ. Accelerometry-based activity spectrum in persons with chronic physical conditions. *Arch. Phys. Med. Rehabil*. 2010;91:1856–61.
26. Rimmer JH, Marques AC. Physical activity for people with disabilities. *Lancet*. 2012;380:193–5.
27. Rimmer JH, Braddock D, Pitetti KH. Research on physical activity and disability: an emerging national priority. *Med. Sci. Sport. Exerc*. 1996;28:1366–72.
28. Khoo S, Li C, Ansari P. The Top 50 Most Cited Publications in Disability Sport: A Bibliometric Analysis. *Percept. Mot. Skills*. 2018;125:525–45.
29. Martin Ginis KA, Ma JK, Latimer-Cheung AE, Rimmer J. A Systematic Review of Review Articles Addressing Factors Related to Physical Activity Participation Among Children and Adults with Physical Disabilities. *Health Psychol. Rev*. 2016;7199:1–31.
30. Rimmer JH, Riley B, Wang E, Rauworth A, Jurkowski J. Physical activity participation among persons with disabilities: Barriers and facilitators. *Am. J. Prev. Med*. 2004;26:419–25.
31. McLeroy KR, Bibeau D, Steckler A, Glanz K. An ecological perspective on health promotion programs. *Health Educ. Q*. 1988;15:351–77.
32. Jaarsma EA, Dekker R, Geertzen JHB, Dijkstra PU. Sports participation after rehabilitation: Barriers and facilitators. *J. Rehabil. Med*. 2016;48:72–9.

References

33. Cervantes CM, Taylor WC. Physical Activity Interventions in Adult Populations With Disabilities: A Review. *Quest*. 2011;63:385–410.
34. Rimmer JH, Chen MD, McCubbin JA, Drum C, Peterson J. Exercise intervention research on persons with disabilities: what we know and where we need to go. *Am J Phys Med Rehabil*. 2010;89:249–63.
35. Alingh RA, Hoekstra F, van der Schans CP, Hettinga FJ, Dekker R, van der Woude LHV. Protocol of a longitudinal cohort study on physical activity behaviour in physically disabled patients participating in a rehabilitation counselling programme: ReSpAct. *BMJ Open*. 2015;5:e007591.
36. Hoekstra F, Hettinga FJ, Alingh RA, Duijf M, Dekker R, van der Woude LHV, et al. The current implementation status of the integration of sports and physical activity into Dutch rehabilitation care. *Disabil. Rehabil*. 2017;39:181–6.
37. Schutte BAM, Haveman-Nies A, Preller L. One-Year Results of the BeweegKuur Lifestyle Intervention Implemented in Dutch Primary Healthcare Settings. *Biomed Res. Int*. 2015;2015:1–7.
38. Berendsen BAJ, Kremers SPJ, Savelberg HHCM, Schaper NC, Hendriks MRC. The implementation and sustainability of a combined lifestyle intervention in primary care: mixed method process evaluation. *BMC Fam. Pract*. 2015;16:37.
39. Gehandicaptensport Nederland. Special Heroes evaluatie 2009/2012 [Special Heroes evaluation 2009/2012]. Bunnik, the Netherlands: 2013.
40. Desveaux L, Beauchamp M, Goldstein R, Brooks D. Community-based Exercise Programs as a Strategy to Optimize Function in Chronic Disease: A systematic review. *Med. Care*. 2014;52:216–26.
41. Van der Ploeg HP, van der Beek AJ, van der Woude LH, van Mechelen W. Physical activity for people with a disability: a conceptual model. *Sports Med*. 2004;34:639–49.
42. World Health Organization. International Classification of Functioning, Disability and Health: ICF. 1st ed. Geneva, Switzerland: 2001.
43. De Vries H, Dijkstra M, Kuhlman P. Self-efficacy: the third factor besides attitude and subjective norm as a predictor of behavioural intentions. *Heal. Educ Res*. 1988;3:273–82.
44. Bartholomew LK, Parcel GS, Kok G, Gottlieb NH, Fernández ME. *Planning Health Promotion Programs; An Intervention Mapping Approach*. San Fransisco: Jossey-Bass; 2011.
45. Glasgow RE, Vogt TM, Boles SM. Evaluating the public health impact of health promotion interventions: the RE-AIM framework. *Am. J. Public Health*. 1999;89:1322–7.
46. Whitehurst DGT, Engel L, Bryan S. Short Form health surveys and related variants in spinal cord injury research: A systematic review. *J Spinal Cord Med*. 2014;37:128–38.
47. Hays RD, Morales LS. The RAND-36 measure of health-related quality of life. *Ann. Med*. 2001;33:350–7.

48. Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med. Care.* 1992;30:473–83.
49. Hays RD, Sherbourne CD, Mazel RM. The RAND 36-Item Health Survey 1.0. *Health Econ.* 1993;2:217–27.
50. Van der Zee KI, Sanderman R. Measuring general health status with the RAND-36. users manual [Het meten van de algemene gezondheidstoestand met de RAND-36. een handleiding]. 2nd edition. Groningen, the Netherlands: Research Institute SHARE. 2012.
51. Van der Zee KI, Sanderman R, Heyink JW. Psychometric qualities of the MOS 36-item Short Form Health Survey (SF-36) in a Dutch population [De psychometrische kwaliteiten van de MOS 36-item Short Form health Survey (SF-36) in een Nederlandse populatie]. *T.Soc.Gezondheidsz.* 1993;71:183–91.
52. Van Baalen B, Odding E, van Woensel MP, Roebroek ME. Reliability and sensitivity to change of measurement instruments used in a traumatic brain injury population. *Clin Rehabil.* 2006;20:686–700.
53. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet.* 1986;1:307–10.
54. Weir JPJ. Quantifying test-retest reliability using the intraclass correlation coefficient and the SEM. *J. Strength Cond. Res.* 2005;19:231–40.
55. Ruta DA, Abdalla MI, Garratt AM, Coutts A, Russell IT. SF 36 health survey questionnaire: I. Reliability in two patient based studies. *Qual. Health Care.* 1994;3:180–5.
56. Vickrey BG, Hays RD, Genovese BJ, Myers LW, Ellison GW. Comparison of a generic to disease-targeted health-related quality-of-life measures for multiple sclerosis. *J. Clin. Epidemiol.* 1997;50:557–69.
57. Dorman P, Slattery J, Farrell B, Dennis M, Sandercock P. Qualitative comparison of the reliability of health status assessments with the EuroQol and SF-36 questionnaires after stroke. United Kingdom Collaborators in the International Stroke Trial. *Stroke.* 1998;29:63–8.
58. Saleh KJ, Radosevich DM, Kassim RA, Moussa M, Dykes D, Bottolfson H, et al. Comparison of commonly used orthopaedic outcome measures using palm-top computers and paper surveys. *J. Orthop. Res.* 2002;20:1146–51.
59. Dunn KM, Jordan K, Croft PR. Does questionnaire structure influence response in postal surveys? *J. Clin. Epidemiol.* 2003;56:10–6.
60. Boyer F, Morrone I, Laffont I, Dizien O, Etienne JC, Novella JL. Health related quality of life in people with hereditary neuromuscular diseases : An investigation of test - retest agreement with comparison between two generic questionnaires , the Nottingham health profile and the short form-36 items. *Neuromuscul. Disord.* 2006;16:99–106.
61. Ferrer M, Pellisé F, Escudero O, Alvarez L, Pont A, Alonso J, et al. Validation of a minimum outcome core set in the evaluation of patients with back pain. *Spine.* 2006;31:1372–9.

References

62. Lin MR, Hwang HF, Chen CY, Chiu WT. Comparisons of the brief form of the World Health Organization Quality of Life and Short Form-36 for persons with spinal cord injuries. *Am. J. Phys. Med. Rehabil.* 2007;86:104-13.
63. Steffen T, Seney M. Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-item short-form health survey, and the unified Parkinson disease rating scale in people with parkinsonism. *Phys. Ther.* 2008;88:733-46.
64. Marx RG, Menezes A, Horovitz L, Jones EC, Warren RF. A comparison of two time intervals for test-retest reliability of health status instruments. *J Clin Epidemiol.* 2003;56:730-5.
65. McHorney CA, Tarlov AR. Individual-patient monitoring in clinical practice: are available health status surveys adequate? *Qual. Life Res.* 1995;4:293-307.
66. Terwee CB, Bot SDM, de Boer MR, van der Windt DAWM, Knol DL, Dekker J, et al. Quality criteria were proposed for measurement properties of health status questionnaires. *J. Clin. Epidemiol.* 2007;60:34-42.
67. Aaronson NK, Muller M, Cohen PDA, Essink-Bot ML, Fekkes M, Sanderman R, et al. Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *J. Clin. Epidemiol.* 1998;51:1055-68.
68. Streiner DL, Norman GR. Health measurement scales, a practical guide to their development and use. United States: Oxford University Press; 2008.
69. Engel GL. The need for a new medical model: a challenge for biomedicine. *Science.* 1977;196:129-36.
70. Nicassio PM, Kay MA, Custodio MK, Irwin MR, Olmstead R, Weisman MH. An evaluation of a biopsychosocial framework for health-related quality of life and disability in rheumatoid arthritis. *J. Psychosom. Res.* 2011;71:79-85.
71. Gavelova M, Nagyova I, Rosenberger J, Krokavcova M, Gdovinova Z, Groothoff JW, et al. Importance of an individual's evaluation of functional status for health-related quality of life in patients with multiple sclerosis. *Disabil. Health J.* 2015;8:372-9.
72. Schrier E, Schrier I, Geertzen JH, Dijkstra PU. Quality of life in rehabilitation outpatients: normal values and a comparison with the general Dutch population and psychiatric patients. *Qual. Life Res.* 2016;25:135-42.
73. Kivisild A, Sabre L, Tomberg T, Ruus T, Korv J, Asser T, et al. Health-related quality of life in patients with traumatic spinal cord injury in Estonia. *Spinal Cord.* 2014;52:570-5.
74. Schoppen T, Boonstra A, Groothoff JW, De Vries J, Goeken LN, Eisma WH. Job satisfaction and health experience of people with a lower-limb amputation in comparison with healthy colleagues. *Arch. Phys. Med. Rehabil.* 2002;83:628-34.
75. McCutcheon T, Knepp T, Richards N, Sparks M. Comparison of quality of life of persons who have experienced amputations and persons who have had bowel resections. *Gastroenterol. Nurs.* 2005;28:221-6.

76. Van der Schans CP, Geertzen JH, Schoppen T, Dijkstra PU. Phantom pain and health-related quality of life in lower limb amputees. *J Pain Symptom Manage.* 2002;24:429–36.
77. Aymerich M, Guillamon I, Jovell AJ. Health-related quality of life assessment in people with multiple sclerosis and their family caregivers. A multicenter study in Catalonia (Southern Europe). *Patient Prefer. Adherence.* 2009;3:311–21.
78. Klevan G, Jacobsen CO, Aarseth JH, Myhr KM, Nyland H, Glad S, et al. Health related quality of life in patients recently diagnosed with multiple sclerosis. *Acta Neurol. Scand.* 2014;129:21–6.
79. Miller A, Dishon S. Health-related quality of life in multiple sclerosis: The impact of disability, gender and employment status. *Qual. Life Res.* 2006;15:259–71.
80. Riazi A, Hobart JC, Lamping DL, Fitzpatrick R, Freeman JA, Jenkinson C, et al. Using the SF-36 measure to compare the health impact of multiple sclerosis and Parkinson's disease with normal population health profiles. *J. Neurol. Neurosurg. Psychiatry.* 2003;74:710–4.
81. Fernandez-Munoz JJ, Moron-Verdasco A, Cigaran-Mendez M, Munoz-Hellin E, Perez-de-Heredia-Torres M, Fernandez-de-Las-Penas C. Disability, quality of life, personality, cognitive and psychological variables associated with fatigue in patients with multiple sclerosis. *Acta Neurol. Scand.* 2015;132:118-24.
82. Baumstarck K, Pelletier J, Butzkueven H, Fernandez O, Flachenecker P, Idiman E, et al. Health-related quality of life as an independent predictor of long-term disability for patients with relapsing-remitting multiple sclerosis. *Eur. J. Neurol.* 2013;20:907–14.
83. Grytten N, Aarseth JH, Espeset K, Berg Johnsen G, Wehus R, Lund C, et al. Health-related quality of life and disease-modifying treatment behaviour in relapsing-remitting multiple sclerosis--a multicentre cohort study. *Acta Neurol. Scand.* 2012;126:51–7.
84. Fernandez O, Baumstarck-Barrau K, Simeoni MC, Auquier P, group M study. Patient characteristics and determinants of quality of life in an international population with multiple sclerosis: assessment using the MusiQoL and SF-36 questionnaires. *Mult. Scler.* 2011;17:1238–49.
85. Szilasiova J, Krokavcova M, Gdovinova Z, Rosenberger J, Van Dijk JP. Quality of life in patients with multiple sclerosis in Eastern Slovakia. *Disabil. Rehabil.* 2011;33:1587–93.
86. Drulovic J, Pekmezovic T, Matejic B, Mesaros S, Manigoda M, Dujmovic I, et al. Quality of life in patients with multiple sclerosis in Serbia. *Acta Neurol. Scand.* 2007;115:147–52.
87. Forbes A, While A, Mathes L, Griffiths P. Health problems and health-related quality of life in people with multiple sclerosis. *Clin. Rehabil.* 2006;20:67–78.
88. Hermann BP, Vickrey B, Hays RD, Cramer J, Devinsky O, Meador K, et al. A comparison of health-related quality of life in patients with epilepsy, diabetes and multiple sclerosis. *Epilepsy Res.* 1996;25:113–8.

References

89. Arango-Lasprilla JC, Nicholls E, Olivera SL, Perdomo JL, Arango JA. Health-related quality of life in individuals with spinal cord injury in Colombia, South America. *NeuroRehabilitation*. 2010;27:313–9.
90. Ebrahimzadeh MH, Golhasani-Keshtan F, Shojaee BS. Correlation between health-related quality of life in veterans with chronic spinal cord injury and their caregiving spouses. *Arch. trauma Res*. 2014;3:e16720.
91. Rognoni C, Fizzotti G, Pistarini C, Quaglini S. Quality of life of patients with spinal cord injury in Italy: preliminary evaluation. *Stud. Health Technol. Inform*. 2014;205:935–9.
92. Tramonti F, Gerini A, Stampacchia G. Individualised and health-related quality of life of persons with spinal cord injury. *Spinal Cord*. 2014;52:231–5.
93. Lidal IB, Veenstra M, Hjeltnes N, Biering-Sorensen F. Health-related quality of life in persons with long-standing spinal cord injury. *Spinal Cord*. 2008;46:710–5.
94. Forchheimer M, McAweeney M, Tate DG. Use of the SF-36 among persons with spinal cord injury. *Am. J. Phys. Med. Rehabil*. 2004;83:390–5.
95. Leduc BE, Lepage Y. Health-related quality of life after spinal cord injury. *Disabil. Rehabil*. 2002;24:196–202.
96. Andresen EM, Fouts BS, Romeis JC, Brownson CA. Performance of health-related quality-of-life instruments in a spinal cord injured population. *Arch. Phys. Med. Rehabil*. 1999;80:877–84.
97. Barrera M, Teall T, Barr R, Silva M, Greenberg M. Health related quality of life in adolescent and young adult survivors of lower extremity bone tumors. *Pediatr. Blood Cancer*. 2012;58:265–73.
98. Sinha R, van den Heuvel WJ, Arokiasamy P. Factors affecting quality of life in lower limb amputees. *Prosthet. Orthot. Int*. 2011;35:90–6.
99. Davidson JH, Khor KE, Jones LE. A cross-sectional study of post-amputation pain in upper and lower limb amputees, experience of a tertiary referral amputee clinic. *Disabil. Rehabil*. 2010;32:1855–62.
100. Taghipour H, Moharamzad Y, Mafi AR, Amini A, Naghizadeh MM, Soroush MR, et al. Quality of life among veterans with war-related unilateral lower extremity amputation: a long-term survey in a prosthesis center in Iran. *J. Orthop. Trauma*. 2009;23:525–30.
101. Nordeman L, Gunnarsson R, Mannerkorpi K. Prevalence and characteristics of widespread pain in female primary health care patients with chronic low back pain. *Clin. J. Pain*. 2012;28:65–72.
102. Lauche R, Cramer H, Langhorst J, Dobos G. Cupping for chronic nonspecific neck pain: a 2-year follow-up. *Forsch. Komplementmed*. 2013;20:328–33.
103. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR, Tudor-Locke C, et al. 2011 compendium of physical activities: A second update of codes and MET values. *Med. Sci. Sports Exerc*. 2011;43:1575–81.

104. Van der Zee KI, Sanderman R, Heyink JW, de Haes H. Psychometric qualities of the RAND 36-Item Health Survey 1.0: a multidimensional measure of general health status. *Int. J. Behav. Med.* 1996;3:104–22.
105. Hazell ML, Morris JA, Linehan MF, Frank PI, Frank TL. Factors influencing the response to postal questionnaire surveys about respiratory symptoms. *Prim. Care Respir. J.* 2009;18:165–70.
106. Norman GR, Sloan JA, Wyrwich KW. Interpretation of changes in health-related quality of life: the remarkable universality of half a standard deviation. *Med. Care.* 2003;41:582–92.
107. Haskell WL, Lee I-M, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med. Sci. Sports Exerc.* 2007;39:1423–34.
108. Mulroy SJ, Thompson L, Kemp B, Hatchett PP, Newsam CJ, Lupold DG, et al. Strengthening and optimal movements for painful shoulders (STOMPS) in chronic spinal cord injury: a randomized controlled trial. *Phys. Ther.* 2011;91:305–24.
109. Hicks AL, Martin Ginis KA, Ditor DS, Latimer AE, Craven C, Bugaresti J, et al. Long-term exercise training in persons with spinal cord injury: effects on strength, arm ergometry performance and psychological well-being. *Spinal Cord.* 2003;41:34–43.
110. Rau B, Bonvin F, de Bie R. Short-term effect of physiotherapy rehabilitation on functional performance of lower limb amputees. *Prosthet. Orthot. Int.* 2007;31:258–70.
111. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep.* 1985;100:126–31.
112. Rijksinstituut voor Volksgezondheid en Milieu. Centrum Gezond Leven [National institute for health and environment. Center for healthy life]. 2016 [cited 2016 Jul 7]; Available from: <https://www.loketgezondleven.nl/leefstijlinterventies/interventiedatabase-gezonden-actief-leven>
113. Kenniscentrum Sport. Netwerk in Beweging [Knowledge center Sport. Network in movement]. Available from: www.netwerkinbeweging.nl/interventies
114. Race KE, Hotch DF, Packer T. Rehabilitation Programm Evaluation; Use of Focus Groups to Empower Clients. *Eval. rev.* 1994;18:730–40.
115. Braun V, Clarke V. Using Thematic Analysis in Psychology. *Qual Res Psych.* 2006;3:77–101.
116. Braun V, Clarke V. What can “thematic analysis” offer health and wellbeing researchers? *Int J Qual. Stud Heal. Well-being.* 2014;9:26152.
117. Rimmer JH, Vanderbom KA, Bandini LG, Drum CE, Luken K, Suarez-Balcazar Y, et al. GRAIDs: a framework for closing the gap in the availability of health promotion programs and interventions for people with disabilities. *Implement. Sci.* 2014;9:100.

118. Murphy NA, Carbone PS. Promoting the participation of children with disabilities in sports, recreation, and physical activities. *Pediatrics*. 2008;121:1057–61.
119. Baker PR, Francis DP. Cochrane review: Community-wide interventions for increasing physical activity. *J. Evid. Based. Med.* 2015;8:57–8.
120. Geidl W, Semrau J, Pfeifer K. Health behaviour change theories: contributions to an ICF-based behavioural exercise therapy for individuals with chronic diseases. *Disabil. Rehabil.* 2014;36:2091-10.
121. Ajzen I. The theory of planned behavior. *Organ. Behav. Hum. Decis. Process.* 1991;50:179–211.
122. Payne W, Reynolds M, Brown S, Fleming A. Sports Role Models and Their Impact on Participation in Physical Activity: a Literature Review. University of Ballarat. 2003.
123. Chang FH, Wang YH, Jang Y, Wang CW. Factors associated with quality of life among people with spinal cord injury: application of the International Classification of Functioning, Disability and Health model. *Arch. Phys. Med. Rehabil.* 2012;93:2264–70.
124. Ekelund U, Steene-Johannessen J, Brown WJ, Fagerland MW, Owen N, Powell KE, et al. Can physical activity eliminate the detrimental association of sitting time with mortality? A harmonized meta-analysis of data from more than one million men and women. *Lancet*. 2016;6736:1–10.
125. Krops LA, Hols DH, Folkertsma N, Dijkstra PU, Geertzen JH, Dekker R. Requirements on a community-based intervention for stimulating physical activity in physically disabled people: a focus group study amongst experts. *Disabil. Rehabil.* 2017;Epub ahead of print.
126. van der Ploeg HP, Streppel KRM, van der Beek AJ, van der Woude LHV., Vollenbroek-Hutten MMR, van Harten WH, et al. Successfully Improving Physical Activity Behavior after Rehabilitation. *Am. J. Heal. Promot.* 2007;21:153–9.
127. Morris JH, Macgillivray S, McFarlane S. Interventions to promote long-term participation in physical activity after stroke: A systematic review of the literature. *Arch. Phys. Med. Rehabil.* 2014;95:956–67.
128. Oliveira CB, Franco MR, Maher CG, Tiedemann A, Silva FG, Damato TM, et al. The efficacy of a multimodal physical activity intervention with supervised exercises, health coaching and an activity monitor on physical activity levels of patients with chronic, nonspecific low back pain (Physical Activity for Back Pain (PAYBACK) trial. *Trials*. 2018;19:40.
129. Coultas DB, Jackson BE, Russo R, Peoples J, Singh KP, Sloan J, et al. Home-Based Physical Activity Coaching, Physical Activity, and Healthcare Utilization in COPD: COPD-SMART Secondary Outcomes. *Ann. Am. Thorac. Soc.* 2018;15:470-8.
130. Nooijen CFJ, Stam HJ, Bergen MP, Bongers-Janssen HMH, Valent L, van Langeveld S, et al. A behavioural intervention increases physical activity in people with subacute spinal cord injury: A randomised trial. *J. Physiother.* 2016;62:35–41.

131. Ang DC, Kaleth AS, Bigatti S, Mazzuca SA, Jensen MP, Hilligoss J, et al. Research to encourage exercise for fibromyalgia (REEF): use of motivational interviewing, outcomes from a randomized-controlled trial. *Clin J Pain*. 2013;29:296–304.
132. Plotnikoff RC, Costigan SA, Karunamuni ND, Lubans DR. Community-based physical activity interventions for treatment of type 2 diabetes: A systematic review with meta-analysis. *Front. Endocrinol*. 2013;4:3.
133. Hale LA, Smith C, Mulligan H, Treharne GJ. “Tell me what you want, what you really really want...”: asking people with multiple sclerosis about enhancing their participation in physical activity. *Disabil. Rehabil*. 2012;34:1887–93.
134. Crowe S, Gilis C. Making patient relevant clinical research a reality. *BMJ*. 2016;23:i6627.
135. Bate P, Robert G. Experience-based design: from redesigning the system around the patient to co-designing services with the patient. *Qual. Saf. Health Care*. 2006;15:307–10.
136. Yardley L, Morrison L, Bradbury K, Muller I. The person-based approach to intervention development: Application to digital health-related behavior change interventions. *J. Med. Internet Res*. 2015;17:e30.
137. Pope C, Mays N. Qualitative Research: Reaching the parts other methods cannot reach: an introduction to qualitative methods in health and health services research. *BMJ*. 1995;311:42–5.
138. Newitt R, Barnett F, Crowe M. Understanding factors that influence participation in physical activity among people with a neuromusculoskeletal condition: a review of qualitative studies. *Disabil. Rehabil*. 2016;38:1–10.
139. Kerstin W, Gabriele B, Richard L. What promotes physical activity after spinal cord injury? An interview study from a patient perspective. *Disabil. Rehabil*. 2006;28:481–8.
140. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): A 32-item checklist for interviews and focus groups. *Int. J. Qual. Heal. Care*. 2007;19:349–57.
141. DiCicco-Bloom B, Crabtree BF. The qualitative research interview. *Med. Educ*. 2006;40:314–21.
142. Kemper H, Ooijendijk W, Stiggelbout M. Consensus over de Nederlandse norm voor gezond bewegen [Consensus on the Dutch norm for healthy exercise]. *Tijdschr. voor Soc. Gezondheidszorg*. 2000;78:180–3.
143. Hoekman R, Hoenderkamp K, van der Poel H, Bakker S, Romijn D, van der Werff H. *Sportaccommodaties in beeld [Sports accommodations in the picture]*. Utrecht, the Netherlands: Mulier Instituut / Arko Sports Media; 2013.
144. Hoekstra F, Roberts L, van Lindert C, Martin Ginis KA, van der Woude LHV, McColl MA. National approaches to promote sports and physical activity in adults with disabilities: examples from the Netherlands and Canada. *Disabil. Rehabil*. 2018;Epub ahead of print.

References

145. Martin Ginis KA, Latimer AE, McKechnie K, Ditor DS, McCartney N, Hicks AL, et al. Using exercise to enhance subjective well-being among people with spinal cord injury: The mediating influences of stress and pain. *Rehabil. Psychol.* 2003;48:157–64.
146. Getz M, Hutzler Y, Vermeer A. The effects of aquatic intervention on perceived physical competence and social acceptance in children with cerebral palsy. *Eur. J. Spec. Needs Educ.* 2007;22:217–28.
147. Lai B, Young HJ, Bickel CS, Motl RW, Rimmer JH. Current Trends in Exercise Intervention Research, Technology, and Behavioral Change Strategies for People With Disabilities: A Scoping Review. *Am. J. Phys. Med. Rehabil.* 2017;96:748–61.
148. Foster C. Guidelines for Health-Enhancing Physical Activity Promotion Programmes [Internet]. Tampere, Finland: 2000. Available from: <http://www.panh.ch/hepaeurope/materials/Guidelines HEPA Europe.pdf>
149. Hildebrandt H, Ooijendijk M, Hopman M. Trendrapport Bewegen en Gezondheid [Trend report on physical activity and health]. Leiden, The Netherlands: TNO, De Brink, Leiden; 2010.
150. Krops LA, Jaarsma EA, Dijkstra PU, Geertzen JHB, Dekker R. Health related quality of life in a dutch rehabilitation population: Reference values and the effect of physical activity. *PLoS One.* 2017;12:e0169169.
151. Jaarsma EA, Dijkstra PU, Geertzen JHB, Dekker R. Barriers to and facilitators of sports participation for people with physical disabilities: A systematic review. *Scand. J. Med. Sci. Sports.* 2014;24:871–81.
152. Krops LA, Folkertsma N, Hols DHJ, Geertzen JHB, Dijkstra PU, Dekker R. Target population's requirements on a community-based intervention for stimulating physical activity in hard-to-reach physically disabled people: an interview study. *Disabil. Rehabil.* 2018; Epub ahead of print.
153. Lustria ML, Cortese J, Noar SM, Glueckauf RL. Computer-tailored health interventions delivered over the web: Review and analysis of key components. *Patient Educ. Couns.* 2009;74:156–73.
154. Bandura A. Social foundations of thought and action : a social cognitive theory. Englewood Cliffs, NJ: Prentice-Hall; 1986.
155. Petty RE, Barden J, Wheeler SC. The Elaboration Likelyhood Model of Persuasion: Developing health promotions for sustained behavioral change. In: *Emerging theories in health promotion practice and research.* San Fransisco: Jossey-Bass; 2009. p. 185–214.
156. Bandura A. Self-efficacy: The exercise of control. New York: W.H. Freeman; 1997.
157. Rogers EM. Diffusion of innovations. New York: Free Press; 2003.
158. Kazdin AE. Behavior modification in applied settings. 6th ed. Long Grove, IL: Waveland Press; 2008.

159. Creer TL. Self-management of chronic illness. In: Handbook of self-regulation. Dan Diego: Academic Press; 2000. p. 601–29.
160. Zimmerman MA. Empowerment theory: Psychological, organizational and community levels of analysis. In: Handbook of community psychology. New York: Kluwer Academic/ Plenum Publishers; 2000. p. 43–62.
161. Latham GP, Locke EA. New developments in and directions for goal-setting research. *Eur. Psychol.* 2007;12:290–300.
162. Heaney CA, Israel BA. Social networks and social support. In: Health behavior and health education: Theory, research and practice. San Fransisco: Jossey-Bass; 2008. p. 189–210.
163. Martins RK, Mcneil DW. Review of Motivational Interviewing in promoting health behaviors. *Clin. Psychol. Rev.* 2009;29:283–93.
164. Kenniscentrum Sport. Beweegcoach [Knowledge center Sport. Activity coach] [Internet]. 2017 [cited 2017 Jul 7];Available from: <https://www.netwerkinbeweging.nl/interventies/overzicht/22530?mode=Mode-all&trefwoorden=beweegcoach&leeftijds categorie=&beoordeling=&thema=&doelgroep=&setting=&sport=>
165. Elbers RG, Rietberg MB, Van Wegen EEH, Verhoef J, Kramer SF, Terwee CB, et al. Self-report fatigue questionnaires in multiple sclerosis, Parkinson's disease and stroke: A systematic review of measurement properties. *Qual. Life Res.* 2012;21:925–44.
166. Nooijen CFJ, Post MWM, Spijkerman DCM, Bergen MP, Stam HJ, Van Den Berg-Emons RJG. Exercise self-efficacy in persons with spinal cord injury: Psychometric properties of the Dutch translation of the exercise self-efficacy scale. *J. Rehabil. Med.* 2013;45:347–50.
167. Post MWM, de Witte LP, Reichrath E, Verdonschot MM, Wijnhuizen GJ, Perenboom RJM. Development and validation of impact-s, an ICF-based questionnaire to measure activities and participation. *J. Rehabil. Med.* 2008;40:620–7.
168. Seves BL, Hoekstra F, Schoenmakers JWA, Hoekstra T, Hettinga FJ, Dekker R, et al. Test-retest reliability and concurrent validity of the Short Questionnaire to Assess Health-enhancing physical activity (SQUASH), adapted for adults with a physical disability or chronic disease. Submitted.
169. Tyson S, Connell L. The psychometric properties and clinical utility of measures of walking and mobility in neurological conditions: a systematic review. *Clin. Rehabil.* 2009;23:1018–33.
170. Collen F, Wade D, Bradshaw C. Mobility after stroke: Reliability of measures of impairment and disability. *Disabil. Rehabil.* 1990;12:6–9.
171. Bellet RN, Adams L, Morris NR. The 6-minute walk test in outpatient cardiac rehabilitation: Validity, reliability and responsiveness-a systematic review. *Physiother.* 2012;98:277–87.

172. Verschuren O, Ketelaar M, De Groot J, Vila Nova F, Takken T. Reproducibility of two functional field exercise tests for children with cerebral palsy who self-propel a manual wheelchair. *Dev. Med. Child Neurol.* 2013;55:185–90.
173. Berg K, Wood-Dauphinee S, Williams JI, Gayton D. Measuring balance in the elderly: Preliminary development of an instrument. *Physiother.* 1989;41:304–11.
174. Blum L, Korner-Bitensky N. Usefulness of the Berg Balance Scale in stroke rehabilitation: a systematic review. *Phys. Ther.* 2008;88:559–66.
175. Horemans HLD, Kooijmans H, Van Den Berg-Emons RJG, Bussmann JBJ. The Activ8 activity monitor: validation of posture and movement classification. Submitted.
176. Ada L, Dean CM, Hall JM, Bampton J, Crompton S. A treadmill and overground walking program improves walking in persons residing in the community after stroke: A placebo-controlled, randomized trial. *Arch. Phys. Med. Rehabil.* 2003;84:1486–91.
177. Horodyska K, Luszczynska A, van den Berg M, Hendriksen M, Roos G, De Bourdeaudhuij I, et al. Good practice characteristics of diet and physical activity interventions and policies: an umbrella review. *BMC Public Health.* 2015;15:19.
178. Jordan JL, Holden MA, Mason EE, Foster NE. Interventions to improve adherence to exercise for chronic musculoskeletal pain in adults. *Cochrane database Syst. Rev.* 2010;CD005956.
179. Hickey AM, Freedson PS. Utility of Consumer Physical Activity Trackers as an Intervention Tool in Cardiovascular Disease Prevention and Treatment. *Prog. Cardiovasc. Dis.* 2016;58:613–9.
180. Finkelstein EA, Haaland BA, Bilger M, Sahasranaman A, Sloan RA, Nang EEK, et al. Effectiveness of activity trackers with and without incentives to increase physical activity (TRIPPA): a randomised controlled trial. *Lancet Diabetes Endocrinol.* 2016;4:983–95.
181. Kohl HW, Craig CL, Lambert EV, Inoue S, Alkandari JR, Leetongin G, et al. The pandemic of physical inactivity: Global action for public health. *Lancet.* 2012;380:294–305.
182. Carroll DD, Courtney-Long EA, Stevens AC, Sloan ML, Lullo C, Visser SN, et al. Vital signs: disability and physical activity--United States, 2009-2012. *Morb. Mortal. Wkly. Rep.* 2014;63:407–13.
183. Mendoza-Vasconez AS, Linke S, Munoz M, Pekmezi D, Ainsworth C, Cano M, et al. Promoting Physical Activity among Underserved Populations. *Curr. Sports Med. Rep.* 2016;15:290–7.
184. Krops LA, Dekker R, Geertzen JHB, Dijkstra PU. Development of an intervention to stimulate physical activity in hard-to-reach physically disabled people and design of a pilot-implementation: an Intervention Mapping approach. *BMJ Open.* 2018;8:e020934.
185. Segura-Ortí E, Martínez-Olmos FJ. Test-Retest Reliability and Minimal Detectable Change Scores for Sit-to-Stand-to-Sit Tests, the Six-Minute Walk Test, the One-Leg Heel-Rise Test, and Handgrip Strength in People Undergoing Hemodialysis. *Phys. Ther.* 2011;91:1244–52.

186. Tyson S, Connell L. The psychometric properties and clinical utility of measures of walking and mobility in neurological conditions: a systematic review. *Clin. Rehabil.* 2009;23:1018–33.
187. Krops LA, Wolthuizen L, Dijkstra PU, Jaarsma EA, Geertzen JH, Dekker R. Reliability of translation of the RAND 36-item health survey in a post-rehabilitation population. *Int J Rehabil Res.* 2017;41:128-137.
188. Rosti-Otajärvi E, Hämäläinen P, Wiksten A, Hakkarainen T, Ruutiainen J. Validity and reliability of the Fatigue Severity Scale in Finnish multiple sclerosis patients. *Brain Behav.* 2017;7:e00743.
189. Greaves CJ, Sheppard KE, Abraham C, Hardeman W, Roden M, Evans PH, et al. Systematic review of reviews of intervention components associated with increased effectiveness in dietary and physical activity interventions. *BMC Public Health.* 2011;11:119.
190. Michie S, Abraham C, Whittington C, McAteer J, Gupta S. Effective techniques in healthy eating and physical activity interventions: a meta-regression. *Heal. Psychol.* 2009;28:690–701.
191. Avery L, Flynn D, Van Wersch A, Sniehotta FF, Trenell MI. Changing physical activity behavior in type 2 diabetes: A systematic review and meta-analysis of behavioral interventions. *Diabetes Care.* 2012;35:2681–9.
192. Visscher TLS, Seidell JC. Time trends (1993-1997) and seasonal variation in body mass index and waist circumference in the Netherlands. *Int. J. Obes. Relat. Metab. Disord.* 2004;28:1309–16.
193. Lang JT, Kassin TO, Devaney LL, Colon-Semenza C, Joseph MF. Test-retest reliability and minimal detectable change for the 10-meter walk test in older adults with Parkinson's disease. *J. Geriatr. Phys. Ther.* 2016;39:165–70.
194. Holland AE, Hill CJ, Rasekaba T, Lee A, Naughton MT, McDonald CF. Updating the Minimal Important Difference for Six-Minute Walk Distance in Patients With Chronic Obstructive Pulmonary Disease. *Arch. Phys. Med. Rehabil.* 2010;91:221–5.
195. Gervasoni E, Jonsdottir J, Montesano A, Cattaneo D. Minimal Clinically Important Difference of Berg Balance Scale in People With Multiple Sclerosis. *Arch. Phys. Med. Rehabil.* 2017;98:337–340.e2.
196. Wyrwich KW, Tierney WM, Babu AN, Kroenke K, Wolinsky FD. A comparison of clinically important differences in health-related quality of life for patients with chronic lung disease, asthma, or heart disease. *Heal. Serv Res.* 2005;40:577–91.
197. CentraalBureauvoor deStatistiek.Laagenlangduriglaaginkomen;huishoudenskenmerken en regio [Statistics Netherlands. Low- and longterm low incomes; household characteristics and region]. [Internet]. 2017 [cited 2018 Jan 23];Available from: <https://www.volksgezondheidenzorg.info/onderwerp/sociaaleconomische-status/regionaal-internationaal/regionaal#node-huishoudens-met-laag-inkomen-gemeente>

References

198. Van Zon SKR, Snieder H, Bültmann U, Reijneveld SA. The interaction of socioeconomic position and type 2 diabetes mellitus family history: A cross-sectional analysis of the Lifelines Cohort and Biobank Study. *BMJ Open*. 2017;7:e015275.
199. Giles-Corti B, Donovan RJ. Socioeconomic status differences in recreational physical activity levels and real and perceived access to a supportive physical environment. *Prev. Med*. 2002;35:601–11.
200. Sattelmair J, Pertman J, Ding EL, Kohl HW, Haskell W, Lee IM. Dose response between physical activity and risk of coronary heart disease: A meta-analysis. *Circulation*. 2011;124:789–95.
201. Kahn EB, Ramsey LT, Brownson RC, Heath GW, Howze EH, Powell KE, et al. The Effectiveness of Interventions to Increase Physical Activity: A Systematic Review. *Am. J. Prev. Med*. 2010;22:73–107.
202. Yancey AK, McCarthy WJ, Harrison GG, Wong WK, Siegel JM, Leslie J. Challenges in Improving Fitness: Results of a Community-Based, Randomized, Controlled Lifestyle Change Intervention. *J. Women's Heal*. 2006;15:412–29.
203. Cleland CL, Tully MA, Kee F, Cupples ME. The effectiveness of physical activity interventions in socio-economically disadvantaged communities: A systematic review. *Prev. Med*. 2012;54:371–80.
204. Lai B, Kim Y, Wilroy J, Bickel CS, Rimmer JH, Motl RW. Sustainability of exercise intervention outcomes among people with disabilities: a secondary review. *Disabil. Rehabil*. 2018;Epub ahead of print.
205. Bartholomew LK, Parcel GS, Kok G. Intervention mapping: a process for developing theory- and evidence-based health education programs. *Health Educ. Behav*. 1998;25:545–63.
206. Jansen MWJ, van Oers HAM, Kok G, de Vries NK. Public health: Disconnections between policy, practice and research. *Heal. Res. Policy Syst*. 2010;8-37.
207. Briss PA, Zaza S, Pappaioanou M, Fielding J, Wright-De Agüero L, Truman BI, et al. Developing an evidence-based Guide to Community Preventive Services - Methods. *Am. J. Prev. Med*. 2000;18:35–43.
208. Green LW, Glasgow RE. Evaluating the relevance, generalization, and applicability of research: Issues in external validation and translation methodology. *Eval. Heal. Prof*. 2006;29:126–53.
209. Goldman N. Social inequalities in health: Disentangling the underlying mechanisms. *Ann. N. Y. Acad. Sci*. 2001;954:118–39.
210. Rollnick S, Miller WR. What is Motivational Interviewing? *Behav. Cogn. Psychother*. 1995;23:325–34.
211. United Nations. Convention on the rights of persons with disabilities [Internet]. 2006. Available from: <https://www.un.org/development/desa/disabilities/convention-on-the-rights-of-persons-with-disabilities.html>

212. Ministerie van Volksgezondheid Welzijn en Sport. Landelijke nota gezondheidsbeleid: Gezondheid Dichtbij [Ministry of Health, Welfare and Sport. National health policy document: Health nearby]. 2011.
213. Schippers El, van Rijn MJ. Kamerbrief over landelijke nota gezondheidsbeleid 2016-2019 [Letter to the parliament on national health policy document 2016-2019]. The Hague, The Netherlands: 2015.
214. Rijksoverheid. Alles is gezondheid. Het nationaal programma preventie 2014-2016 [National government. Everything is health. National program on prevention 2014-2016]. 2013.
215. Schippers El. Kamerbrief over gehandicaptensportbeleid [Letter to the parliament on disability sports policy]. The Hague, The Netherlands: 2015.

