

University of Groningen

## The crystal structure of a hyperthermoactive exopolygalacturonase from *Thermotoga maritima* reveals a unique tetramer

Pijning, Tjaard; van Pouderooyen, Gertie; Kluskens, Leon; van der Oost, John; Dijkstra, Bauke W.

*Published in:*  
FEBS Letters

*DOI:*  
[10.1016/j.febslet.2009.10.047](https://doi.org/10.1016/j.febslet.2009.10.047)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2009

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Pijning, T., van Pouderooyen, G., Kluskens, L., van der Oost, J., & Dijkstra, B. W. (2009). The crystal structure of a hyperthermoactive exopolygalacturonase from *Thermotoga maritima* reveals a unique tetramer. *FEBS Letters*, 583(22), 3665-3670. <https://doi.org/10.1016/j.febslet.2009.10.047>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Supplementary data




Fig. S1. Sequence alignment of polygalacturonases, figure prepared by ESPrift. *Tma* = *Thermotoga maritima*, *Yen* = *Yersinia enterocolitica* (β-helical domain), *Eca* = *Erwinia carotovora*, *Ani* = *Aspergillus niger*, *Spu* = *Sterum purpureum*, *Clu* = *Colletotrichum lupine*, *Aac* = *Aspergillus aculeatus*, *Fmo* = *Fusarium moniliforme*. Fully conserved residues are boxed in red; the three catalytic aspartates are marked with ▲. Secondary structure elements are coded as follows: α = α-helix, β = β-strand, η = 310-helix, T = hydrogen bonded turn. Turns and loops lining the active site are indicated with their coil number and turn type.