

University of Groningen

Flavoprotein oxidases

Dijkman, Willem P.; de Gonzalo, Gonzalo; Mattevi, Andrea; Fraaije, Marco W.

Published in:
Applied Microbiology and Biotechnology

DOI:
[10.1007/s00253-013-4925-7](https://doi.org/10.1007/s00253-013-4925-7)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dijkman, W. P., de Gonzalo, G., Mattevi, A., & Fraaije, M. W. (2013). Flavoprotein oxidases: classification and applications. *Applied Microbiology and Biotechnology*, 97(12), 5177-5188.
<https://doi.org/10.1007/s00253-013-4925-7>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Flavoprotein oxidases: classification and applications

Willem P. Dijkman · Gonzalo de Gonzalo ·
Andrea Mattevi · Marco W. Fraaije

Received: 13 February 2013 / Revised: 10 April 2013 / Accepted: 11 April 2013 / Published online: 3 May 2013
© Springer-Verlag Berlin Heidelberg 2013

Abstract This review provides an overview of oxidases that utilise a flavin cofactor for catalysis. This class of oxidative flavoenzymes has shown to harbour a large number of biotechnologically interesting enzymes. Applications range from their use as biocatalysts for the synthesis of pharmaceutical compounds to the integration in biosensors. Through the recent developments in genome sequencing, the number of newly discovered oxidases is steadily growing. Recent progress in the field of flavoprotein oxidase discovery and the obtained biochemical knowledge on these enzymes are reviewed. Except for a structure-based classification of known flavoprotein oxidases, also their potential in recent biotechnological applications is discussed.

Keywords Flavoproteins · Oxidases · Biocatalysis · Biosensors · Flavin · Oxygen

Introduction

Selective oxidations are often key in synthetic routes towards valuable chemicals. Chemical oxidation methods typically require harsh conditions and polluting reagents and often suffer from poor chemoselectivity and enantioselectivity. Nature, on the other hand, has come up with a versatile set of enzymes performing a wide variety of selective oxidation

reactions. Known oxidative enzyme classes are the oxidases, oxygenases and dehydrogenases. Oxygenases require reducing equivalents [e.g. NAD(P)H] and molecular oxygen for activity, while dehydrogenases utilise organic coenzymes (for example pyrroloquinoline quinone, quinones or NAD⁺) as electron acceptors. The coenzyme dependence of these enzymes can compromise cost-effective biotechnological applications. Instead of (often expensive) coenzymes, oxidases merely require molecular oxygen as oxidant (electron acceptor) for catalysis. This feature makes them valuable enzymes for industrial applications that range from oxidase-based biosensors to their application as biocatalysts in the synthesis of valuable chemicals. For an organism, however, oxidases are much less favourable. By the direct reduction of molecular oxygen, electrons are lost and cannot be used in subsequent metabolic events, as in the case of dehydrogenases. Another physiological relevant issue concerning oxidases is their inherent reactivity to produce reduced dioxygen species. While some oxidases are capable to reduce dioxygen to harmless water, most oxidases generate instead hydrogen peroxide as by-product, and even in some cases, the even more toxic superoxide O₂⁻ is formed. The features above provide logic for why oxidases are relatively rare enzymes.

Most oxidases rely on a tightly bound cofactor for their activity, and only a few examples of cofactor independent oxidases have been described (Fetzner and Steiner 2010). This is due to the fact that amino acids are very poor in mediating redox reactions. To equip enzymes with oxidising power, nature has evolved several different redox cofactors. For oxidases, there is a bias towards two types of cofactors resulting in two main oxidase families. One family utilises copper in mono- and trinuclear centers or a copper atom combined with a quinone cofactor. The reader is referred to many reviews on this topic (Guengerich 2012; Ridge et al. 2008). This review will focus on the other major oxidase family: the flavin-containing oxidases.

W. P. Dijkman · G. de Gonzalo · M. W. Fraaije (✉)
Molecular Enzymology Group, Groningen Biomolecular Sciences
and Biotechnology Institute, University of Groningen,
Nijenborgh 4,
9747 AG Groningen, The Netherlands
e-mail: m.w.fraaije@rug.nl
URL: www.rug.nl/staff/m.w.fraaije/research

A. Mattevi
Department of Biology and Biotechnology,
University of Pavia, Via Ferrata 1,
27100 Pavia, Italy

flavoprotein oxidases. Based on sequence homology and the available structural information, a comprehensive classification is proposed (Table 1). The distinct flavoprotein oxidase families will be described in more detail in the next paragraphs, while the last paragraphs will describe several recent advances on applications of flavoprotein oxidases.

The GMC-type oxidase family

The glucose–methanol–choline oxidoreductase (GMC) flavoprotein family contains several well-known oxidases. The protein sequence of each member of this family entails a conserved N-terminal FAD binding domain, called GMC_oxred_N (Pfam00732) in the Pfam database (Finn et al. 2010). This domain includes the typical GxGxxG/A sequence motif, which is indicative for the Rossmann fold that is involved in binding the ADP moiety of the FAD. Less conserved is the C-terminal region, which forms the substrate binding domain (Kiess et al. 1998). It contains only one generally conserved residue, an active-site histidine, which can assist in substrate oxidation and FAD reoxidation by molecular oxygen. Mutagenesis studies on glucose oxidase has revealed that protonation of this histidine (His516 in GO) is essential for catalysing dioxygen reduction (Roth and Klinman 2003). Nevertheless, exceptions to this rule have already been discovered. For instance, in choline oxidase, the function of the histidine has been taken over by the positively charged amine moiety of the choline substrate molecule (Gadda 2012).

Known GMC-type oxidases act on primary and secondary alcohols, forming the corresponding aldehydes or ketones. Examples are glucose oxidase (EC 1.1.3.4; Fig. 2a), cholesterol oxidase (EC 1.1.3.6), pyranose oxidase (EC 1.1.3.10), methanol oxidase (EC 1.1.3.13) and aryl alcohol oxidase (EC 1.1.3.7). Choline oxidase (EC 1.1.3.17) is a notable exception as it is able to perform two sequential oxidations on choline, yielding the carboxylic acid, trimethylglycine, as shown in Fig. 3 (Fan et al. 2004).

Several oxidases were found to contain a covalently linked FAD cofactor (choline oxidase and pyranose oxidase), where the flavin is attached via an 8α -N³-histidyl bond. Nevertheless, the majority of GMC-type oxidases contain a dissociable FAD cofactor that is tightly bound to the protein. Mechanistic details have been established for a number of GMC-type oxidases, including glucose oxidase, pyranose oxidase, aryl alcohol oxidase and choline oxidase. The mechanism by which GMC-type oxidases oxidise their substrates involves a direct hydride transfer from the substrate to the N5 atom of the FAD cofactor. This reaction is facilitated by proton abstraction, typically promoted by a histidine acting as a general base.

In addition to oxidases, the GMC flavoprotein family also contains other types of flavoenzymes. A number of flavoprotein dehydrogenases (for example glucose dehydrogenase and cellobiose dehydrogenase) and even non-redox flavoproteins (for example hydroxynitrile lyase) are members of this superfamily. In contrast to the sequence-related oxidases, these enzymes are ineffective in utilising molecular oxygen as electron acceptor. At present, there is no sequence motif known that can discriminate between GMC-type oxidases and other GMC-type enzymes. The same holds true for most of the flavoprotein oxidase families below. More knowledge on the precise mechanism by which flavoproteins can tune their oxygen reactivity may change this situation and would perhaps enable more reliable annotation of flavoprotein-encoding genes into oxidases, facilitating effective genome mining for novel flavoprotein oxidases.

The VAO-type oxidase family

The second flavoprotein family, which includes oxidases, is the vanillyl alcohol oxidase (VAO) flavoprotein family, which was named after a fungal oxidase, VAO (EC 1.1.3.38; Fraaije et al. 1998; Leferink et al. 2008; Mattevi

Table 1 Classification of the major flavoprotein oxidase families

Flavoprotein oxidase family	GMC	VAO	AO	SO	ACO	HAO
Cofactor type	FAD	FAD	FAD	FAD	FAD	FMN
Cofactor binding mode:						
Noncovalent	+	+	+	+	+	+
Covalent	+	+	+	–	–	–
Typical substrates	Alcohols	Alcohols Amines	Amines	Thiols	Acyl CoA	Hydroxyacids
Oxidase prototype (PDB)	Glucose oxidase (1CF3)	Vanillyl alcohol oxidase (1VAO)	D-Amino acid oxidase (1KIF)	Erv2p sulfhydryl Oxidase (1JRA)	Acyl CoA oxidase (1IS2)	Glycolate oxidase (1GOX)




Fig. 2 Structures of representative members of the different flavin-dependent oxidases with the FAD or FMN cofactor highlighted in gold. **a** Glucose oxidase (pdb 1CF3). **b** Vanillyl alcohol oxidase (pdb

1VAO). **c** D-amino acid oxidase (pdb 1KIF). **d** Sulfhydryl oxidase Ery2p (pdb 1JRA). **e** acyl CoA oxidase (pdb 1FS2). **f** Glycolate oxidase (pdb 1GOX)

et al. 1997). Similar to GMC-type flavoproteins, all VAO-type flavoproteins contain a distinct FAD binding domain (FAD_binding_4 domain, Pfam01565) in the N-terminal half of the protein (Fig. 2b). This domain includes the so-called PP loop where the pyrophosphate moiety of the FAD interacts with the protein (Mattevi et al. 1997). The vanillyl alcohol oxidase crystal structure revealed for the first time how a FAD cofactor is covalently tethered to a protein via a histidine residue. Indeed, the VAO flavoprotein family is relatively rich in covalent flavoproteins (estimated 25 % contain a histidyl-FAD). It has been established that, by covalent attachment of the FAD cofactor to a histidine, the redox potential of the cofactor is increased to a large extent

(Fraaije et al. 1999). As a result, the number of feasible electron acceptors is limited, leaving molecular oxygen as one of the few candidates. This explains why many VAO flavoproteins act as oxidases. Recently, it has been discovered that several VAO-type oxidases even contain a bicovalently linked FAD cofactor. In these cases, in addition to the histidyl bond, the FAD is also tethered at the 6' position to a cysteine (Heuts et al. 2009). Except for an effect on the redox potential, it is hypothesised that the second covalent bond allows the protein to create a very open active site, while retaining the FAD cofactor in position for catalysis through the bicovalent anchoring. This is in line with the observation that the described bicovalent flavoprotein oxidases all accept rather bulky substrates (Heuts et al. 2007a).




Fig. 3 Choline oxidase-catalysed sequential oxidation of choline into trimethylglycine

For VAO-type oxidases, some details on the oxygen reactivity have been elucidated. For the VAO-type cholesterol oxidase and alditol oxidase (EC 1.1.3.41) specific oxygen channels have been described, guiding dioxygen to the FAD (Baron et al. 2009; Coulombe et al. 2001). In alditol oxidase, all proposed channels are situated in the C-terminal part of the protein and lead to one oxygen entry site

in the active site. Based on this finding, a mutant of an L-galactonolactone dehydrogenase (EC 1.3.2.3) has been engineered in which such an oxygen entry in the active site was created. The mutation indeed resulted in the creation of an oxidase (Leferink et al. 2009).

In the last decade, a large number of novel VAO-type oxidases have been discovered and studied (Leferink et al. 2008; Winter and Fraaije 2012). In addition to alcohol oxidations, leading in most cases to aldehydes or ketones, many other bonds can be oxidised. Vanillyl alcohol oxidase itself was shown to perform alcohol and amine oxidations, hydroxylations of alkyl-phenols and ether bond cleavage (Fraaije et al. 1995). Another remarkable example of a VAO-type oxidase is reticuline oxidase, also known as berberine bridge enzyme (EC 1.21.3.3). This enzyme is able to form a C–C bond, creating the intramolecular bond between a methyl group and an aromatic carbon in the conversion from (*S*)-reticuline to (*S*)-scoulerine, a plant alkaloid (Winkler et al. 2008). Elucidation of the crystal structure of this plant enzyme has revealed that the reaction is triggered by hydride transfer from the substrate to the flavin cofactor. VAO-type oxidases that can catalyse an oxidative C–C bond formation are not restricted to plants. Prosolanapyrone-II oxidase (EC 1.1.3.42) from the fungus *Alternaria solani* performs an intramolecular Diels–Alder reaction, a [4+2] cycloaddition in which a conjugated diene reacts with a carbon–carbon double bond (Kasahara et al. 2010). In addition, other microbial C–C and C–O forming VAO-type oxidases have been described in recent literature (Winter and Fraaije 2012).

The amine oxidase family

The amine oxidase (AO)-type oxidases belong to another distinct structural FAD-containing flavoprotein family. As the name suggests, most enzymes are solely active on amines. Prototype oxidases for this family of flavoprotein oxidases are microbial amino acid oxidases (Fig. 2c) and eukaryotic monoamine oxidases. Beside C–N oxidising oxidases, a few alcohol and thioether oxidases have been found that belong to this family. This shows that AO-type oxidases cover a wide range of different reactions. The individual enzymes, however, can in most cases perform only one type of reaction.

All members of this family (Pfam01593 and Pfam01266) share a similar domain in the N-terminal half of the protein, responsible for FAD binding. Covalent binding through a histidyl or cysteinyl bond on the 8 α position of FAD is frequently observed. Most AO-type oxidases act on a carbon–nitrogen bond of a primary or secondary amine. The catalytic cycle of these oxidases starts with proton abstraction from the amine or ammonium group by an active site base like histidine (Trickey et al. 1999) or a catalytic dyad of

histidine and tyrosine (Leys et al. 2003). Hydride transfer to the enzyme-bound FAD results in the formation of an imine product intermediate, which can spontaneously hydrolyse to an aldehyde and amine (Fitzpatrick 2010; Trickey et al. 1999).

Among the known AO-type oxidases, also thioether oxidases have been identified: prenylcysteine lyase (EC 1.8.3.5) and farnesylcysteine lyase (EC 1.8.3.6). The proposed catalytic mechanism for prenylcysteine oxidase is quite distinct from other FAD-catalysed oxidations. The C–S bond is not activated by an active site base; the first step is hydride transfer to FAD. The resulting thiocarbenium is hydrolysed to farnesal, an aldehyde and cysteine (Fig. 4) (Digits et al. 2002). The overall products of the reaction, therefore, resemble the ones obtained upon secondary amine oxidation.

Another unusual AO-type oxidase is *nikD*, which is involved in the biosynthesis of the antibiotic nikkomycin. This oxidase produces the aromatic compound picolinate by a 4-electron oxidation of Δ -piperidine-2-carboxylate (Fig. 5). In the proposed mechanism, the reaction is initiated with oxidation of a carbon–nitrogen bond. After tautomerisation and isomerisation of the formed imine, the same carbon–nitrogen bond is oxidised again resulting in the final, aromatic product (Bruckner and Jorns 2009).

In contrast to all other members of this flavoprotein oxidase family, including monomeric sarcosine oxidase (EC 1.5.3.1), heterotetrameric sarcosine oxidase does not rely only on FAD as cofactor. The α -subunit of this tetrameric enzyme binds NAD^+ , but has no described catalytic function. The β -subunit resembles monomeric sarcosine oxidase, but contains both FAD and covalently bound FMN. In this oxidase subunit, sarcosine is oxidised to the corresponding imine by FAD. With the help of the γ and δ subunits, heterotetrameric sarcosine oxidase can interact with 5,10- $\text{CH}_2\text{-H}_4$ folate synthase, forming a channel, which prevents hydrolysis of the imine. The reaction of the imine product reacts with tetrahydrofolate resulting in the formation of 5,10- CH_2 -tetrahydrofolate and glycine (Chen et al. 2006).

The sulfhydryl oxidase family

A number of FAD-containing oxidases acting on thiols have been discovered in the last decade. The most important role of these so-called sulfhydryl oxidases is the formation of disulphide bonds in proteins by oxidising cysteine residues. Crystal structures of several sulfhydryl oxidases have been solved, including human, viral and yeast variants (Fig. 2d; Guo et al. 2012; Gross et al. 2004). Based on this, two subfamilies can be identified: *Erv*-like (Pfam04777) and *Ero*-like (Pfam04137) sulfhydryl oxidases. The first subfamily is named after *Erv1* (EC 1.8.3.2), a protein that was discovered

Another recently discovered ACO-type oxidase is the fungal nitroalkane oxidase (EC 1.7.3.1). This FAD-containing oxidase oxidises primary and secondary nitroalkanes to the corresponding aldehydes and ketones with the release of nitrite (Fig. 6). The first step, abstraction of the α -proton from the nitroalkane, is similar to the acyl-CoA oxidase mechanism. The carbanion obtained forms a covalent adduct with the N5 of the flavin, which generates a cationic flavin imine. Attack by water leads to the reduced flavin and the final product, the aldehyde or ketone (Tormos et al. 2010). Reoxidation of the reduced flavin cofactor by molecular oxygen completes the catalytic cycle.

The 2-hydroxyacid oxidase family

A number of flavoprotein oxidases use FMN as prosthetic group. The majority of these oxidases belong to one distinct structural flavoprotein family. Many members of this oxidase family oxidise aromatic or aliphatic 2-hydroxy acids (grouped in EC 1.1.3.5), forming the respective 2-oxoacids (Cunane et al. 2005). Known examples of this oxidase family are glycolate oxidase and L-lactate oxidase. The structure of several members of this family has been solved, providing insights into the fold and the catalytically important residues. The proteins have a $\beta 8/\alpha 8$ TIM barrel structure (Fig. 2f; Umena et al. 2006). A conserved arginine in the active site typically accommodates the carboxyl group of the substrate. The catalytic mechanism of the reductive half reaction for 2-hydroxyacid oxidase (HAO)-type oxidases is still under debate. Studies seem to support both the formation of a carbanion and a hydride transfer mechanism (Furuichi et al. 2008). In both cases, however, a histidine (His265 in lactate oxidase) is involved, abstracting the C α -proton in the case of the carbanion mechanism and the proton from the C α -hydroxyl group in the hydride transfer mechanism.

The crystal structure of a FMN-containing nitroalkane oxidase has recently been elucidated (Li et al. 2011). It revealed no structural resemblance with the above-mentioned ACO-type oxidases but rather to the HAO-type flavoprotein oxidase family. This shows that members of the HAO family can perform other reactions than the oxidation

of 2-hydroxyacids. The active site of this FMN-containing TIM barrel protein also contains a histidine residue that was found to be crucial for catalysis.

Unique flavoprotein oxidases

Several flavoprotein oxidases have been identified for which no or only a few homologs with other activities have been described. These individual oxidases are discussed below. Pyruvate oxidase is an exceptional flavin-containing oxidase as it contains two cofactors, the vitamin B1 and B2 derivatives, thiamine diphosphate (ThDP) and FAD, respectively. The combination of ThDP and FAD in one enzyme is rare, but not restricted to pyruvate oxidase. Acetolactate synthase, also acting on pyruvate, is another example in which both cofactors are bound (Choi et al. 2007). The crystal structure of pyruvate oxidase from *Lactobacillus plantarum* (Muller et al. 1994) has revealed that each monomer of the homotetramer consist of three domains. The FAD cofactor is not buried within the enzyme but is located at the surface between the N-terminal α domain and the middle β domain. ThDP is coordinated by a magnesium ion and located at the C-terminal γ domain. The α and γ domains have a similar topology, whereas the β domain differs (Arjunan et al. 1996). Pyruvate oxidase catalyses the decarboxylation of pyruvate to acetyl phosphate and carbon dioxide, consuming oxygen and yielding hydrogen peroxide. The proposed catalytic mechanism involves both cofactors, ThDP and FAD. The activated C2 carbanion of the thiazolium ring of ThDP attacks pyruvate, leading to decarboxylation. FAD subsequently oxidises the ThDP-substrate adduct. The attack by phosphate releases acetyl phosphate from ThDP, and FADH₂ is oxidised by dioxygen to complete the catalytic cycle (Tittmann et al. 2005).

Pyridoxal 5'-phosphate oxidase (PNPO, EC 1.4.3.5) is an essential enzyme in the metabolism of vitamin B6. The enzyme oxidises the amine group of pyridoxamine 5'-phosphate to an aldehyde (Di Salvo et al. 2011), a reaction similar to those catalysed by the AO-type oxidases. The enzyme is also active on pyridoxine 5'-phosphate in which the alcohol is oxidised to the aldehyde. In addition, activity on imine and ester moieties has been described (Kazarinoff and McCormick 1973; Kazarinoff and McCormick 1975). PNPO contains a FMN as flavin cofactor and displays no structural resemblance with any of the above discussed flavoprotein oxidases. The crystal structures of human, yeast and two bacterial homologs have been solved. All PNPOs are dimers, with each monomer composed of two domains. The N-terminal domain (Pfam01243) forms a six stranded β -barrel with antiparallel strands. The smaller C-terminal domain (Pfam 10590) adopts a helix-loop-helix conformation. FMN binds in the β -barrel of the protein with the




Fig. 6 General scheme of the oxidation of primary nitroalkanes by nitroalkane oxidase in order to obtain aldehydes

phosphate moiety near the N-terminus (Biswal et al. 2005). Mechanistic studies have revealed that, as for many other flavoprotein oxidases, oxidation of the substrate is achieved by a direct hydride transfer from the substrate to the N5 of the FMN cofactor (Di Salvo et al. 2011).

Another FAD-containing oxidase that catalyses only one type of reaction is NADPH oxidase (NOX). This enzyme is implicated in a large number of physiological processes in humans. A crucial catalytic property is the ability to generate superoxide as reduced oxygen species. The enzyme is composed of multiple subunits and is membrane bound. Although the crystal structure of NOX has not yet been elucidated, a structure of the subunit gp91^{phox} has been proposed based on sequence alignments and homology models. This shows that the protein contains an N-terminal membrane domain composed of six α -helices. The C-terminal domain is exposed in the cytosol and contains both the NADPH and the FAD binding sites (Debeurme et al. 2010). NOXs have also been described in prokaryotes, being both FAD- and FMN-dependent enzymes depending on the microbial source. These enzymes can be produced as recombinant enzymes, being valuable biocatalysts for regeneration of the NAD(P)⁺ cofactor.


Xanthine oxidase (XO, EC 1.17.3.2) represents yet another flavin-containing oxidase. The oxidase does not only rely on FAD for activity but also contains two non-identical [2Fe–2S] clusters and a molybdenum cofactor. In fact, the oxidation of the organic substrate occurs at the molybdopterin. The [2Fe–2S] clusters assist in catalysis by facilitating electron transfer from the molybdenum site to the flavin cofactor (Pauff et al. 2007). The FAD cofactor is merely used to transfer the electrons to a suitable electron acceptor. This also explains the atypical oxidations that are catalysed by XO: It can catalyse hydroxylations of a large number of compounds, including pyrimidines, purines and pterins. In contrast with typical oxygenases, the inserted oxygen is derived from water. XO is an excellent example of the thin line between oxidases and dehydrogenases. The oxidase can efficiently use dioxygen as electron acceptor, but is unable to reduce NAD⁺. This is in sharp contrast with xanthine dehydrogenase (XDH) for which the opposite is true. Yet, both XO and XDH are essentially the same protein (Nishino et al. 2008). XDH can reversibly be interconverted to XO when two disulphide bonds are formed (Nishino et al. 2005). The disulphide bond formation causes a conformational change which disrupts the cavity needed for NAD⁺ to reach FAD. Irreversible formation of XO from XDH happens upon proteolytic cleavage of the linker between the FAD and the molybdenum binding domain, again changing the environment of the NAD⁺ binding site, leaving the protein only active towards dioxygen. Another special feature of XO is that, depending on the conditions, it produces hydrogen peroxide or superoxide as reduced dioxygen species.

Flavoprotein oxidases in biotechnology

Flavoprotein oxidases have been widely employed in biocatalytic processes. In most cases, the targeted reactions involved the selective oxidation of alcohols or amines. In the last decade, amine-oxidising enzymes have become very popular as biocatalysts. Both monoamine oxidases and amino acid oxidases have been extensively applied in the selective oxidation of amines and amino acids under mild reaction conditions. For instance, a D-amino acid oxidase from yeast has been employed in the industrial preparation of 7-aminocephalosporanic acid (Pillone and Pollegioni 2002), and an engineered bacterial glycine oxidase could be engineered to convert *N*-(phosphonomethyl)glycine (glyphosate; Pedotti et al. 2009). Another development is the use of monoamine oxidases and amino acid oxidases in chemo-enzymatic deracemisation of racemic amines. This concept relies on the ability of the oxidases to oxidise only one enantiomer into the corresponding imine or imino acid, which is chemically (nonselectively) reduced back into the amine. By performing this cycle of enzymatic oxidation and chemical reduction in one pot (cyclic deracemisation), a 100% theoretical yield of an optically pure amine is feasible. For examples of this elegant approach for deracemisation of amines, the reader is referred to a recent review of Turner (2011). Amine-oxidising oxidases have also been combined with other enzymes for the preparation of enantiopure amines (Patel 2011). For example, in 2011, enantiopure (*S*)-amino-3-[3-{6-(2methylphenyl)}pyridyl]-propionic acid was synthesised in 72% yield from the racemic amino acid by combining the (*R*)-amino acid oxidase from *Trigonopsis variabilis* with (*S*)-aminotransferase from *Sporosarcina urea* (Patel et al. 2007). The racemic amino acid mixture was converted by the oxidase into the (*S*)-enantiomer and a keto acid, of which the latter was converted by the aminotransferase into the (*S*)-amino acid. An (*S*)-aminotransferase from *Burkholderia* sp. was also employed with a global yield of 85% (Fig. 7). The process was scaled up to 100 L. An (*S*)-amino acid oxidase has been used for production of 2-amino-3(7-methyl-1*H*-indazol-5-yl)-propionic acid, a key intermediate for the preparation of anti-migraine drugs. Enantiopure (*R*)-amino acid was prepared with 79% global yield by the selective oxidation with the (*S*)-amino acid oxidase from *Proteus mirabilis* followed by the transamination of the keto acid intermediate by a commercially available (*R*)-transaminase (Hanson et al. 2008).

One recent example of a flavoprotein oxidase that can be used for synthetic chemistry is the plant flavoprotein oxidase, reticuline oxidase. This VAO-type oxidase enzyme catalyses the biotransformation of (*S*)-reticuline into (*S*)-scoulerine through the formation of an intramolecular C–C bond at the expense of molecular oxygen. Scoulerine and related compounds often exhibit interesting biological

Fig. 7 Enzymatic synthesis of (*S*)-amino-3-[3-{6-(2-methylphenyl)}pyridyl]-propionic acid by cyclic deracemisation using two biocatalysts



properties (Schrittwiesser et al. 2011b). Therefore, reticuline oxidase has been thoroughly studied for its use in the synthesis of such alkaloids. The flavoenzyme has been shown to be a potent biocatalyst for performing oxidative coupling reactions to form a range of benzyloisoquinolines derivatives with high selectivity. Initial experiments showed that catalase was required in the reaction medium, as the formation of hydrogen peroxide during the oxidative process led to inactivation of the oxidase. Nevertheless, the enzyme is able to maintain its activity in a broad range of pH values (8–11) and temperatures (30–60 °C). As most of the reticuline oxidase substrates are poorly soluble in water, the effect of different organic solvents and ionic liquids on the biocatalyst properties was studied. This revealed that the oxidase is highly tolerant towards solvents. It enabled working at solvent concentrations of 70% *v/v* and substrate concentrations of 20 g/L (Resch et al. 2011). Thus, a set of racemic 1-substituted tetrahydroisoquinolines was synthesised starting from the corresponding *N*-phenylmethanamines and phenylacetic acid derivatives and subjected to reticuline oxidase-catalysed resolutions (Schrittwiesser et al. 2011a). These oxidase-mediated conversions yielded optically pure products. Oxidation of (*S*)-reticuline catalyzed by wild-type reticuline oxidase occurs with total regioselectivity, leading to the formation of the 9-hydroxy functionalised regioisomer (*S*)-scoulerine, while the 11-hydroxy regioisomer (*S*)-coreximine was not formed (Fig. 8). The reticuline oxidase E417Q mutant was able to produce a 30% of the 11-hydroxy regioisomer, at the expense of a huge decrease in the reaction rate for the major regioisomer, due to the substrate hydroxyl group deprotonation and phenyl moiety activation induced by the residue Glu417. As some of the 11-hydroxy functionalised


products present biological activity, such as (*S*)-isocoreximine, (*S*)-corytenchine and phellodendrine, reticuline oxidase and its mutant were employed in the oxidative coupling of reticuline analogues, but only traces of the non-expected regioisomers were found. Only substrate modification, by introducing fluoro substituents, led to the synthesis of the 11-hydroxy compounds as sole products with good yields and excellent selectivities (Resch et al. 2012).

Oxidases have also attractive features for use in biosensors. A well-known example of a flavoprotein oxidase that has been used for biosensor development is glucose oxidase (Cash and Clark 2010), which has been extensively applied to monitor the blood glucose levels in diabetics due to its catalytic ability to oxidise glucose. The generated electrons can be used in a direct or indirect manner for readout of a sensor. Another example of an oxidase that can be used for diagnostic purposes is fructosyl amino acid oxidase (EC 1.5.3.x) which oxidises the C–N bond of an amino acid–carbohydrate adduct, resulting in formation of the respective glucosone and amino acid. The oxidase is of interest for detecting glycated proteins, a marker for hyperglycemia in diabetes patients (Qian et al. 2013). Furthermore, *D*-amino acid oxidases are widely used for the quantification of *D*-amino acids in biological samples by virtue of their strict stereospecificity (Netto et al. 2013).

Concluding remarks

Oxidases are widespread among a large number of structural flavoprotein families. This shows that nature has found many ways to exploit the ability of the reduced flavin

Fig. 8 Reticuline oxidase-catalysed oxidative C–C coupling leading to the 9-hydroxy (major) or 11-hydroxy (minor) stereoisomer



cofactor to use molecular oxygen as oxidant. Interestingly, depending on the structural fold, certain oxidation reactions are preferred. For example, for GMC-type oxidases, there is a strong bias towards alcohol oxidation while members of the sulfhydryl oxidase (SO) family exclusively act on thiols. Yet, oxidases from different structural families can have similar activities. A notable example is cholesterol oxidase for which two structurally distinct representatives have been discovered, one belonging to the VAO oxidase family and the other is a GMC-type oxidase. Another interesting case of shared substrate specificity among two flavoprotein oxidase families has been found for 6-hydroxynicotine oxidases. For each enantiomer of this compound, a specific and structurally distinct oxidase has been found: (*S*)- and (*R*)-6-hydroxynicotine oxidase (respectively EC 1.5.3.5 and EC 1.5.3.6). These enzymes oxidise the pyrrolidine ring in the substrate to form the enamine 6-hydroxy-*N*-methylmyosmine. Hydration leads to the final product (*R*)-6-hydroxy-pseudooxynicotine (Decker and Dai 1967). Interestingly, the other enantiomer (*S*)-6-hydroxynicotine is oxidised by an AO-type oxidase.

The flavoprotein oxidases that encompass the broadest range of oxidation activities are the VAO-type oxidases. Reactions of these oxidases include alcohol oxidations, amine oxidation, oxidative ether-bond cleavage, hydroxylations and oxidative C–C and C–O bond formation (Winter and Fraaije 2012). Moreover, the recently discovered VAO-type oxidases that harbour a bicovalently bound FAD have shown to be able act on relatively large substrates. This makes the VAO-type oxidases an interesting oxidase family when looking for new oxidative biocatalysts. With the available sequence and structure data, it has become feasible to discover new oxidases by genome mining. In the last decade, we have been successful in obtaining a number of novel flavoprotein oxidases by this approach: alditol oxidase (Heuts et al. 2007b), eugenol oxidase (Jin et al. 2007), putrescine oxidase (Van Hellemond et al. 2008), and chitooligosaccharide oxidase (Heuts et al. 2007). Genome mining has also shown to be effective in identifying thermostable variants of known oxidases, as shown in *L*-aspartate oxidase (Bifulco et al. 2013) or alditol oxidase (Winter et al. 2012). This shows that it is becoming relatively easy to identify and produce new oxidases for biotechnological applications. For fine tuning or improving the enzyme properties, ample methods are available to engineer oxidases. It has been demonstrated that glucose oxidase can be optimised for biosensor and biofuel cells, whereas glycine oxidase has been engineered towards converting glyphosate. We have shown that, by enzyme engineering, it is feasible to introduce peroxidase activity into an oxidase (Winter and Fraaije 2012) and to turn a dehydrogenase into an oxidase (Lefेरink et al. 2009). These examples show that the

number of available flavoprotein oxidases is rapidly expanding, which will fuel the development of oxidase-based biotechnological applications.

Acknowledgements M.W. Fraaije and A. Mattevi received support from the European Union's Seventh Framework Programme (FP7/2007-2013), project Oxygreen (grant agreement no. 212281).

References

- Araki K, Inaba K (2012) Structure, mechanism, and evolution of Ero1 family enzymes. *Antioxid Redox Signal* 16:790–799
- Arjunan P, Umland T, Dyda F, Swaminathan S, Furey W, Sax M, Farrenkopf B, Gao Y, Zhang D, Jordan F (1996) Crystal structure of the thiamin diphosphate-dependent enzyme pyruvate decarboxylase from the yeast *Saccharomyces cerevisiae* at 2.3 Å resolution. *J Biol Chem* 256:590–600
- Bankar SB, Bule MV, Singhal RS, Ananthanaryan L (2009) Glucose oxidase: an overview. *Biotechnol Adv* 27:489–501
- Baron R, Riley C, Chenprakhon P, Thotsaporn K, Winter RT, Alfieri A, Forneris F, Van Berkel WJH, Chaiyen P, Fraaije W, Mattevi A, McCammon JA (2009) Multiple pathways guide oxygen diffusion into flavoenzyme active sites. *PNAS* 106:10603–10608.
- Binzak B, Willard J, Vockley J (1998) Identification of the catalytic residue of human short/branched chain acyl-CoA dehydrogenase by in vitro mutagenesis. *Biochim Biophys Acta* 1382:137–142
- Bifulco D, Pollegioni L, Tessaro D, Servi S, Molla G (2013) A thermostable *L*-aspartate oxidase: a new tool for biotechnological applications. *Appl Microbiol Biotechnol*. doi:10.1007/s00253-013-4688-1
- Biswal BK, Cherney MM, Wang M, Garen C, James MNG (2005) Structure of *Mycobacterium tuberculosis* pyridoxine 5'-phosphate oxidase and its complexes with flavin mononucleotide and pyridoxal 5'-phosphate. *Acta Crystallogr D: Biol Crystallogr* 61:1492–1499
- Bruckner RC, Jorns MS (2009) Spectral and kinetic characterization of intermediates in the aromatization reaction catalysed by nikD, an unusual amino acid oxidase. *Biochemistry* 48:4455–4465
- Cash KJ, Clark HA (2010) Nanosensors and nanomaterials for monitoring glucose in diabetes. *Trends Mol Med* 16:584–593
- Chaiyen P, Fraaije MW, Mattevi A (2012) The enigmatic reaction of flavins with oxygen. *Trends BiochemSci* 37:373–380
- Chen Z, Hassan-Abdulah A, Zhao G, Schuman Jorns M, Mathews FS (2006) Heterotetrameric sarcosine oxidase: structure of a diflavin metalloenzyme at 1.85 Å resolution. *J Mol Biol* 360:1000–1018
- Choi KJ, Noh KM, Kim DE, Ha BH, Kim EE, Yoon MY (2007) Identification of the catalytic subunit of acetohydroxyacid synthase in *Haemophilus influenzae* and its potent inhibitors. *Arch Biochem Biophys* 466:24–30
- Coulombe R, Yue KQ, Ghisla S, Vrieling A (2001) Oxygen access to the active site of cholesterol oxidase through a narrow channel is gated by an Arg–Glu pair. *J Biol Chem* 276:30435–30441
- Cunane LM, Barton JD, Chen Z, Diép Lê KH, Amar D, Lederer F, Mathews FS (2005) Crystal structure analysis of recombinant rat kidney long chain hydroxyl acid oxidase. *Biochem* 44:1521–1531
- Debeurme F, Picciocchi A, Dagher MC, Grunwald D, Beaumel S, Fieschi F, Stasia MJ (2010) Regulation of NADPH oxidase activity in phagocytes relationship between FAD/NADPH binding and oxidase complex assembly. *J Biol Chem* 285:33197–33208
- Decker K, Dai VD (1967) Mechanism and specificity of *L*- and *D*-6-hydroxynicotine oxidase. *Eur J Biochem* 3:132

- Digits JA, Pyun H-J, Coates RM, Casey PJ (2002) Stereospecificity and kinetic mechanism of human prenylcysteine lyase, an usual thioether oxidase. *J Biol Chem* 277:41086–41093
- Di Salvo ML, Contestabile R, Safo MK (2011) Vitamin B6 salvage enzymes: mechanism, structure and regulation. *Biochim Biophys Acta* 1814:1597–1608
- Endo T, Yamano K, Kawano S (2010) Structural basis for the disulfide relay system in the mitochondrial intermembrane space. *Antioxid Redox Sign* 13:1359–1373
- Fan F, Ghanem M, Gadda G (2004) Cloning, sequence analysis, and purification of choline oxidase from *Arthrobacter globiformis*: a bacterial enzyme involved in osmotic stress tolerance. *Arch Biochim Biophys* 421:149–158
- Fass D (2008) The Erv family of sulfhydryl oxidases. *Biochim Biophys Acta* 1783:557–566
- Fetzner S, Steiner RA (2010) Cofactor-independent oxidases and oxygenases. *Appl Microbiol Biotechnol* 86:791–804
- Finn RD, Mistry J, Tate J, Coggill P, Heger A, Pollington JE, Gavin OL, Gunasekaran P, Ceric G, Forslund K, Holm L, Sonnhammer EL, Eddy SR, Bateman A (2010) The Pfam protein families database. *Nucleic Acids Res* 38:D211–222
- Fitzpatrick PF (2010) Oxidation of amines by flavoproteins. *Arch Biochim Biophys* 493:13–25
- Fraaije MW, Veeger C, van Berkel WJH (1995) Substrate specificity of flavin-dependent vanillyl-alcohol oxidase from *Penicillium simplicissimum*. *Eur J Biochem* 234:271–277
- Fraaije MW, van Berkel WJ, Benen JA, Visser J, Mattevi A (1998) A novel oxidoreductase family sharing a conserved FAD-binding domain. *Trends Biochem Sci* 23:206–207
- Fraaije MW, van den Heuvel RHH, van Berkel WJH, Mattevi A (1999) Covalent flavinylation is essential for efficient redox catalysis in vanillyl-alcohol oxidase. *J Biol Chem* 274:35514–35520
- Fraaije MW, Mattevi A (2000) Flavoenzymes: diverse catalysts with recurrent features. *Trends Biochem Sci* 25:126–132
- Furuichi M, Suzuki N, Dhakshnamoorthy B, Minagawa H, Yamagishi R, Watanabe Y, Goto Y, Kaneko H, Yoshida Y, Yagi H, Waga I, Kuamr PKR, Mizuno H (2008) X-ray structure of *Aerococcus viridans* lactate oxidase and its complex with D-lactate at pH 4.5 show an α -hydroxyacid oxidation mechanism. *J Mol Biol* 378:436–446
- Gadda G (2012) Oxygen activation in flavoprotein oxidases: the importance of being positive. *Biochemistry* 51:2662–2669
- Gross E, Kastner DB, Kaiser CA, Fass D (2004) Structure of Ero1p, source of disulphide bonds for oxidative protein folding in the cell. *Cell* 117:601–610
- Guengerich FP (2012) Thematic minireview series: Metals in Biology 2012. *J Biol Chem* 287:13508–13509, and references therein
- Guo PC, Ma YL, Jiang YL, Wang SJ, Bao ZZ, Yu XJ, Chen Y, Zhou CZ (2012) Structure of yeast sulfhydryl oxidase Erv1 reveals electron transfer of the disulfide relay system in the mitochondrial intermembrane space. *J Biol Chem* 287:34961–34969
- Hanson RL, Davis BL, Goldberg SL, Johnston RM, Parker WL, Tully TP, Montana MA, Patel RN (2008) Enzymatic preparation of a D-amino acid from a racemic amino acid or keto acid. *Org Proc Res Dev* 12:1119–1129
- van Hellemond EW, van Dijk M, Heuts DPHM, Janssen DB, Fraaije MW (2008) Discovery and characterization of a putrescine oxidase from *Rhodococcus erythropolis* NCIMB 11540. *Appl Microbiol Biotechnol* 78:455–463
- Heuts DPHM, Janssen DB, Fraaije MW (2007a) A) Changing the substrate specificity of a chitoooligosaccharide oxidase from *Fusarium graminearum* by model-inspired site-directed mutagenesis. *FEBS L* 581:4905–4909
- Heuts DPHM, van Hellemond EW, Janssen DB, Fraaije MW (2007b) B) Discovery, characterization, and kinetic analysis of an alditol oxidase from *Streptomyces coelicolor*. *J Biol Chem* 282:20283–20291
- Heuts DPHM, Scrutton NS, McIntire WS, Fraaije MW (2009) What's in a covalent bond? On the role and formation of covalently bound flavin cofactors. *FEBS J* 276:3405–3427
- Jin J, Mazon H, van den Heuvel RHH, Janssen DB, Fraaije MW (2007) Discovery of a eugenol oxidase from *Rhodococcus* sp. strain RHA1. *FEBS J* 274:2311–2321
- Kasahara K, Miyamoto T, Fujimoto T, Oguri H, Tokiwano T, Oikawa H, Ebizuka Y, Fujii I (2010) Solanapyrone synthase, a possible Diels–Alderase and iterative type I polyketide synthase encoded in a biosynthetic gene cluster from *Alternaria solani*. *Chem Bio Chem* 11:1245–1252
- Kazarinoff MN, McCormick DB (1973) N-(5'-phospho-4'-pyridoxyl) amines as substrates for pyridoxine (pyridoxamine) 5'-phosphate oxidase. *Biochem Biophys Res Commun* 52:440–446
- Kazarinoff MN, McCormick DB (1975) Rabbit liver pyridoxamine (pyridoxine) 5'-phosphate oxidase. Purification and properties. *J Biol Chem* 250:3436–3442
- Kiess M, Hecht HJ, Kalisz HM (1998) Glucose oxidase from *Penicillium amagasakiense*. Primary structure and comparison with the other glucose-methanol-choline (GMC) oxidoreductases. *Eur J Biochem* 252:90–99
- Leferink NGH, Heuts DPHM, Fraaije MW, van Berkel WJH (2008) The growing VAO flavoprotein family. *Arch Biochim Biophys* 474:292–301
- Leferink NGH, Fraaije MW, Joosten HJ, Schaap PJ, Mattevi A, van Berkel WJH (2009) Identification of a gatekeeper residue that prevents dehydrogenases from acting as oxidases. *J Biol Chem* 284:4392–4397
- Leys D, Basran J, Scrutton NS (2003) Channeling and formation of 'active' formaldehyde in dimethylglycine oxidase. *EMBO J* 22:4038–4048
- Li Y, Gao Z, Hou H, Zhang J, Yang H, Dong Y, Tan H (2011) Crystal structure and site-directed mutagenesis of a nitroalkane oxidase from *Streptomyces ansochromogenes*. *Biochem Biophys Res Commun* 405:344–348
- Macheroux P, Kappes B, Ealick SE (2011) Flavogenomics—a genomic and structural view of flavin-dependent proteins. *FEBS J* 278:2625–2634
- Mattevi A, Fraaije MW, Coda A, van Berkel WJH (1997) Crystallization an preliminary x-ray analysis of the flavoenzyme vanillyl-alcohol oxidase from *Penicillium simplicissimum*. *Proteins* 27:601–603
- Mattevi A (2006) To be or not to be an oxidase: challenging the oxygen reactivity of flavoenzymes. *Trends Biochem Sci* 31:276–283
- Muller YA, Schumacher G, Rudolph R, Schultz GE (1994) The refined structure of a stabilized mutant and of wild-type pyruvate oxidase from *Lactobacillus plantarum*. *J Mol Biol* 237:315–335
- Netto CGCM, Toma HE, Andrade LH (2013) Superparamagnetic nanoparticles as versatile carriers and supporting materials for enzymes. *J Mol Catal B Enzym* 85–86:71–92
- Nishino T, Okamoto K, Kawaguchi Y, Hori H, Matsumura T, Eger BT, Pai EF, Nishino T (2005) Mechanism of the conversion of xanthine dehydrogenase to xanthine oxidase. *J Biol Chem* 280:2488–24894
- Nishino T, Okamoto K, Eger BT, Pai EF, Nishino T (2008) Mammalian xanthine oxidoreductases—mechanism of transition from xanthine dehydrogenase to xanthine oxidase. *FEBS J* 275:3278–3289
- Pauff JM, Hemann CF, Junemann N, Leimkuhler S, Hille R (2007) The role of arginine 310 in catalysis and substrate specificity in xanthine dehydrogenase from *Rhodobacter capsulatus*. *J Biol Chem* 282:12785–12790
- Patel RN, Chen Y, Goldberg SL, Hanson RL, Goswami A, Tully TP, Parker WL (2007) PCT Int. Appl WO 2007112299 A2 20071004, CAN147:425649:1120352.

- Patel RN (2011) Biocatalysis: synthesis of key intermediates for development of pharmaceuticals. *ACS Catal* 1:11056–1074
- Pedotti M, Rosini E, Molla G, Moschetti T, Savino C, Vallone PL (2009) Glyphosate resistance by engineering the flavoenzyme glycine oxidase. *J BiolChem* 284:36415–36423
- Pillone MS, Pollegioni L (2002) D-amino acid oxidase as an industrial biocatalyst. *BiocatBiotrans* 20:145–159
- Pollard MG, Travers KJ, Weissman JS (1998) Ero1P: novel and ubiquitous protein with an essential role in oxidative protein folding in the endoplasmic reticulum. *Mol Cell* 1:171–82
- Pollegioni L, Molla G, Sacchi S, Rosini E, Verga R, Pilone MS (2008) Properties and applications of microbial D-amino acid oxidases: current state and perspectives. *Appl Microbiol Biotech* 78:1–16
- Qian Y, Zheng J, Lin Z (2013) Loop engineering of amadoriase II and mutational cooperativity. *Appl Microbiol Biotechnol*. doi:10.1007/s00253-013-4705-4
- Resch V, Schrittwieser JH, Wallner S, Macheroux P, Kroutil W (2011) Biocatalytic oxidative C–C bond formation catalysed by the berberine bridge enzyme: optimal reaction conditions. *Adv Synth Catal* 353:2377–2383
- Resch V, Lechner H, Schrittwieser JH, Wallner S, Gruber K, Macheroux P, Kroutil W (2012) Inverting the regioselectivity of the Berberine Bridge Enzyme by employing customized fluorine-containing substrates. *Chem Eur J* 18:13173–13179
- Ridge PG, Zhang Y, Gladyshev VN (2008) Comparative genomic analyses of copper transporters and cuproproteomes reveal evolutionary dynamics of copper utilization and its link to oxygen. *PLoS One* 3:e1378
- Roth JP, Klinman JP (2003) Catalysis of electron transfer during activation of O₂ by the flavoprotein glucose oxidase. *PNAS* 100:62–67
- Schrittwieser JH, Resch V, Sattler JH, Lienhart WD, Durchschein K, Winkler A, Gruber K, Macheroux P, Kroutil W (2011a) Biocatalytic enantioselective oxidative C–C coupling by aerobic C–H activation. *Angew Chem Int Ed* 50:1068–1071
- Schrittwieser JH, Resch V, Wallner V, Lienhart WD, Sattler JH, Resch J, Macheroux P, Kroutil W (2011b) Biocatalytic organic synthesis of optically pure (S)-scoulerine and berberine and benzyloquinoline alkaloids. *J Org Chem* 76:6703–6714
- Thorpe C, Hooper KL, Raje S, Glynn NM, Burnside J, Turi GK, Coppock DL (2002) Sulfhydryl oxidases: emerging catalysts of protein disulphide bond formation in eukaryotes. *Arch Biochem Biophys* 405:1–12
- Tittmann K, Wille G, Golbik R, Weidner A, Ghisla S, Hübner G (2005) Radical phosphate transfer mechanism for the thiamin diphosphate- and FAD-dependent pyruvate oxidase from *Lactobacillus plantarum*. Kinetic coupling of intercofactor electron transfer with phosphate transfer to acetyl-thiamin diphosphate via transient FAD semiquinone/hydroxyethyl-THDP radical pair. *Biochemistry* 44:13291–13303
- Tokuoka K, Nakajima Y, Hirotsu K, Miyahara I, Nishina Y, Shiga K, Tamaoki H, Setoyama C, Tojo H, Miura R (2006) Three-dimensional structure of rat-liver Acyl-CoA oxidase in complex with a fatty acid: insights into substrate-recognition and reactivity toward molecular oxygen. *J Biochem* 139:789–795
- Tormos JR, Taylor AB, Daubner SC, Hart PJ, Fitzpatrick PF (2010) Identification of a hypothetical protein from *Podospira anserine* as a nitroalkane oxidase. *Biochemistry* 24:5035–5041
- Trickey P, Wagner MA, Schuman Jorns M, Mathews FS (1999) Monomeric sarcosine oxidase: structure of a covalently flavinylated amine oxidizing enzyme. *Structure* 7:331–345
- Turner NJ (2011) Enantioselective oxidation of C–O and C–N bonds using oxidases. *Chem Rev* 111:4073–4087
- Umena Y, Yorita K, Matsuoka K, Kita A, Fukui K, Morimoto Y (2006) The crystal structure of L-lactate oxidase from *Aerococcus viridans* at 2.1 Å resolution reveals the mechanism of strict substrate recognition. *Biochemistry* 350:249–256
- Winkler A, Lyskowski A, Riedl S, Puhl M, Kutchan TM, Macheroux P, Gruber K (2008) A concerted mechanism for berberine bridge enzyme. *Nat Chem Biol* 4:739–741
- Winter RT, Fraaije MW (2012) Applications of flavoprotein oxidases in organic synthesis: novel reactivities that go beyond amine and alcohol oxidation. *Curr Org Chem* 16:2542–2550
- Winter RT, Heuts DPHM, Rijpkema EMA, van Bloois E, Wijma HJ, Fraaije MW (2012) Hot or not? Discovery and characterization of a thermostable alditol oxidase from *Acidothermus cellulolyticus* 11B. *Appl Microbiol Biotech* 95:389–403