

University of Groningen

First Ionization Potentials of Fm, Md, No, and Lr

Sato, Tetsuya K.; Asai, Masato; Borschevsky, Anastasia; Beerwerth, Randolph; Kaneya, Yusuke; Makii, Hiroyuki; Mitsukai, Akina; Nagame, Yuichiro; Osa, Akihiko; Toyoshima, Atsushi

Published in:
Journal of the American Chemical Society

DOI:
[10.1021/jacs.8b09068](https://doi.org/10.1021/jacs.8b09068)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Sato, T. K., Asai, M., Borschevsky, A., Beerwerth, R., Kaneya, Y., Makii, H., ... Stora, T. (2018). First Ionization Potentials of Fm, Md, No, and Lr: Verification of Filling-Up of 5f Electrons and Confirmation of the Actinide Series. *Journal of the American Chemical Society*, 140(44), 14609-14613. <https://doi.org/10.1021/jacs.8b09068>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



First Ionization Potentials of Fm, Md, No, and Lr: Verification of Filling-Up of 5f Electrons and Confirmation of the Actinide Series

Tetsuya K. Sato,^{*,†,Ⓜ} Masato Asai,[†] Anastasia Borschevsky,[‡] Randolph Beerwerth,^{§,⊥} Yusuke Kaneya,^{†,||} Hiroyuki Makii,[†] Akina Mitsukai,^{†,||} Yuichiro Nagame,^{†,||} Akihiko Osa,[†] Atsushi Toyoshima,[†] Kazuaki Tsukada,[†] Minoru Sakama,[#] Shinsaku Takeda,[#] Kazuhiro Ooe,[∇] Daisuke Sato,[∇] Yudai Shigekawa,[⊗] Shin-ichi Ichikawa,[◆] Christoph E. Düllmann,^{¶,▲,×} Jessica Grund,^{¶,▲} Dennis Renisch,^{¶,▲} Jens V. Kratz,[¶] Matthias Schädel,[×] Ephraim Eliav,[□] Uzi Kaldor,[□] Stephan Fritzsche,^{§,⊥} and Thierry Stora[○]

[†]Japan Atomic Energy Agency (JAEA), Tokai, Ibaraki 319-1195, Japan

[‡]The Van Swinderen Institute for Particle Physics and Gravity, University of Groningen, 9700 AB Groningen, The Netherlands

[§]Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität, 07743 Jena, Germany

[⊥]Helmholtz-Institut Jena, 07743 Jena, Germany

^{||}Graduate School of Science and Engineering, Ibaraki University, Mito, Ibaraki 310-8512, Japan

[#]Graduate School of Biomedical Sciences, Tokushima University, Tokushima 770-8503, Japan

[∇]Graduate School of Science and Technology, Niigata University, Niigata 910-2181, Japan

[⊗]Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

[◆]Nishina Center for Accelerator-Based Science, RIKEN, Wako, Saitama 351-0198, Japan

[¶]Institut für Kernchemie, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany

[▲]Helmholtz-Institut Mainz, 55099 Mainz, Germany

[×]GSI Helmholtzzentrum für Schwerionenforschung, 64291 Darmstadt, Germany

[□]School of Chemistry, Tel Aviv University, 69978 Tel Aviv, Israel

[○]ISOLDE, CERN, 1211 Geneva, Switzerland

Supporting Information

ABSTRACT: We report the first ionization potentials (IP₁) of the heavy actinides, fermium (Fm, atomic number Z = 100), mendelevium (Md, Z = 101), nobelium (No, Z = 102), and lawrencium (Lr, Z = 103), determined using a method based on a surface ionization process coupled to an online mass separation technique in an atom-at-a-time regime. The measured IP₁ values agree well with those predicted by state-of-the-art relativistic calculations performed alongside the present measurements. Similar to the well-established behavior for the lanthanides, the IP₁ values of the heavy actinides up to No increase with filling up the 5f orbital, while that of Lr is the lowest among the actinides. These results clearly demonstrate that the 5f orbital is fully filled at No with the [Rn]5f¹⁴7s² configuration and that Lr has a weakly bound electron outside the No core. In analogy to the lanthanide series, the present results unequivocally verify that the actinide series ends with Lr.

Extending the periodic table and classifying newly discovered heavy elements are among the most fundamental and exciting aspects of the chemical sciences. This leads to architect the periodic table and revise its structure

in the heavy element region. The most recent revision of the structure of the periodic table took place in the 1940s when Glenn T. Seaborg introduced the ground-breaking actinide concept,^{1,2} placing a new actinide series below the lanthanides. In this new series, the 5f electron shell is filled in a manner similar to the filling of the 4f electron shell in lanthanides. The actinide concept did not only allow for the immediate discoveries of the elements 95, americium, and 96, curium, but was also instrumental for the discovery of heavier ones. Chemical properties of weighable amounts of nuclear-reactor-produced actinides up to Fm have been extensively studied.³ However, much less is known about the heavier actinides due to stringent limitation on experimental procedures⁴ with increasing atomic number as these heavy elements are available in decreasing quantities of only one atom at a time.^{5,6}

The first ionization potential (IP₁) of an atom is one of the most fundamental chemical and physical quantities of every element. The first measurements of IP₁ of actinides were performed by a surface ionization technique.⁷ Then laser spectroscopy and resonance ionization mass spectroscopy of macroscopically available actinides up to einsteinium have been conducted to measure accurate IP₁ values.^{8–11}

Received: September 7, 2018

Published: October 25, 2018



Table 1. IP_1^* Obtained from I_{eff} and N at Temperature T

element	T (K)	I_{eff} (%)	N	IP_1^* (eV)	$kT \ln(Q_i/Q_0)$ (eV)	IP_1^a (eV)
^{100}Fm	2900 ± 100	1.3 ± 0.4	71 ± 20	6.39 ± 0.13	0.13 ± 0.02	6.52 ± 0.13
^{101}Md	2900 ± 100	1.2 ± 0.3	71 ± 20	6.43 ± 0.13	0.16 ± 0.01	6.59 ± 0.13
^{102}No	2850 ± 80 3000 ± 100	0.54 ± 0.09 0.77 ± 0.10	43 ± 8 34 ± 7	6.44 ± 0.08 $6.45^{+0.09}_{-0.10}$	0.17 ± 0.01 0.18 ± 0.01	6.61 ± 0.08 $6.63^{+0.08}_{-0.10}$
^{103}Lr	2550 ± 50 2850 ± 50	23 ± 5 39 ± 6	35 ± 3 47 ± 3	$5.31^{+0.09}_{-0.06}$ $5.30^{+0.09}_{-0.05}$	$-0.37^{+0.06}_{-0.04}$ $-0.32^{+0.06}_{-0.04}$	$4.99^{+0.10}_{-0.07}$ $4.94^{+0.10}_{-0.07}$

^aThe IP_1^* and the temperature-dependent correction factor, $kT \ln(Q_i/Q_0)$, give IP_1 (see text).

Recently, we reported the successful measurement of IP_1 of Lr in an atom-at-a-time scale experiment using a method based on surface ionization coupled to mass separation and α -particle detection techniques.¹² The result suggested that Lr has the lowest IP_1 value of all actinide elements, although those of other heavy actinides, Fm, Md, and No, have not yet been determined experimentally. According to the systematic variation of the IP_1 values of heavy actinides, an increasing trend is anticipated up to No due to filling electrons up in the 5f orbital.^{13–16} Nobelium is expected to have the highest IP_1 among the actinides due to the closed-shell structure of $[\text{Rn}]5f^{14}7s^2$. Very recently laser resonance ionization spectroscopy of No, using ^{254}No (half-life, $T_{1/2} = 51.2$ s) in one-atom-at-a-time quantities, was performed and the IP_1 has been measured to be 6.62621 ± 0.00005 eV,^{17,18} supporting the scenario of closed 5f and 7s atomic shells in No. However, to unequivocally confirm the filling of the 5f electron shell in the heavy actinides, it is indispensable to experimentally determine the successive IP_1 values from Fm to Lr.

In the present study, we have applied the earlier developed surface-ionization method¹² to determine the IP_1 values of Fm, Md, and No. In addition, IP_1 of Lr has been also measured to improve the accuracy of the previously reported IP_1 .¹² Surface

installed at the JAEA-ISOL (Isotope Separator Online) by the He/CdI₂ gas-jet transport system.²¹ Transported products were injected into the ionization cavity of the ion-source. Metallic tantalum (Ta) was selected as the cavity material in this work. The products were surface-ionized on the hot surface of the Ta cavity kept at a temperature between 2550 and 3000 K. Produced ions are extracted and mass separated in the ISOL. The number of collected ions after the mass-separation was determined by α spectrometry.^{12,20} The I_{eff} value was calculated from a ratio of the number of mass-separated ions to that of directly collected atoms transported by the gas-jet system.²⁰

The α spectra after surface ionization and following mass-separation are shown in Supplement Figures 1–4. The measured I_{eff} values for ^{249}Fm , ^{251}Md , ^{257}No , and ^{256}Lr are listed in Table 1 with the related surface temperature. On the basis of the S-L equation,^{19,23} I_{eff} in a small cavity configuration can be expressed as^{12,24}

$$I_{\text{eff}} = \frac{N \exp\left(\frac{\phi - IP_1^*}{kT}\right)}{1 + N \exp\left(\frac{\phi - IP_1^*}{kT}\right)}, \quad (1)$$

where N is a parameter that depends on the effective number of atom–surface interactions in the cavity, and k is the Boltzmann constant. IP_1^* , the effective IP_1 , is directly related to the IP_1 as^{19,23}

$$IP_1^* = IP_1 - kT \ln\left(\frac{Q_i}{Q_0}\right) \quad (2)$$

where Q_i and Q_0 are the partition functions for the ion and atoms at a given temperature, which can be calculated using excitation energies and statistical weights of their ground and excited states. Thus, IP_1^* can be calculated from the experimentally determined I_{eff} value of the isotope of interest via eq 1. Then, IP_1^* can be converted to IP_1 using eq 2.

To confirm the correlation between I_{eff} and IP_1^* in the present system, I_{eff} values of short-lived lanthanides, an alkali metal, and a chromium isotope were measured. The short-lived isotopes, $^{143\text{m}}\text{Sm}$, $^{142\text{m},143}\text{Eu}$, $^{148\text{m}}\text{Tb}$, $^{153,154}\text{Ho}$, ^{157}Er , ^{162}Tm , ^{165}Yb , ^{168}Lu , ^{80}Rb , and ^{49}Cr were employed. Figure 1 shows the typical plot of the measured I_{eff} values vs IP_1^* of these elements at $T = 3000$ K. The IP_1^* values of the above elements were calculated via eq 2 using their known IP_1 values compiled in the National Institute of Standard and Technology (NIST) atomic spectra database (ASD).²⁵ Low-lying excited states for the calculation of Q_i and Q_0 were also taken from NIST ASD. Values of the parameter N were obtained by a best-fit with eq 1 to the measured I_{eff} values for the isotopes; summarized with the other quantities in Table 1. The determination of $IP_1^* = 6.45$ eV for No from $I_{\text{eff}} = 0.77\%$ at $T = 3000$ K is depicted in Figure 1. The I_{eff} vs IP_1^* plot at 2900 K for Md and Fm is shown in Supplement Figure 5.




Figure 1. Ionization efficiency (I_{eff}) of various short-lived isotopes as a function of the effective IP_1 , IP_1^* , at 3000 K. The red-dashed curve is obtained by fitting eq 1 to the experimental data.

ionization process takes place on a solid surface kept at a high temperature and can be described by the Saha–Langmuir (S-L) equation.¹⁹ The ionization efficiency (I_{eff}) depends on the work function of the ionizing material, ϕ (eV), the temperature of the material surface, T (K), and IP_1 of the element. The detailed experimental setup and the analytical method used in this work have been described in our previous papers.^{12,20,21}

Short-lived isotopes ^{249}Fm ($T_{1/2} = 2.6$ min), ^{251}Md ($T_{1/2} = 4.27$ min), ^{257}No ($T_{1/2} = 24.5$ s), and ^{256}Lr ($T_{1/2} = 27$ s) were produced in nuclear fusion reactions (Supplement Table 1). The produced atoms, recoiling from the target, were transported via a Teflon capillary^{20–22} to a surface ion-source

Table 2. Experimental and Theoretical IP₁ Values

ref	method	IP ₁ (eV)			
		Fm	Md	No	Lr
		<i>Theoretical</i>			
Sugar ¹³	semiempirical	6.50	6.58	6.65	
Rajnak and Shore ¹⁴	semiempirical	6.46	6.57	6.67	
Liu et al. ¹⁵	QRPP-CASSCF+ACPF ^d	6.26	6.10	6.14	5.28
Cao et al. ³¹	RPP-CASSCF+ACPF ^b	6.13	6.23	6.27	4.79
Borschevsky et al. ^{12,28}	IHFSCC			6.632	4.963(15)
				ref 28	ref 12 ^d
Pantazis and Neese ¹⁶	DKH2-B3LYP ^c	6.45	6.54	6.64	4.56
Dzuba et al. ³²	CI+SD ^d			6.743	4.9
present work	CCSD(T)	6.469	6.557	6.638	
	MCDF	6.22			
		<i>Experimental</i>			
literature		<6.76		6.62621(5)	4.96 ^{+0.08} _{-0.07}
		ref 33		ref 18	ref 12
present work		6.52 ± 0.13	6.59 ± 0.13	6.62 ^{+0.06} _{-0.07}	4.96 ^{+0.05} _{-0.04}

^aQuasirelativistic ab initio pseudopotential (QRPP) complete active space self-consistent field (CASSCF) calculations combined with averaged coupled-pair functional (ACPF) and corrected for spin-orbit coupling. ^bRelativistic ab initio pseudopotential (RPP) CASSCF calculations combined with ACPF and corrected for spin-orbit coupling. ^cSecond order Douglas-Kroll-Hess approach combined with density functional theory (B3LYP functional). ^dRelativistic configuration interaction (CI) combined with the linearized single-double coupled cluster method.




Figure 2. Variation of the experimental IP₁ values of heavy actinides and heavy lanthanides with atomic numbers. Closed circles indicate the values obtained in the present work.

To calculate the IP₁ values of Fm, Md, No, and Lr from their IP₁* values, excitation energies and statistical weights of the low-lying states of each atom and ion are required. As no experimental data on excited states in the heavy actinides are available, we calculated these values using relativistic computational methods. The intermediate-Hamiltonian Fock space coupled cluster (IHFSCC) method²⁶ was applied to calculations of the atomic and ionic states of Md and No (for some of the levels the single reference coupled cluster with single, double, and perturbative triple excitations (CCSD(T)) was employed), while the Multi-Configuration Dirac-Fock (MCDF) method, as implemented in the Graps2k code,²⁷ was used for the Fm atom and its ion. Excited states of No and Lr were taken from refs 28, 29, where they were also calculated within the IHFSCC approach. The methods used here were also applied to the lower excitation energies of the lanthanide

homologues of the elements of interest (Er, Tm, Yb, and Lu). For the lighter elements, we can compare our results to the available experimental data, thus assessing the accuracy of our calculations and of our predictions for Fm through Lr. The obtained values are compiled in the Supplement Table 2. Although several low-lying states were found in Fm⁺, only one state should be considered for the Fm atom (Supporting Information). There are no excited states in the range of interest for Md, while one state is present for Md⁺. In No and No⁺, only the ground states are expected to contribute. The errors in the energy of the excited states were evaluated from relative errors of the calculated values compared to the experimental transition energies of the respective lanthanide homologues. The $kT \ln(Q_i/Q_0)$ values are presented in Table 1. For the case of Lr, the values were obtained in the same manner in ref 12. The IP₁ values of Fm, Md, and No are determined to be 6.52 ± 0.13, 6.59 ± 0.13, and 6.62^{+0.06}_{-0.07} eV, respectively, where IP₁ of No was obtained by taking a weighted average of the IP₁ values listed in Table 1. A more accurate IP₁ of Lr of 4.96^{+0.05}_{-0.04} eV was determined by also taking a weighted average of our previous¹² and present values. Errors in IP₁ mainly come from counting statistics, surface temperature, and fitting procedure with eq 1.

In parallel to the measurements, we calculated the IP₁ values of Fm, Md, and No within the relativistic CCSD(T) approach, corrected for the Breit term and the higher order quantum electrodynamic (QED) corrections, using a similar scheme to that employed in ref 30.

The experimental and theoretical IP₁ values obtained in the present work are summarized in Table 2 together with earlier theoretical predictions^{12–16,28,31,32} and measurements.^{12,18,33} The present experimental values for Md and No agree with the semiempirical values^{13,14} as well as with the more recent relativistic calculations²⁸ for No and the DKH2-B3LYP calculations¹⁶ for both atoms. Our result on IP₁ of No also agrees with the recent value from laser-spectroscopic measurements,¹⁸ thus providing independent validation to our experimental method. The calculated IP₁ of Er, the homologue

of Fm, is 5.94 eV as obtained from the present MCDF calculation, that is significantly lower than the experimental value of 6.11 eV. Therefore, the MCDF prediction probably also underestimates the IP_1 value of Fm. The CCSD(T) + Breit + QED calculations of IP_1 agree well with the measurements for all elements investigated here.

The variation of the IP_1 values of the heavy actinides with atomic number in comparison with those of the heavy lanthanides is shown in Figure 2. As expected from the prediction,^{13–16} the IP_1 values increase up to No via Fm and Md with filling of the 5f orbital in analogy to the heavy lanthanides. We take this as an indication that the 5f orbital is fully filled at No. The lowest IP_1 value of Lr is confirmed; the ground-state electronic configuration of the Lr atom has closed 5f¹⁴ and 7s² shells with an additional weakly bound electron in the valence orbital. The results unambiguously confirm that the actinide series end with Lr.³⁴

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.8b09068.

Experimental details of the nuclear reactions for the production of the short-lived actinide isotopes, ²⁴⁹Fm, ²⁵¹Md, ²⁵⁷No, and ²⁵⁶Lr and for production of various isotopes used to determine a relationship between IP_1^* and I_{eff} in the present system; alpha spectra of ²⁴⁹Fm, ²⁵¹Md, ²⁵⁷No, and ²⁵⁶Lr measured after mass-separation; ionization efficiencies of various short-lived isotopes as a function of the IP_1^* at 2900 K, which was used for IP_1 calculations of Md and Fm; and summary of computed low-lying level energies for the Fm, Md, and No atoms and ions together with those of respective lanthanide homologues, Er, Tm, and Yb (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*sato.tetsuya@jaea.go.jp

ORCID

Tetsuya K. Sato: 0000-0002-5490-9178

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors would like to thank the JAEA tandem accelerator crew for supplying intense and stable beams for the experiments. The ²⁴⁹Cf was produced in the form of ²⁴⁹Bk through the former Transplutonium Element Production Program at Oak Ridge National Laboratory (ORNL) under the auspices of the Director, Office of Science, Office of Basic Energy Sciences, Chemical Sciences, Geosciences, and Biosciences Division of US Department of Energy. It was made available by H. Nitsche of University of California, Berkeley, USA, and we acknowledge the LBNL Nuclear Science Division's R. F. Fairchild II, N. E. Reeves, J. A. Van Wart, and the Radiation Protection Group of the Environmental Health and Safety Division for their support with the preparation and execution of the ²⁴⁹Cf shipment to Germany. Financial support by the Helmholtz-Institut Mainz is gratefully acknowledged. This work has been partly supported by the Grant-in-Aid for Scientific Research (A) No. 16H02130, (B)

No. 26288028, and (C) No. 26390119 of the Ministry of Education, Science, Sports and Culture (MEXT).

■ REFERENCES

- (1) Seaborg, G. T. The Chemical and Radioactive Properties of the Heavy Elements. *Chem. Eng. News* **1945**, *23*, 2190.
- (2) Seaborg, G. T. The Transuranium Elements. *Science* **1946**, *104*, 379.
- (3) *The Chemistry of the Actinide and Transactinide Elements*, 3rd ed.; Morss, L. R., Edelstein, N. M., Fuger, J., Katz, J. J., Eds.; Springer: Dordrecht, 2006.
- (4) Silva, R. In *The Chemistry of the Actinide and Transactinide Elements*, 3rd ed.; Morss, L. R.; Edelstein, N. M., Fuger, J., Katz, J. J., Eds.; Springer: Dordrecht, 2006; Vol. 3, Chapter 13, pp 1621–1651.
- (5) Hoffman, D. C.; Lee, D. N. Chemistry of the Heaviest Elements—One Atom at a Time. *J. Chem. Educ.* **1999**, *76*, 331.
- (6) Türlér, A.; Pershina, V. Advances in the Production and Chemistry of the Heaviest Elements. *Chem. Rev.* **2013**, *113*, 1237.
- (7) (a) Hertel, G. R. Surface Ionization. II. The First Ionization Potential of Uranium. *J. Chem. Phys.* **1967**, *47*, 335. (b) Smith, D. H.; Hertel, G. R. First Ionization Potentials of Th, Np, and Pu by Surface Ionization. *J. Chem. Phys.* **1969**, *51*, 3105.
- (8) Worden, E. F.; Blaise, J.; Fred, M.; Trautmann, N.; Wyart, J.-F. In *The Chemistry of the Actinide and Transactinide Elements*, 3rd ed.; Morss, L. R., Edelstein, N. M., Fuger, J., Katz, J. J., Eds.; Springer: Dordrecht, 2006; Vol. 3, Chapter 16, pp 1836–1892.
- (9) Köhler, S.; Deisenberger, R.; Eberhardt, K.; Erdmann, N.; Herrmann, G.; Huber, G.; Kratz, J. V.; Nunnemann, M.; Passler, G.; Rao, P. M.; Riegel, J.; Trautmann, N.; Wendt, K. Determination of the first ionization potential of actinide elements by resonance ionization mass spectroscopy. *Spectrochim. Acta, Part B* **1997**, *52*, 717.
- (10) Peterson, J. R.; Erdmann, N.; Nunnemann, M.; Eberhardt, K.; Huber, G.; Kratz, J. V.; Passler, G.; Stetzer, O.; Thörle, P.; Trautmann, N.; Waldek, A. Determination of the first ionization potential of einsteinium by resonance ionization mass spectroscopy (RIMS). *J. Alloys Compd.* **1998**, *271–273*, 876.
- (11) Wendt, K.; Gottwald, T.; Mattolat, C.; Raeder, S. Ionization potentials of the lanthanides and actinides – towards atomic spectroscopy of super-heavy elements. *Hyperfine Interact.* **2014**, *227*, 55.
- (12) Sato, T. K.; Asai, M.; Borschevsky, A.; Stora, T.; Sato, N.; Kaneya, Y.; Tsukada, K.; Düllmann, Ch. E.; Eberhardt, K.; Eliav, E.; Ichikawa, S.; Kaldor, U.; Kratz, J. V.; Miyashita, S.; Nagame, Y.; Ooe, K.; Osa, A.; Renisch, D.; Runke, J.; Schädel, M.; Thörle-Pospiech, P.; Toyoshima, A.; Trautmann, N. Measurement of the first ionization potential of lawrencium, element 103. *Nature* **2015**, *520*, 209.
- (13) (a) Sugar, J. Ionization energies of the neutral actinides. *J. Chem. Phys.* **1973**, *59*, 788. (b) Sugar, J. Revised ionization energies of the neutral actinides. *J. Chem. Phys.* **1974**, *60*, 4103.
- (14) Rajnak, K.; Shore, B. W. Regularities in s-electron binding energies in $f^N s^M$ configurations. *J. Opt. Soc. Am.* **1978**, *68*, 360.
- (15) Liu, W.; Küchle, W.; Dolg, M. Ab initio pseudopotential and density-functional all-electron study of ionization and excitation energies of actinide atoms. *Phys. Rev. A: At., Mol., Opt. Phys.* **1998**, *58*, 1103.
- (16) Pantazis, D. A.; Neese, F. All-Electron Scalar Relativistic Basis Sets for the Actinides. *J. Chem. Theory Comput.* **2011**, *7*, 677.
- (17) Laatiaoui, M.; Lauth, W.; Backe, H.; Block, M.; Ackermann, D.; Cheal, B.; Chhetri, P.; Düllmann, Ch. E.; Van Duppen, P.; Even, J.; Ferrer, R.; Giacoppo, F.; Götz, S.; Heßberger, F. P.; Huyse, M.; Kaleja, O.; Khuyagbaatar, J.; Kunz, P.; Lautenschläger, F.; Mistry, A. K.; Raeder, S.; Ramirez, E. M.; Walther, T.; Wraith, C.; Yakushev, A. Atom-at-a-time laser resonance ionization spectroscopy of nobelium. *Nature* **2016**, *538*, 495.
- (18) Chhetri, P.; Ackermann, D.; Backe, H.; Block, M.; Cheal, B.; Droese, C.; Düllmann, Ch. E.; Even, J.; Ferrer, R.; Giacoppo, F.; Götz, S.; Heßberger, F. P.; Huyse, M.; Kaleja, O.; Khuyagbaatar, J.; Kunz, P.; Laatiaoui, M.; Lautenschläger, F.; Lauth, W.; Lecesne, N.; Lens, L.; Minaya Ramirez, E.; Mistry, A. K.; Raeder, S.; Van Duppen, P.;

Walther, Th.; Yakushev, A.; Zhang, Z. Precision Measurement of the First Ionization Potential of Nobelium. *Phys. Rev. Lett.* **2018**, *120*, 263003.

(19) Zandberg, É. Ya.; Ionov, N. I. Surface ionization. *Sov. Phys. Usp.* **1959**, *2*, 255.

(20) Sato, T. K.; Sato, N.; Asai, M.; Tsukada, K.; Toyoshima, A.; Ooe, K.; Miyashita, S.; Schädel, M.; Kaneya, Y.; Nagame, Y.; Osa, A.; Ichikawa, S.; Stora, T.; Kratz, J. V. First successful ionization of Lr ($Z = 103$) by a surface-ionization technique. *Rev. Sci. Instrum.* **2013**, *84*, 023304.

(21) Sato, T. K.; Asai, M.; Sato, N.; Tsukada, K.; Toyoshima, A.; Ooe, K.; Miyashita, S.; Kaneya, Y.; Osa, A.; Schadel, M.; Nagame, Y.; Ichikawa, S.; Stora, T.; Kratz, J. V. Development of a He/CdI₂ gas-jet system coupled to a surface-ionization type ion-source in JAEA-ISOL: towards determination of the first ionization potential of Lr ($Z = 103$). *J. Radioanal. Nucl. Chem.* **2015**, *303*, 1253.

(22) Ichikawa, S.; Asai, M.; Tsukada, K.; Osa, A.; Ikuta, T.; Shinohara, N.; Iimura, H.; Nagame, Y.; Hatsukawa, Y.; Nishinaka, I.; Kawade, K.; Yamamoto, H.; Shibata, M.; Kojima, Y. Mass separation of neutron-rich isotopes using a gas-jet coupled thermal ion source. *Nucl. Instrum. Methods Phys. Res., Sect. A* **1996**, *374*, 330.

(23) Stora, T. Radioactive Ion Sources, Report No. CERN-2013-007, p 331; <http://cds.cern.ch/record/1693046>.

(24) Kirchner, R. On the thermoionization in hot cavities. *Nucl. Instrum. Methods Phys. Res., Sect. A* **1990**, *292*, 203.

(25) Kramida, A.; Ralchenko, Yu.; Reader, J., NIST ASD Team; *NIST Atomic Spectra Database*, version 5.5.2; National Institute of Standards and Technology: Gaithersburg, MD; <https://physics.nist.gov/asd> [accessed Jan 30, 2018].

(26) Eliav, E.; Vilkas, M. J.; Ishikawa, Y.; Kaldor, U. Extrapolated intermediate Hamiltonian coupled-cluster approach: Theory and pilot application to electron affinities of alkali atoms. *J. Chem. Phys.* **2005**, *122*, 224113.

(27) Jönsson, P.; Bieroń, J.; Fischer, C. F.; Grant, I.; Gaigalas, G. Newversion: Grasp2K Relativistic atomics structure package. *Comput. Phys. Commun.* **2013**, *184*, 2197.

(28) Borschevsky, A.; Eliav, E.; Vilkas, M. J.; Ishikawa, Y.; Kaldor, U. Predicted spectrum of atomic nobelium. *Phys. Rev. A: At., Mol., Opt. Phys.* **2007**, *75*, 042514.

(29) Borschevsky, A.; Eliav, E.; Vilkas, M. J.; Ishikawa, Y.; Kaldor, U. Transition energies of atomic lawrencium. *Eur. Phys. J. D* **2007**, *45*, 115.

(30) Borschevsky, A.; Paštka, L. F.; Pershina, V.; Eliav, E.; Kaldor, U. Ionization potentials and electron affinities of the superheavy elements 115–117 and their sixth-row homologues Bi, Po, and At. *Phys. Rev. A: At., Mol., Opt. Phys.* **2015**, *91*, 020501.

(31) Cao, X.; Dolg, M.; Stoll, H. Valence basis sets for relativistic energy-consistent small-core actinide pseudopotentials. *J. Chem. Phys.* **2003**, *118*, 487.

(32) Dzuba, V. A.; Safronova, M. S.; Safronova, U. I. Atomic properties of superheavy elements No, Lr, and Rf. *Phys. Rev. A: At., Mol., Opt. Phys.* **2014**, *90*, 012504.

(33) Sewtz, M.; Backe, H.; Dretzke, A.; Kube, G.; Lauth, W.; Schwamb, P.; Eberhardt, K.; Gruning, C.; Thorle, P.; Trautmann, N.; Lassen, J.; Passler, G.; Dong, C. Z.; Fritzsche, S.; Kunz, P.; Haire, R. G. First Observation of Atomic Levels for the Element Fermium ($Z = 100$). *Phys. Rev. Lett.* **2003**, *90*, 163002.

(34) A debate has emerged over the places of the last lanthanide Lu and the last actinide Lr in the periodic table: Jensen, W. B. The positions of lanthanum (actinium) and lutetium (lawrencium) in the periodic table: an update. *Found. Chem.* **2015**, *17*, 23. An IUPAC task group has been formed to make a recommendation regarding the membership of group 3 of the periodic table, Project No. 2015-039-2-200: Scerri, E. Which Elements Belong to Group 3 of the Periodic Table. *Chem. Int.* **2016**, *38*, 22.