

University of Groningen

The Authors' Reply to Letter to the Editor, Re

Matton, Alix P M; de Meijer, Vincent E; Lisman, Ton; Porte, Robert J

Published in:
 Transplantation

DOI:
[10.1097/TP.0000000000002863](https://doi.org/10.1097/TP.0000000000002863)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Final author's version (accepted by publisher, after peer review)

Publication date:
 2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Matton, A. P. M., de Meijer, V. E., Lisman, T., & Porte, R. J. (2020). The Authors' Reply to Letter to the Editor, Re: Biliary Bicarbonate, pH, and Glucose Are Suitable Biomarkers of Biliary Viability During Ex Situ Normothermic Machine Perfusion of Human Donor Livers. *Transplantation*, 104(1), E41-E41. <https://doi.org/10.1097/TP.0000000000002863>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

OPEN

Transplantation Publish Ahead of Print

DOI: 10.1097/TP.0000000000002863

The Authors' Reply to Letter to the Editor,

**Re: Biliary Bicarbonate, pH and Glucose Are Suitable Biomarkers of Biliary Viability
During Ex Situ Normothermic Machine Perfusion of Human Donor Livers**

Alix P.M. Matton, MD^{1,2}, Vincent E. de Meijer, MD, PhD¹, Ton Lisman, PhD², Robert J. Porte, MD, PhD¹

¹ Section of Hepatobiliary Surgery and Liver Transplantation, Department of Surgery,

² Surgical Research Laboratory, Department of Surgery, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands.

*Re: Biliary Bicarbonate, pH and Glucose Are Suitable Biomarkers of Biliary Viability
During Ex Situ Normothermic Machine Perfusion of Human Donor Livers*

Raphael P.H. Meier, Shareef Syed, John P. Roberts

Transplant Surgery, University of California San Francisco, San Francisco, CA, USA

Corresponding author: Robert J. Porte, MD, PhD, Department of Surgery, Division of HPB Surgery and Liver Transplantation, University Medical Center Groningen, P.O. Box 30.001 9700 RB Groningen, The Netherlands. E-mail: r.j.porte@umcg.nl, Telephone: +31 50 361 2896, Fax: +31 50 361 4873

AUTHORSHIP

Writing of the first version of the reply: APMM, VEdM, RP

Critically revised the manuscript: APMM, VEdM, TL, RP

DISCLOSURES: The authors declare no conflicts of interest.

FUNDING: This study was not financially supported by any third party.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

ACCEPTED

To the editor:

We thank Meier and colleagues for their interest in our recent article.^{1,2} The authors comment that in our previous paper the histological parameters (i.e., the presence of stroma necrosis, injury of the extramural peribiliary glands and injury of the perivascular plexus) in pretransplant bile duct biopsies were only individually correlated with the post-transplant development of nonanastomotic strictures.³ In the present work, we assessed the degree of bile duct injury per histological item and combined the scores into a composite outcome to separate the group into livers with a low and high degree of bile duct injury. The identified biomarkers of bile duct injury (i.e., biliary bicarbonate, biliary pH, biliary glucose, the bile/perfusate glucose ratio and biliary LDH) were subsequently used and validated as cholangiocellular viability criteria, in addition to hepatocellular viability criteria, in a subset of patients from a prospective clinical trial (Figure 5 of our manuscript).² By using the identified biomarkers from the present preclinical work, we were able to safely select and transplant previously nationwide declined donor livers. The current follow-up time is 16.5 (interquartile range 15.8 – 18.4) months with a 100% patient and graft survival. Histology of bile duct biopsies obtained in this study will provide a definite answer to the predictive ability of the combined bile duct injury score.

The reasons why the research livers were declined for transplantation have been published previously.⁴ We would like to clarify that the intention of this preclinical study was to investigate cholangiocellular viability criteria to assess livers that were declined for transplantation, to potentially allow safe transplantation. The characteristics of the livers used in the present preclinical study were, therefore, very similar to the cohort of declined livers assessed in the prospective clinical trial.⁵

The author's call for a large randomized controlled trial using a combination of biliary viability criteria and bile duct histology on frozen sections as a preimplantation diagnostic tool compared to standard of care seems unfounded. Histological quality of frozen sections from bile duct biopsies is inferior to paraffin-embedded sections.

ACCEPTED

References

1. Meier RPH, Syed S, Roberts JP. Re: Biliary Bicarbonate, pH and Glucose Are Suitable Biomarkers of Biliary Viability During Ex Situ Normothermic Machine Perfusion of Human Donor Livers. IN PRESS.
2. Matton APM, de Vries Y, Burlage LC, et al. Biliary Bicarbonate, pH and Glucose Are Suitable Biomarkers of Biliary Viability During Ex Situ Normothermic Machine Perfusion of Human Donor Livers. *Transplantation*. 2018.
3. op den Dries S, Westerkamp AC, Karimian N, et al. Injury to peribiliary glands and vascular plexus before liver transplantation predicts formation of non-anastomotic biliary strictures. *J Hepatol*. 2014;60(6):1172-1179.
4. Matton APM, Burlage LC, van Rijn R, et al. Normothermic machine perfusion of donor livers without the need for human blood products. *Liver Transpl*. 2018;24(4):528-538.
5. de Vries Y, Matton APM, Nijsten MWN, et al. Pretransplant sequential hypo- and normothermic machine perfusion of suboptimal livers donated after circulatory death using a hemoglobin-based oxygen carrier perfusion solution. *Am J Transplant*. 2019;19(4):1202-1211.